

EQUIVARIANT CW-COMPLEXES AND THE ORBIT CATEGORY

IAN HAMBLETON, SEMRA PAMUK, AND ERGÜN YALÇIN

ABSTRACT. We give a general framework for studying G -CW complexes via the orbit category. As an application we show that the symmetric group $G = S_5$ admits a finite G -CW complex X homotopy equivalent to a sphere, with cyclic isotropy subgroups.

1. INTRODUCTION

A good algebraic setting for studying actions of a group G with isotropy in a given family of subgroups \mathcal{F} is provided by the category of R -modules over the orbit category $\Gamma_G = \text{Or}_{\mathcal{F}} G$, where R is a commutative ring with unit. This theory was established by Bredon [6], tom Dieck [10] and Lück [20], and further developed by many authors (see, for example, Jackowski-McClure-Oliver [17, §5], Brady-Leary-Nucinkis [5], Symonds [34], [35], Grodal [14], Grodal-Smith [15]). In particular, the category of $R\Gamma_G$ -modules is an abelian category with Hom and tensor product, and has enough projectives for standard homological algebra.

In this paper, we will study finite group actions on spheres with non-trivial isotropy, generalizing the approach of Swan [32] to the spherical space form problem through periodic projective resolutions. A finite group is said to have *rank* k if k is the largest integer such that G has an elementary abelian subgroup $C_p \times \cdots \times C_p$ of rank k for some prime p . A rank 1 group G has periodic cohomology, and Swan showed that this was a necessary and sufficient condition for the existence of a finite free G -CW complex X , homotopy equivalent to a sphere.

The work of Adem-Smith [1] concerning free actions on products of spheres led to the following open problem:

Question. If G is a rank 2 finite group, does there exist a finite G -CW complex $X \simeq S^n$ with rank 1 isotropy ?

If G is a finite p -group of rank 2, then there exist orthogonal linear representations V so that $S(V)$ has rank 1 isotropy (see [12]). If G is not of prime power order, representation spheres with rank 1 isotropy do not exist in general: a necessary condition is that G has a p -effective character for each prime p dividing $|G|$ (see [18, Thm. 47]). In [18, Prop. 48] it is claimed that this condition is also sufficient for an affirmative answer to the G -CW question above, but the discussion on [18, p. 831] does not provide a construction for X .

Date: April 14, 2010 (final version).

Research partially supported by NSERC Discovery Grant A4000. The third author is partially supported by TÜBİTAK-BDP and TÜBA-GEBİP/2005-16.

Our main result concerns the first non-trivial case: the permutation group $G = S_5$ of order 120, which has rank 2 but no linear action with rank 1 isotropy on any sphere, although it does admit p -effective characters for $p = 2, 3, 5$.

Theorem A. *The permutation group $G = S_5$ admits a finite G -CW complex $X \simeq S^n$, such that $X^H \neq \emptyset$ implies that H is a rank 1 subgroup of 2-power order.*

Remark 1.1. It is an interesting problem for future work to decide if the group $G = S_5$ can act *smoothly* on S^n with rank 1 isotropy.

In order to prove this result we develop further techniques over the orbit category, which may have some independent interest. A well-known theorem of Rim [29] shows that a module M over the group ring $\mathbf{Z}G$ is projective if and only if its restriction $\text{Res}_P^G M$ to any p -Sylow subgroup is projective. Over the orbit category we have a similar statement localized at p (see Theorem 3.9).

Theorem B. *Let G be a finite group and let $R = \mathbf{Z}_{(p)}$. Then an $R\Gamma_G$ -module M has a finite projective resolution with respect to a family of p -subgroups if and only if its restriction $\text{Res}_P^G M$ has a finite projective resolution over any p -Sylow subgroup $P \leq G$.*

Remark 1.2. For modules over the group ring RG , those having finite projective resolutions are already projective. Over the orbit category, these two properties are distinct.

Another useful feature of homological algebra over group rings is the detection of group cohomology by restriction to the p -Sylow subgroups. Here is an important concept in group cohomology (see for example [33]).

Definition 1.3. For a given prime p , we say that a subgroup $H \subseteq G$ *controls p -fusion* provided that

- (i) $p \nmid |G/H|$, and
- (ii) whenever $Q \subseteq H$ is a p -subgroup, and there exists $g \in G$ such that $Q^g := g^{-1}Qg \subseteq H$, then $g = ch$ where $c \in C_G(Q)$ and $h \in H$.

One reason for the importance of this definition is the fact that the restriction map

$$H^*(G; \mathbb{F}_p) \rightarrow H^*(H; \mathbb{F}_p)$$

is an isomorphism if and only if H controls p -fusion in G (see [25], [33]). We have the following generalization (see Theorem 5.1) for functors of cohomological type over the orbit category (with respect to any family \mathcal{F}).

Theorem C. *Let G be a finite group, $R = \mathbf{Z}_{(p)}$, and $H \leq G$ a subgroup which controls p -fusion in G . If M is an $R\Gamma_G$ -module and N is a cohomological Mackey functor, then the restriction map*

$$\text{Res}_H^G: \text{Ext}_{R\Gamma_G}^n(M, N) \rightarrow \text{Ext}_{R\Gamma_H}^n(\text{Res}_H^G M, \text{Res}_H^G N)$$

is an isomorphism for $n \geq 0$, provided that the centralizer $C_G(Q)$ of any p -subgroup $Q \leq H$, with $Q \in \mathcal{F}$, acts trivially on $M(Q)$ and $N(Q)$.

The construction of the G -CW complex X for $G = S_5$ and the proof of Theorem A is carried out in Section 9. We first construct finite projective chain complexes $\mathbf{C}^{(p)}$ over the orbit categories $R\Gamma_G$, with $R = \mathbf{Z}_{(p)}$, separately for the prime $p = 2, 3, 5$ dividing $|G|$. In each case, the isotropy family \mathcal{F} consists of the rank 1 subgroups of 2-power order in G .

The chain complexes $\mathbf{C}^{(p)}$ all have the same *dimension function* (see Definition 8.2). We prescribe a non-negative function $\underline{n}: \mathcal{F} \rightarrow \mathbf{Z}$, with the property that $\underline{n}(K) \leq \underline{n}(H)$ whenever H is conjugate to a subgroup of K . Then, by construction, each complex $\mathbf{C}^{(p)}$ has the R -homology of an \underline{n} -sphere: for each $K \in \mathcal{F}$, the complexes $\mathbf{C}^{(p)}(K)$ have homology $H_i = R$ only in two dimensions $i = 0$ and $i = \underline{n}(K)$. In other words, the complexes $\mathbf{C}^{(p)}$ are algebraic versions of tom Dieck's *homotopy representations* [10, II.10].

In the case $p = 2$, we start with the group $H = S_4$ acting by orthogonal rotations on the 2-sphere. A regular H -equivariant triangulation of an inscribed cube or octahedron gives a finite projective chain complex over $R\Gamma_H$. Then we use Proposition 6.4, a chain complex version of Theorem C, to lift it to a finite projective complex over $R\Gamma_G$. For $p = 3$ and $p = 5$, the p -rank of S_5 is 1, and there exists a periodic complex over the group ring RG (see Swan [32, Theorem B]). We start with a periodic complex over RG and add chain complexes to this complex, for every nontrivial subgroup $K \in \mathcal{F}$, to obtain the required complex $\mathbf{C}^{(p)}$ over $R\Gamma_G$.

We use the theory of algebraic Postnikov sections by Dold [11] to glue the complexes together to form a finite projective $\mathbf{Z}\Gamma_G$ chain complex (see Section 6). We complete the chain complex construction by varying the finiteness obstruction to obtain a complex of free $\mathbf{Z}\Gamma_G$ -modules, and then we prove a realization theorem (see Section 8) to construct the required G -CW complex $X \simeq S^n$.

Throughout the paper, a *family* of subgroups will always mean a collection of subgroups which is closed under conjugation and taking subgroups. Also, unless otherwise stated, all modules are *finitely generated*.

Acknowledgement. The authors would like to thank the referee for many valuable criticisms and suggestions. The third author would also like to thank McMaster University for the support provided by a H. L. Hooker Visiting Fellowship, and the Department of Mathematics & Statistics at McMaster for its hospitality while this paper was written.

2. MODULES OVER SMALL CATEGORIES

Our main source for the material in this section is Lück [20, §9, §17] (see also [10, §I.10, §I.11]). We include it here for the convenience of the reader.

Let R be a commutative ring. We denote the category of R -modules by $R\text{-Mod}$. For a small category Γ (i.e., the objects $\text{Ob}(\Gamma)$ of Γ form a set), the category of right $R\Gamma$ -modules is defined as the category of contravariant functors $\Gamma \rightarrow R\text{-Mod}$, where the objects are functors $M(-): \Gamma \rightarrow R\text{-Mod}$ and morphisms are natural transformations. Similarly, we define the category of left $R\Gamma$ -modules as the category of covariant functors $N(-): \Gamma \rightarrow R\text{-Mod}$. We denote the category of right $R\Gamma$ -modules by $\text{Mod-}R\Gamma$ and the category of left $R\Gamma$ -modules by $R\Gamma\text{-Mod}$.

The category of covariant or contravariant functors from a small category to an abelian category has the structure of abelian category which is object-wise induced from the

abelian category structure on abelian groups (see [23, Chapter 9, Prop. 3.1]). Hence the category of $R\Gamma$ -modules is an abelian category where the notions submodule, quotient module, kernel, image, and cokernel are defined object-wise. The direct sum of $R\Gamma$ -modules is given by taking the usual direct sum object-wise.

Example 2.1. The most important example for our applications is the orbit category of a finite group. Let G be a finite group and let \mathcal{F} be a family of subgroups of G which is closed under conjugation and taking subgroups. The orbit category $\text{Or}(G)$ is the category whose objects are subgroups H of G or coset spaces G/H of G , and the morphisms $\text{Mor}(G/H, G/K)$ are given by the set of G -maps $f: G/H \rightarrow G/K$.

The category $\Gamma_G = \text{Or}_{\mathcal{F}} G$ is defined as the full subcategory of $\text{Or}(G)$ where the objects satisfy $H \in \mathcal{F}$. The category of right $R\Gamma_G$ -modules is the category of contravariant functors from $\text{Or}_{\mathcal{F}} G$ to R -modules. A right $R\Gamma_G$ -module M is often called a *coefficient system* [35]. We will sometimes denote $M(G/H)$ by $M(H)$ if the group G is clear from the context. When $\mathcal{F} = \{e\}$, $R\Gamma_G\text{-Mod}$ is just the category of left RG -modules and $\text{Mod-}R\Gamma_G$ is just the category of right RG -modules. \square

Now, we will introduce the tensor product and Hom functors for modules over small categories. Let Γ be a small category and let $M \in \text{Mod-}R\Gamma$ and $N \in R\Gamma\text{-Mod}$. The tensor product over $R\Gamma$ is given by

$$M \otimes_{R\Gamma} N = \bigoplus_{x \in \text{Ob}(\Gamma)} M(x) \otimes N(x) / \sim$$

where \sim is the equivalence relation defined by $\varphi^*(m) \otimes n \sim m \otimes \varphi_*(n)$ for every morphism $\varphi: x \rightarrow y$. For $R\Gamma$ -modules M and N , we mean by $\text{Hom}_{R\Gamma}(M, N)$ the R -module of $R\Gamma$ -homomorphisms from M to N . In other words,

$$\text{Hom}_{R\Gamma}(M, N) \subseteq \bigoplus_{x \in \text{Ob}(\Gamma)} \text{Hom}_R(M(x), N(x))$$

is the submodule satisfying the relations $f(x) \circ \varphi^* = \varphi^* \circ f(y)$, for every morphism $\varphi: x \rightarrow y$. We sometimes consider a second tensor product, namely the tensor product over R , which is defined for $R\Gamma$ -modules M and N which are both left modules or both right modules. The tensor product $M \otimes_R N$ is defined by the formula

$$[M \otimes_R N](x) = M(x) \otimes_R N(x)$$

on objects $x \in \text{Ob}(\Gamma)$ and on morphisms, one has $[M \otimes_R N](f) = M(f) \otimes_R N(f)$.

The tensor product over $R\Gamma$ and $\text{Hom}_{R\Gamma}$ are adjoint to each other. This can be described in the following way:

Proposition 2.2. *Given two small categories Γ and Λ , the category of $R\Gamma$ - $R\Lambda$ -bimodules is defined as the category of functors $\Gamma \times \Lambda^{\text{op}} \rightarrow R\text{-Mod}$. For a right $R\Gamma$ -module M , an $R\Gamma$ - $R\Lambda$ -bimodule B , and a right $R\Lambda$ -module N , one has a natural transformation*

$$\text{Hom}_{R\Lambda}(M \otimes_{R\Gamma} B, N) \cong \text{Hom}_{R\Gamma}(M, \text{Hom}_{R\Lambda}(B, N)).$$

Proof. See [20, 9.21.] \square

We will be using this isomorphism later when we are discussing induction and restriction.

2A. Free and finitely generated modules. For a small category Γ , a sequence

$$M' \rightarrow M \rightarrow M''$$

of $R\Gamma$ -modules is exact if and only if

$$M'(x) \rightarrow M(x) \rightarrow M''(x)$$

is exact for all $x \in \text{Ob}(\Gamma)$. Recall that a module P in $\text{Mod-}R\Gamma$ is projective if the functor

$$\text{Hom}_{R\Gamma}(P, -): \text{Mod-}R\Gamma \rightarrow R\text{-Mod}$$

is exact. For an object $x \in \Gamma$, we define a right $R\Gamma$ -module $R\Gamma(?, x)$ by setting

$$R\Gamma(?, x)(y) = R\text{Mor}(y, x)$$

for all $y \in \text{Ob}(\Gamma)$. Here, $R\text{Mor}(y, x)$ denotes the free abelian group on the set of morphisms $\text{Mor}(y, x)$ from y to x . As a consequence of the Yoneda lemma, we have

$$\text{Hom}_{R\Gamma}(R\Gamma(?, x), M) \cong M(x).$$

So, for each $x \in \text{Ob}(\Gamma)$, the module $R\Gamma(?, x)$ is a projective module. When working with modules over small categories one uses the following notion of free modules.

Definition 2.3. Let Γ be a small category. A $\text{Ob}(\Gamma)$ -set is defined as a set S together with a map $\beta: S \rightarrow \text{Ob}(\Gamma)$. We say a $R\Gamma$ -module M is free if it is isomorphic to a module of the form

$$R\Gamma(S) = \bigoplus_{b \in S} R\Gamma(?, \beta(b))$$

for some $\text{Ob}(\Gamma)$ -set S . A free module $R\Gamma(S)$ is called finitely generated if the set S is finite.

Note that for every $R\Gamma$ -module M , there is a free $R\Gamma$ -module $R\Gamma(S)$ and a map $f: R\Gamma(S) \rightarrow M$ such that f is surjective. We can take such a free module by choosing a set of generators S_x for the R -module $M(x)$ for each $x \in \text{Ob}(\Gamma)$, and then taking S as the $\text{Ob}(\Gamma)$ -set which has the property $\beta^{-1}(x) = S_x$. A free module $R\Gamma(S)$ which maps surjectively on M is called a *free cover* of M . A $R\Gamma$ -module is called *finitely generated* if it has a finitely generated free cover.

It is clear from our description of free modules that an $R\Gamma$ -module M is projective if and only if it is a direct summand of a free module. This shows that the module category of a small category has enough projectives. We will later give a more detailed description of projective $R\Gamma$ -modules.

Example 2.4. For the orbit category $\Gamma = \text{Or}(G)$, the free modules described above have a more specific meaning. For any subgroup $K \leq G$, the $R\Gamma$ -module $R\Gamma(?, G/K)$ is given by

$$R\Gamma(?, G/K)(G/H) = R\text{Mor}(G/H, G/K) = R[(G/K)^H]$$

where $R[(G/K)^H]$ is the free abelian group on the set of fixed points of the H action on G/K . Because of this, we denote the free module $R\Gamma(?, G/K)$ by $R[G/K]^?$.

If \mathcal{F} is a family of subgroups, and $\Gamma_G = \text{Or}_{\mathcal{F}} G$, we obtain free $R\Gamma_G$ -modules $R[G/K^?]$ by restriction whenever $K \in \mathcal{F}$. The *constant* $R\Gamma_G$ -module \underline{R} defined by $\underline{R}(H) = R$, for all $H \in \mathcal{F}$, is just the restriction to $R\Gamma_G$ of the module $\underline{R} = R[G/G^?]$. This shows that the constant module \underline{R} is projective if $G \in \mathcal{F}$. More generally, if $K \in \mathcal{F}$, a non-empty fixed set

$$(G/K)^H = \{gK \mid g^{-1}Hg \subseteq K\} \neq \emptyset$$

implies $H \in \mathcal{F}$, since \mathcal{F} is closed under conjugation and taking subgroups. Therefore, $R[G/K^?](H) = 0$ for $H \notin \mathcal{F}$, whenever $K \in \mathcal{F}$.

2B. Induction and Restriction. We now recall the definitions and terminology for these terms presented in Lück [20, 9.15]. Let Γ and Λ be two small categories. Given a covariant functor $F: \Lambda \rightarrow \Gamma$, we define an $R\Lambda$ - $R\Gamma$ -bimodule

$$R(??, F(?)): \Lambda \times \Gamma^{\text{op}} \rightarrow R\text{-Mod}$$

on objects by $(x, y) \rightarrow R\text{Hom}(y, F(x))$. We define the restriction map

$$\text{Res}_F: \text{Mod-}R\Gamma \rightarrow \text{Mod-}R\Lambda$$

as the composition with F . The induction map

$$\text{Ind}_F: \text{Mod-}R\Lambda \rightarrow \text{Mod-}R\Gamma$$

is defined by

$$\text{Ind}_F(M)(??) = M \otimes_{R\Lambda} R(??, F(?))$$

for every $R\Lambda$ -module M . For every right $R\Gamma$ -module N , the $R\Lambda$ -module

$$\text{Hom}_{R\Gamma}(R(??, F(?)), N)$$

is the same as the composition $\Lambda \xrightarrow{F} \Gamma \xrightarrow{N} R\text{-Mod}$. So, by Proposition 2.2, we can conclude the following:

Proposition 2.5. *Induction and restriction are adjoint functors: for any $R\Gamma$ -module M and $R\Lambda$ -module N , there is a natural isomorphism*

$$\text{Hom}_{R\Gamma}(\text{Ind}_F M, N) = \text{Hom}_{R\Lambda}(M, \text{Res}_F N).$$

The induction functor respects direct sum, finitely generated, free, and projective but it is not exact in general. The restriction functor is exact but does not respect finitely generated, free, or projective in general.

Now we will define functors which are special cases of the restriction and induction functors. Let Γ be a small category. For $x \in \text{Ob}(\Gamma)$, we define $R[x] = R\text{Aut}(x)$ to be the group ring of the automorphism group $\text{Aut}(x)$ and denote the category of right $R[x]$ -modules by $\text{Mod-}R[x]$. Let Γ_x denote the full subcategory of Γ with single object x and let $F: \Gamma_x \rightarrow \Gamma$ be the inclusion natural transformation. The restriction functor associated to F gives a functor

$$\text{Res}_x: \text{Mod-}R\Gamma \rightarrow \text{Mod-}R[x]$$

which is called the **restriction functor**. This functor behaves like an evaluation map $\text{Res}_x(M) = M(x)$. In the other direction, the induction functor associated to F gives a functor

$$E_x: \text{Mod-}R[x] \rightarrow \text{Mod-}R\Gamma$$

which is called the **extension functor**. For a $R[x]$ -module M , we define $E_x(M)(y) = M \otimes_{R[x]} R\text{Mor}(y, x)$ for every $y \in \text{Ob}(\Gamma)$. They form an adjoint pair: for every $R[x]$ -module M and an $R\Gamma$ -module N , we have

$$\text{Hom}_{R\Gamma}(E_x M, N) \cong \text{Hom}_{R[x]}(M, \text{Res}_x N).$$

By general properties of restriction and induction, the functor Res_x is exact and E_x takes projectives to projectives. In general, E_x is not exact and Res_x does not take projectives to projectives. But in some special cases, we can say more. For example, when Γ is free, i.e. $R\text{Mor}(y, x)$ is a free $R[x]$ -module for all $y \in \Gamma$, then it is easy to see that E_x is exact [20, 16.9].

Example 2.6. In the case of an orbit category $\Gamma_G = \text{Or}_{\mathcal{F}} G$, we denote the extension function for $H \in \mathcal{F}$ simply by E_H and the restriction functor by Res_H . In this case, the automorphism group $\text{Aut}(G/H)$ for $H \in \mathcal{F}$ is isomorphic to the quotient group $N_G(H)/H$. The isomorphism $N_G(H)/H \cong \text{Aut}(G/H)$ is given by the isomorphism $nH \rightarrow f_n$ where $f_n(gH) = gn^{-1}H$ for $n \in N_G(H)$ (see [10, Example 11.2]). This isomorphism takes right $R[x]$ -modules to right $R[N_G(H)/H]$ -modules, so given a right $R\Gamma$ -module M , the evaluation at $H \in \mathcal{F}$ gives a right $R[N_G(H)/H]$ -module.

It is easy to see that the morphism set $\text{Mor}(G/K, G/H)$ is a free $[N_G(H)/H]$ -set, so $\text{Or}_{\mathcal{F}} G$ is free in the above sense [20, Example 16.2]. Therefore, the functor E_H is exact and preserves projectives, whereas Res_H is exact but does not necessarily preserve projectives. For example, the module $\mathbf{Z}[G/G^?]$ is free over $\mathbf{Z}\text{Or}(G)$ by definition, but $\text{Res}_H \mathbf{Z}[G/G^?] = \mathbf{Z}$ is not projective whenever $N_G(H)/H \neq 1$.

2C. Inclusion and Splitting Functors. We will introduce two more functors. These are also special cases of induction and restriction, but they are defined through a bimodule rather than just a natural transformation F . We first describe these functors and then give their interpretations as restriction and induction functors.

Let Γ be an EI-category. By this, we mean that Γ is a small category where every endomorphism $x \rightarrow x$ is an isomorphism for all $x \in \text{Ob}(\Gamma)$. This allows us to define a partial ordering on the set $\text{Iso}(\Gamma)$ of isomorphism classes \bar{x} of objects x in Γ . For $x, y \in \text{Ob}(\Gamma)$, we say $\bar{x} \leq \bar{y}$ if and only if $\text{Mor}(x, y) \neq \emptyset$. The EI-property ensures that $\bar{x} \leq \bar{y} \leq \bar{x}$ implies $\bar{x} = \bar{y}$.

For each object $x \in \Gamma$, and $M \in \text{Mod-}R[x]$, the **inclusion functor**,

$$I_x: \text{Mod-}R[x] \rightarrow \text{Mod-}R\Gamma$$

is defined by

$$I_x M(y) = \begin{cases} M \otimes_{R[x]} R\text{Mor}(y, x) & \text{if } \bar{y} = \bar{x} \\ \{0\} & \text{if } \bar{y} \neq \bar{x}. \end{cases}$$

In the other direction, we define the **splitting functor**

$$S_x: \text{Mod-}R\Gamma \rightarrow \text{Mod-}R[x]$$

by $S_x(M) = M(x)/M(x)_s$ where $M(x)_s$ is the R -submodule of $M(x)$ which is generated by the images of $M(f): M(y) \rightarrow M(x)$ for all $f: x \rightarrow y$ with $\bar{x} \leq \bar{y}$ and $\bar{x} \neq \bar{y}$.

There is a $R\Gamma$ - $R[x]$ -bimodule B defined in such a way that the inclusion functor I_x can be described as $M \rightarrow \text{Hom}_{R[x]}(B, M)$ and the splitting functor S_x is the same as the functor $M \rightarrow M \otimes_{R\Gamma} B$ (see [20, page 171] for details). So (S_x, I_x) is an adjoint pair, meaning that

$$\text{Hom}_{R[x]}(S_x M, N) \cong \text{Hom}_{R\Gamma}(M, I_x N)$$

for every $R\Gamma$ -module M and $R[x]$ -module N .

From general properties of induction and restriction, we can conclude that I_x is exact and S_x preserves projectives. Some of the other properties of these functors are listed in [20, Lemma 9.31]. It is interesting to note that the composition $S_x \circ E_x$ is naturally equivalent to the identity functor. Also, the composition $S_y \circ E_x$ is zero when $\bar{x} \neq \bar{y}$. These are used to give a splitting for projective $R\Gamma$ -modules.

Theorem 2.7. *Let P be a finitely generated projective $R\Gamma$ -module. Then*

$$P \cong \bigoplus_{x \in \text{Iso}(\Gamma)} E_x S_x(P).$$

Proof. For proof see [20, Corollary 9.40]. □

In the statement, the notation $\bigoplus_{x \in \text{Iso}(\Gamma)}$ means that the sum is over a set of representatives $x \in \text{Ob}(\Gamma)$ for $\bar{x} \in \text{Iso}(\Gamma)$.

2D. Resolutions for $R\Gamma$ -modules. Let Γ be an EI-category. For a non-negative integer l we define an l -chain c from $x \in \text{Ob}(\Gamma)$ to $y \in \text{Ob}(\Gamma)$ to be a sequence

$$c: \bar{x} = \bar{x}_0 < \bar{x}_1 < \cdots < \bar{x}_l = \bar{y}.$$

Define the length $l(y)$ of $y \in \text{Ob}(\Gamma)$ to be the largest integer l such that there exists an l -chain from some $x \in \text{Ob}(\Gamma)$ to y . The length $l(\Gamma)$ of Γ is $\max\{l(x) \mid x \in \text{Ob}(\Gamma)\}$. Given an $R\Gamma$ -module M , its length $l(M)$ is defined by $\max\{l(x) \mid M(x) \neq 0\}$ if M is not the zero module and $l(\{0\}) = -1$.

We call a category Γ *finite* if $\text{Iso}(\Gamma)$ and $\text{Mor}(x, y)$ are finite for all $x, y \in \text{Ob}(\Gamma)$. Denote by $m(\Gamma)$ the least common multiple of all the integers $|\text{Aut}(x)|$.

Given an $R\Gamma$ -module M , consider the map

$$\phi: \bigoplus_{x \in \text{Iso}(\Gamma)} E_x \text{Res}_x M \rightarrow M$$

where for each $x \in \text{Ob}(\Gamma)$, the map $\phi_x: E_x \text{Res}_x M \rightarrow M$ is the map adjoint to the identity homomorphism. It is easy to see that ϕ is surjective. Let

$$EM := \bigoplus_{x \in \text{Iso}(\Gamma)} E_x \text{Res}_x M$$

and let KM denote the kernel of $\phi: EM \rightarrow M$. Note that if x is an object with $l(x) = l(M)$, then $\text{Res}_x = S_x$ which also gives that

$$\text{Res}_x \phi: \text{Res}_x E_x \text{Res}_x M \rightarrow \text{Res}_x M$$

is an isomorphism. Note that this implies $l(KM) < l(M)$ which allows one to proceed by induction and obtain the following:

Proposition 2.8. *If Γ is finite EI-category, then every nonzero M has a finite resolution of the form*

$$0 \rightarrow EK^t M \rightarrow \cdots \rightarrow EKM \rightarrow EM \rightarrow M \rightarrow 0 .$$

where $t = l(M)$.

Proof. See [20, 17.13]. Here $K^0 M = M$ and $K^s M = K(K^{s-1} M)$. □

From the description above it is easy to see that

$$EK^s M := \bigoplus_{x \in \text{Iso}(\Gamma)} E_x \text{Res}_x K^s M$$

where $\text{Res}_x K^s M$ is isomorphic to a direct sum of $R[x]$ -modules

$$M(c) := M(x_0) \otimes_{R[x_0]} R \text{Mor}(x_1, x_0) \otimes_{R[x_1]} \cdots \otimes_{R[x_{s-1}]} R \text{Mor}(x, x_{s-1})$$

over representatives in $\text{Ob}(\Gamma)$ for all the chains of the form $c : \bar{x} < \bar{x}_{s-1} < \cdots < \bar{x}_0$ (see [20, 17.24]). Note that if Γ is a finite, free EI-category, then the resolution given in Proposition 2.8 will be a finite projective resolution if $M(c)$ is projective as an $R[x]$ -module for every chain c . This gives the following:

Proposition 2.9. *Let M be $R\Gamma_G$ module where $\Gamma_G = \text{Or}_{\mathcal{F}} G$ for some finite group G and R is a commutative ring such that $|G|$ is invertible in R . Suppose also that $M(H)$ is projective as an R -module for all $H \in \mathcal{F}$. Then, M has a projective resolution with length less than or equal to $l(\Gamma)$.*

Proof. See [20, 17.31]. □

In particular, if $R = \mathbf{Z}_{(p)}$ with $p \nmid |G|$ and if M is a $R\Gamma$ -module such that $M(H)$ is R -torsion free for all $H \in \mathcal{F}$, then M has a finite projective resolution of length less than or equal to $l(M)$.

3. THE PROOF OF THEOREM B

The main result of this section is Theorem 3.9, which is an orbit category version of a well-known result of Rim [29]. We first collect some further information about induction and restriction for subgroups.

Let G be a finite group and let H be a subgroup of G . Given a family of subgroups \mathcal{F} of G , we consider the orbit categories $\Gamma_G = \text{Or}_{\mathcal{F}} G$ and $\Gamma_H = \text{Or}_{\mathcal{F}} H$, where the objects of Γ_H are orbits of H with isotropy in $\mathcal{F}_H = \{K \leq H \mid K \in \mathcal{F}\}$. Let $F: \Gamma_H \rightarrow \Gamma_G$ be the functor which takes H/K to G/K and sends an H -map $f: H/K \rightarrow H/L$ to the induced G -map

$$\text{Ind}_H^G(f): G/K = G \times_H H/K \rightarrow G \times_H H/L = G/L$$

for every $K, L \in \mathcal{F}_H$. Note that if f is the map which takes eK to hL , then $\text{Ind}_H^G(f)(gK) = ghL$. The restriction and induction functors (see Proposition 2.5) associated to this functor gives us two adjoint functors

$$\text{Res}_H^G: \text{Mod-}\Gamma_G \rightarrow \text{Mod-}\Gamma_H$$

and

$$\text{Ind}_H^G: \text{Mod-}\Gamma_H \rightarrow \text{Mod-}\Gamma_G.$$

The restriction functor is defined as the composition with F . So, for a $R\Gamma_G$ -module M , we have $(\text{Res}_H^G M)(K) = M(K)$, for all $K \in \mathcal{F}_H$. For the induced module we have the following formula:

Lemma 3.1. *Let N be a $R\Gamma_H$ -module and $K \leq G$. Then,*

$$(\text{Ind}_H^G N)(K) \cong \bigoplus_{gH \in G/H, K^g \leq H} N(K^g)$$

where $K^g = g^{-1}Kg$.

Proof. For a (right) $R\Gamma_H$ -module N , the induced module $\text{Ind}_H^G N$ is defined as the direct sum

$$\bigoplus_{L \leq H} N(L) \otimes_R R\text{Mor}(G/K, G/L)$$

modulo the relations $n \otimes \varphi f \sim \varphi^*(n) \otimes f$ where $n \in N(L)$, $f \in \text{Mor}(G/K, G/L')$ and $\varphi = \text{Ind}_H^G(\phi)$ for some $\phi: H/L' \rightarrow H/L$. Every morphism $G/K \rightarrow G/L$ which satisfies the condition $L \leq H$ can be written as a composition φf_g where $\varphi: G/K^g \rightarrow G/L$ is induced from an H -map and $f_g: G/K \rightarrow G/K^g$ is given by $xK \rightarrow xgK^g$, for some $g \in G$.

This shows that every element in the above sum is equivalent to an element of the form $n \otimes f_g$ where $n \in N(K^g)$ and $f_g: G/K \rightarrow G/K^g$ is as above with $K^g \leq H$. There is one summand for each gH satisfying $K^g \leq H$. \square

Note that we can also express the above formula by

$$(\text{Ind}_H^G N)(K) \cong \bigoplus_{gH \in (G/H)^K} N(K^g).$$

If $J \leq K$, then the argument above can be extended to show that restriction map

$$(\text{Ind}_H^G N)(K) \rightarrow (\text{Ind}_H^G N)(J)$$

is given by the coordinate-wise restriction maps $N(K^g) \rightarrow N(J^g)$. Note that if $gH \in (G/H)^K$, then $gH \in (G/H)^J$. Similarly, the conjugation map

$$(\text{Ind}_H^G N)(K) \rightarrow (\text{Ind}_H^G N)({}^xK)$$

can be described by coordinate-wise conjugation maps. From these, it is easy to see that $\text{Ind}_H^G R \cong R[G/H^?]$. A generalization of this argument gives the following:

Lemma 3.2. [35, Cor. 2.12]. *Let G be a finite group and let H be a subgroup of G . For every $R\Gamma_G$ -module M , we have $\text{Ind}_H^G \text{Res}_H^G M \cong M \otimes_R R[G/H^?]$.*

We also have the following formulas:

Lemma 3.3. *Let G be a finite group and let H be a subgroup of G .*

- (i) *For every $K \leq H$, we have $\text{Ind}_H^G R[H/K] \cong R[G/K]$.*
- (ii) *For every $K \leq G$, we have $\text{Res}_H^G R[G/K] \cong \bigoplus_{K \backslash G/H} R[H/(H \cap {}^g K)]$.*

Proof. Part (i) follows from the fact that $\text{Ind}_H^G \text{Ind}_K^H = \text{Ind}_K^G$ which is a consequence of a more general formula $\text{Ind}_F \text{Ind}_{F'} = \text{Ind}_{F \circ F'}$. We can prove this more general formula by using adjointness and the formula $\text{Res}_{F'} \text{Res}_F = \text{Res}_{F \circ F'}$. For (ii), observe that the definition of $R[G/H]$ can be extended to define a $R\Gamma_G$ -module $R[S]$ for every G -set S , by taking

$$(3.4) \quad R[S](G/K) = R \text{Map}_G(G/K, S)$$

for every $K \in \mathcal{F}$, where $\text{Map}_G(G/K, S)$ denotes the set of G -maps from G/K to S . For G -sets S and T , we have an isomorphism $R[(S \sqcup T)] \cong R[S] \oplus R[T]$. By the definition of restriction map, we get

$$\left(\text{Res}_H^G R[S] \right)(H/K) = R \text{Map}_G(G/K, S) = R \text{Map}_H(H/K, \text{Res}_H^G S).$$

It is easy to see that this induces an $R\Gamma_H$ -module isomorphism

$$\text{Res}_H^G R[S] \cong R[(\text{Res}_H^G S)].$$

Since

$$\text{Res}_H^G(G/K) \cong \coprod_{H \backslash G/K} H/(H \cap {}^g K)$$

as G -sets, we obtain the formula given in (ii). \square

Example 3.5. Let $G = S_5$ be the symmetric group on $\{1, 2, 3, 4, 5\}$ and $H = S_4$ be the subgroup of symmetries that fix 5. Let $C_2 = \langle (12) \rangle$ and $C_3 = \langle (345) \rangle$. The formula in Lemma 3.3 (ii) gives

$$\text{Res}_H^G R[G/(C_2 \times C_3)] = R[H/C_2] \oplus R[H/{}^g C_3]$$

where ${}^g C_3 = \langle (123) \rangle$. From this expression we obtain

$$R[G/(C_2 \times C_3)](G/C_2) \cong RH/C_2 \cong R[N_H(C_2)/C_2],$$

as an $N_H(C_2)/C_2$ -module, where $N_H(C_2) = C_2 \times C_2$. Note that $N_G(C_2) = C_2 \times S_3$ and as an $N_G(C_2)/C_2$ -module $R[G/(C_2 \times C_3)](G/C_2)$ is isomorphic to $R[C_2 \times S_3/C_2 \times C_3]$. \square

We can give a more general formula for $R[G/H](G/K)$ as follows:

Lemma 3.6. *Let G be a finite group, and H and K be two subgroups of G . Then, as an $R[N_G(H)/H]$ -module*

$$R[G/K](G/H) \cong \bigoplus_{v(H,K)} R[N_G(H)/N_{{}^g K}(H)]$$

where the sum is over the set $v(H, K)$ of representatives of K -conjugacy classes of subgroups H^g such that $H^g \leq K$.

Proof. This formula can easily be proved by first determining the orbits of $N_G(H)$ action on $(G/K)^H = \{gK \mid H^g \leq K\}$, and then by calculating the isotropy subgroups for each of these orbits. A similar computation can be found in the proof of Theorem 4.1 in [8]. \square

Proposition 3.7. *Both Res_H^G and Ind_H^G are exact and take projectives to projectives.*

Proof. The fact that Res_H^G is exact and Ind_H^G preserves projectives follows from the general properties of restriction and induction functor associated to a natural transformation F . The fact that Ind_H^G is exact follows from the formula given in Lemma 3.1. Finally, to show that Res_H^G takes projective to projectives, it is enough to show it takes free modules to projective modules. An indecomposable free $R\Gamma_G$ -module M is of the form $R[G/K^?]$ for some $K \in \mathcal{F}$. By Lemma 3.3, $\text{Res}_H^G(R[G/K^?])$ will be projective if $H \cap {}^gK$ is in \mathcal{F} for all $HgK \in H \backslash G/K$. But this is always true since the family \mathcal{F} is closed under conjugation and taking subgroups. \square

A result of Rim [29] relates projectivity over the group ring $\mathbf{Z}G$ to projectivity over the p -Sylow subgroups.

Proposition 3.8 (Rim's Theorem). *Let G be a finite group, and M be a finitely generated $\mathbf{Z}G$ -module. Then M is projective over $\mathbf{Z}G$ if and only if $\text{Res}_P^G M$ is projective over $\mathbf{Z}P$ for any p -Sylow subgroup $P \leq G$.*

Proof. A module M is $\mathbf{Z}G$ -projective if and only if $\text{Ext}_{\mathbf{Z}G}^1(M, N) = 0$ for every $\mathbf{Z}G$ -module N . Therefore M is projective if and only if $\mathbf{Z}_{(p)} \otimes_{\mathbf{Z}} M$ is projective over $\mathbf{Z}_{(p)}G$ for all primes p dividing the order of G .

For any p -Sylow subgroup $P \leq G$, the permutation module $R[G/P] \cong R \oplus N$ splits when $R = \mathbf{Z}_{(p)}$. Therefore, if M is any RG -module, $M \otimes_R R[G/P] \cong M \oplus (M \otimes_R N)$. Since $M \otimes_R R[G/P] \cong \text{Ind}_P^G \text{Res}_P^G M$, the projectivity of M is equivalent to the projectivity of $\text{Res}_P^G M$. \square

Here is an orbit category version of this result.

Theorem 3.9 (Rim's Theorem for the Orbit Category). *Let G be a finite group and let M be a $R\Gamma_G$ -module where $R = \mathbf{Z}_{(p)}$. Suppose that \mathcal{F} is a family of p -subgroups in G . Then M has a finite projective resolution if and only if $\text{Res}_P^G M$ has a finite projective resolution for any p -Sylow subgroup P of G .*

Proof. One direction is clear since Res_P^G is exact and takes projectives to projectives. For the other direction, we will give the proof by induction on the length $l(M)$ of M . Without loss of generality, we can assume that $M(H)$ is R -torsion free for all $H \in \mathcal{F}$. Suppose M is a $R\Gamma_G$ -module with $l(M) = 0$. Then, we can regard M as an RG -module. If $\text{Res}_P^G M$ has a finite projective resolution, then $\text{Res}_P^G M$ must be projective (see [20, page 348]). Then, by Rim's theorem, M is a projective RG -module, hence has finite projective length.

Now, assume M is an $R\Gamma_G$ -module with $l(M) = s > 0$. Let

$$0 \rightarrow P_n \rightarrow \cdots \rightarrow P_0 \rightarrow \text{Res}_P^G M \rightarrow 0$$

be a projective resolution for $\text{Res}_P^G M$. We can assume that $l(P_i) \leq s$ for all i . Then, for $Q \in \mathcal{F}$ with $l(Q) = s$, we have

$$S_Q P_i = \text{Res}_Q P_i = P_i(Q).$$

Since S_Q takes projectives to projectives, the resolution

$$0 \rightarrow P_n(Q) \rightarrow \cdots \rightarrow P_0(Q) \rightarrow (\text{Res}_P^G M)(Q) \rightarrow 0$$

is a finite projective resolution of $(\text{Res}_P^G M)(Q) = M(Q)$ as an $R[N_P(Q)/Q]$ -module. This gives that $M(Q)$ is projective as an $R[N_P(Q)/Q]$ -module.

Lemma 3.10. *For every p -group Q , there is a p -Sylow subgroup P of G such that $N_P(Q)$ is a p -Sylow subgroup of $N_G(Q)$.*

Proof. Let S be a p -Sylow subgroup of $N_G(Q)$, and pick a p -Sylow subgroup P of G containing S . Since $N_P(Q) = N_G(Q) \cap P$ is a p -subgroup of $N_G(Q)$, we have $|N_P(Q)| \leq |S|$. But $S \leq P$ and $S \leq N_G(Q)$ implies $S \leq N_P(Q)$. Therefore $S = N_P(Q)$. \square

We can assume P is a p -Sylow subgroup which has this property. Then, by the p -local version of Rim's theorem, we can conclude that $M(Q)$ is projective as an $R[N_G(Q)/Q]$ -module. Now, consider the map

$$\psi = (\psi_Q): \bigoplus_{Q \in \text{Iso}(\Gamma_G), l(Q)=s} E_Q \circ \text{Res}_Q M \rightarrow M$$

where $\psi_Q: E_Q \circ \text{Res}_Q M \rightarrow M$ is the map adjoint to the identity map $\text{id}: \text{Res}_Q M \rightarrow \text{Res}_Q M$. For every $K \in \mathcal{F}$ with $l(K) = s$, the induced map $\psi(K)$ is an isomorphism. This is because

$$(E_Q \circ \text{Res}_Q M)(K) = \text{Res}_K E_Q \text{Res}_Q M = S_K E_Q \text{Res}_Q M \cong M(K)$$

if K is conjugate to Q and zero otherwise. So, we have $l(\text{coker } \psi) < s$. Therefore, there is a finitely generated projective $R\Gamma_G$ -module P with $l(P) < s$, and a map $\alpha: P \rightarrow M$ such that $\psi \oplus \alpha$ is surjective. If K is the kernel of $\psi \oplus \alpha$, we get an exact sequence of $R\Gamma_G$ -modules

$$0 \rightarrow K \rightarrow P \oplus \bigoplus_{Q \in \text{Iso}(\Gamma_G), l(Q)=s} E_Q \circ \text{Res}_Q M \rightarrow M \rightarrow 0$$

where the middle term is projective as an $R\Gamma_G$ -module, and $l(K) < s$. Note that $\text{Res}_P^G K$ must have a finite projective resolution by [20, Lemma 11.6]. So, by induction, K has a finite projective resolution, and hence M has a finite projective resolution as well. \square

Remark 3.11. The inductive argument we use in the above proof is similar to the argument used by Lück to prove Proposition 17.31 in [20]. By this result, any module M over a finite EI-category Γ which has a finite projective resolution, admits a resolution of length $\leq l(M)$ provided that $M(x)$ is R -projective for all $x \in \text{Ob}(\Gamma)$. \square

It isn't clear to us how to generalize Theorem 3.9 to integer coefficients. For $R = \mathbf{Z}_{(p)}$, the following example shows that the result does not hold for an arbitrary family \mathcal{F} .

Example 3.12. Let $G = S_5$ and $R = \mathbf{Z}_{(2)}$, and take \mathcal{F} as the family of all 2-subgroups and 3-subgroups in G . Consider the $R\Gamma_G$ -module $M = R[G/(C_2 \times C_3)^?]$ where C_2 and C_3 are as in Example 3.5. It is clear that the restriction of M to a 2-Sylow subgroup is projective (since its restriction to $H = S_4$ is already projective), but M does not have a finite projective resolution as an $R\Gamma_G$ -module.

To see this, suppose that M has a finite projective resolution $\mathbf{P} \rightarrow M$. Then, $\mathbf{P}(C_3)$ will be a finite projective resolution for $M(C_3)$ over $R[N_G(C_3)/C_3]$. This is because $C_3 = \langle (123) \rangle$ is a maximal subgroup in \mathcal{F} . This implies

$$M(C_3) \cong R[S_3 \times C_2/C_3 \times C_2] \cong R[C_2]$$

is projective as an $R[N_G(C_3)/C_3]$ -module. But,

$$R[N_G(C_3)/C_3] = R[S_3 \times C_2/C_3] \cong R[C_2 \times C_2],$$

and it is clear that $R[C_2]$ is not projective as an $R[C_2 \times C_2]$ -module. So, M does not have a finite projective resolution. \square

On the other hand, the following holds for modules over orbit categories:

Proposition 3.13. *Let G be a finite group, and \mathcal{F} be a family of subgroups of G . Then, a $\mathbf{Z}\Gamma_G$ -module M has a finite projective resolution if and only if $\mathbf{Z}_{(p)} \otimes_{\mathbf{Z}} M$ has a finite projective resolution over $\mathbf{Z}_{(p)}\Gamma_G$, for all primes p dividing the order of G .*

The proof of this statement follows from Propositions 4.4 and 4.5 in the next section. We end this section with some corollaries of Theorem 3.9.

Corollary 3.14. *Let G be a finite group and $R = \mathbf{Z}_{(p)}$. Suppose that \mathcal{F} is a family of p -subgroups. Then, $R[G/H^?]$ has a finite projective resolution over $R\Gamma_G$ if a p -Sylow subgroup of H is included in \mathcal{F} .*

Proof. If a p -Sylow subgroup of H is in \mathcal{F} , then $\text{Res}_P^G R[G/H^?]$ is a free $R\Gamma_P$ -module for any $P \in \text{Syl}_p(G)$. So, by Theorem 3.9, it has a finite projective resolution. \square

As a special case of this corollary, we obtain the following known result (see [4, 6.8], [35, 2.5 and p. 296], [17], [14]).

Corollary 3.15. *Let G be a finite group and $R = \mathbf{Z}_{(p)}$. Then, \underline{R} has a finite projective resolution over $R\Gamma_G$ relative to the family of all p -subgroups of G .*

Proof. This follows from $\underline{R} = R[G/G^?]$. \square

4. MACKEY STRUCTURES ON $\text{Ext}_{R\Gamma_G}^*(M, N)$

The notation and results of the previous sections will now be used to establish some structural and computational facts about the Ext-groups over the orbit category. Our main sources are Cartan-Eilenberg [7] and tom Dieck [10, §II.9] (see also [17], [14]).

We have seen that the category of right $R\Gamma$ -modules has enough projectives to define the bifunctor

$$\text{Ext}_{R\Gamma}^*(M, N) = H^*(\text{Hom}_{R\Gamma}(\mathbf{P}, N))$$

via any projective resolution $\mathbf{P} \twoheadrightarrow M$ (see [20, Chap. III, §17], [23, Chap. III.6]). The following property is also useful (see Lück [20, 17.21]).

Lemma 4.1. *If Γ is a free EI-category, then $\mathrm{Ext}_{R\Gamma}^*(E_x M, N) \cong \mathrm{Ext}_{R[x]}^*(M, \mathrm{Res}_x N)$.*

Proof. Take a projective resolution \mathbf{P} of M . Since Γ is free, the extension functor E_x is exact [20, 16.9]. In addition, E_x preserves projectives and is adjoint to the restriction functor Res_x by Proposition 2.5. Therefore

$$\cdots \rightarrow E_x P_n \rightarrow \cdots \rightarrow E_x P_1 \rightarrow E_x P_0 \rightarrow E_x M \rightarrow 0$$

is a projective resolution of $E_x M$, and applying Hom over the orbit category gives

$$\begin{aligned} \mathrm{Ext}_{R\Gamma}^n(E_x M, N) &= H^n(\mathrm{Hom}_{R\Gamma}(E_x \mathbf{P}, N)) \\ &\cong H^n(\mathrm{Hom}_{R[x]}(\mathbf{P}, \mathrm{Res}_x N)) = \mathrm{Ext}_{R[x]}^n(M, \mathrm{Res}_x N). \quad \square \end{aligned}$$

In the rest of this section, we assume that $\Gamma_G = \mathrm{Or}_{\mathcal{F}} G$ for a finite group G , where \mathcal{F} is a family of subgroups in G . Note that Γ_G is both finite and free as an EI-category. If there are two groups $H \leq G$, we use the notations $\Gamma_G = \mathrm{Or}_{\mathcal{F}} G$ for the orbit category with respect to the family \mathcal{F} , and $\Gamma_H = \mathrm{Or}_{\mathcal{F}} H$ for the orbit category with respect to the family $\mathcal{F}_H = \{H \cap K \mid K \in \mathcal{F}\}$.

Proposition 4.2. *Let M and N be two $\mathbf{Z}\Gamma_G$ -modules, where $M(H)$ is \mathbf{Z} -torsion free for all $H \in \mathcal{F}$. Then for every $n > l(M)$, the groups $\mathrm{Ext}_{\mathbf{Z}\Gamma_G}^n(M, N)$ are finite abelian, with exponent dividing some power of $|G|$.*

Proof. This follows from the Lemma 4.1, Proposition 2.8, and the corresponding result for modules over finite groups. \square

Note that the Ext-groups in lower dimensions are not finite in general. But, it is still true in all dimensions that the Ext-groups over $\mathbf{Z}\Gamma_G$ vanish if and only if they vanish over $\mathbf{Z}_{(p)}\Gamma_G$, for all primes p . To see this, we note that tensoring over \mathbf{Z} with $\mathbf{Z}_{(p)}$ preserves exactness, and hence

$$(4.3) \quad \mathrm{Ext}_{\mathbf{Z}\Gamma_G}^n(M, N) \otimes_{\mathbf{Z}} \mathbf{Z}_{(p)} \cong \mathrm{Ext}_{\mathbf{Z}_{(p)}\Gamma_G}^n(M \otimes_{\mathbf{Z}} \mathbf{Z}_{(p)}, N \otimes_{\mathbf{Z}} \mathbf{Z}_{(p)}).$$

We also have the following:

Proposition 4.4. *Let M and N be two $\mathbf{Z}\Gamma_G$ -modules, where $M(H)$ is \mathbf{Z} -torsion free for all $H \in \mathcal{F}$. Then, for every $n > l(M)$, there is an isomorphism*

$$\mathrm{Ext}_{\mathbf{Z}\Gamma_G}^n(M, N) \cong \bigoplus_{p \mid |G|} \mathrm{Ext}_{\mathbf{Z}_{(p)}\Gamma_G}^n(M_p, N_p)$$

where $M_p = \mathbf{Z}_{(p)} \otimes_{\mathbf{Z}} M$ and $N_p = \mathbf{Z}_{(p)} \otimes_{\mathbf{Z}} N$.

Proof. From Proposition 4.2 we know that $\mathrm{Ext}_{\mathbf{Z}\Gamma_G}^n(M, N)$ is a finite abelian group with exponent dividing some power of $|G|$, when $n > l(M)$. Now the flatness of $\mathbf{Z}_{(p)}$ over \mathbf{Z} implies as above that $\mathrm{Ext}_{\mathbf{Z}\Gamma_G}^n(M, N)$ is the direct sum of its p -localizations, for all $p \mid |G|$. We then apply the isomorphism (4.3). \square

To complete the proof of Proposition 3.13, we also need the following standard result in homological algebra (see [7, Chap. VI, 2.1] for the case of modules over rings):

Proposition 4.5. *A right $R\Gamma_G$ -module M admits a finite projective resolution if and only if there exists an integer $\ell_0 \geq 0$ such that $\text{Ext}_{R\Gamma_G}^n(M, N) = 0$, for all $n > \ell_0$ and all right $R\Gamma_G$ -modules N .*

Proof. If M admits a finite projective resolution of length k , then $\text{Ext}_{R\Gamma_G}^n(M, N) = 0$ for $n > k$ and any $R\Gamma_G$ -module N . Conversely, if $\text{Ext}_{R\Gamma_G}^n(M, N) = 0$ for $n > \ell_0$ and any N , then consider the kernel Z_m of the boundary map $\partial_m : P_m \rightarrow P_{m-1}$ in the projective resolution \mathbf{P} of M . It follows that

$$\text{Ext}_{R\Gamma_G}^1(Z_m, N) \cong \text{Ext}_{R\Gamma_G}^{m+2}(M, N) = 0$$

for any $R\Gamma_G$ -module N , provided $m+2 > \ell_0$, and so Z_m is projective if we take $m = \ell_0 - 1$. This gives a finite projective resolution of length ℓ_0 over $R\Gamma_G$. \square

We now recall the definition of a Mackey functor (following Dress [13]). Let G be a finite group and $\mathcal{D}(G)$ denote the Dress category of finite G -sets and G -maps. A bivariate functor

$$M = (M^*, M_*): \mathcal{D}(G) \rightarrow R\text{-Mod}$$

consists of a contravariant functor

$$M^*: \mathcal{D}(G) \rightarrow R\text{-Mod}$$

and a covariant functor

$$M_*: \mathcal{D}(G) \rightarrow R\text{-Mod}.$$

The functors are assumed to coincide on objects. Therefore, we write $M(S) = M_*(S) = M^*(S)$ for a finite G -set S . If $f: S \rightarrow T$ is a morphism, we often use the notation $f_* = M_*(f)$ and $f^* = M^*(f)$. If $S = G/H$ and $T = G/K$ with $H \leq K$ and $f: G/H \rightarrow G/K$ is given by $f(eH) = eK$, then we use the notation $f_* = \text{Ind}_H^K$ and $f^* = \text{Res}_H^K$.

Definition 4.6 (Dress [13]). A bivariate functor is called a *Mackey functor* if it has the following properties:

(M1) For each pullback diagram

$$\begin{array}{ccc} X & \xrightarrow{h} & Y \\ g \downarrow & & \downarrow k \\ S & \xrightarrow{f} & T \end{array}$$

of finite G -sets, we have $h_* \circ g^* = k^* \circ f_*$.

(M2) The two embeddings $S \rightarrow S \sqcup T \leftarrow T$ into the disjoint union define an isomorphism $M^*(S \sqcup T) \cong M^*(S) \oplus M^*(T)$.

Remark 4.7. There is a functor $\text{Or}(G) \rightarrow \mathcal{D}(G)$ defined on objects by $H \mapsto G/H$ for every subgroup $H \leq G$, and as the identity on morphism sets. By composition, any contravariant functor $\mathcal{D}(G) \rightarrow R\text{-Mod}$ gives a right $R\Gamma_G$ -module, with respect to any given family of subgroups \mathcal{F} of G .

In the statement of Theorem 4.11 we will use the examples $R[S^?]: \mathcal{D}(G) \rightarrow R\text{-Mod}$, defined in (3.4) for any finite G -set S .

The following example and lemma will be used in the proof of Theorem 5.9.

Example 4.8. Let $Q \in \mathcal{F}$ and let V be a right $R[W_G(Q)]$ -module, where $W_G(Q) = N_G(Q)/Q$. Then we define a bivariant functor $D_Q(V): \mathcal{D}(G) \rightarrow R\text{-Mod}$ on objects by setting

$$D_Q(V)(S) = \text{Hom}_{R[W_G(Q)]}(R[S^Q], V)$$

for any finite G -set S . For any G -map $f: S \rightarrow T$ we have a $W_G(Q)$ -map $f^Q: S^Q \rightarrow T^Q$, which induces a homomorphism

$$f^*: \text{Hom}_{R[W_G(Q)]}(R[T^Q], V) \rightarrow \text{Hom}_{R[W_G(Q)]}(R[S^Q], V)$$

by composition. To define the covariant map f_* , let $\varphi_S: R[S^Q] \rightarrow V$ be an $R[W_G(Q)]$ -homomorphism, and define $f_*(\varphi_S) = \varphi_T$ by

$$f_*(\varphi_S)(t) = \varphi_T(t) = \sum_{s \in S^Q, f(s)=t} \varphi_S(s)$$

It is not hard to verify that $D_Q(V)$ is actually a Mackey functor. The axiom (M1) follows because the Q -fixed sets in a pull-back diagram of G -sets give again a pull-back diagram. The axiom (M2) is immediate.

Definition 4.9. For any $R\Gamma_G$ -module N , we define $DN = \sum_{Q \in \text{Iso}(\Gamma_G)} D_Q(N(Q))$ and define $j: N \rightarrow DN$ as the direct sum of the adjoints of $\text{id}: N(Q) \rightarrow N(Q)$, for each $Q \in \text{Iso}(\Gamma_G)$. Let CN denote the cokernel of j . For $k \geq 0$, define inductively $C^0N = N$ and $C^kN = C(C^{k-1}N)$, together with the induced maps $C^k \rightarrow DC^k$.

Here is a dual construction to the E-resolution given in [20, 17.13].

Lemma 4.10. *For any $R\Gamma_G$ -module N , the finite length sequence*

$$0 \rightarrow N \xrightarrow{j} DN \rightarrow DCN \rightarrow \dots \rightarrow DC^mN \rightarrow 0$$

is an exact coresolution of Mackey functors, for some $m \geq 0$.

Proof. For any $R\Gamma_G$ -module N , the map $j: N \rightarrow DN$ defined above is injective, so we have a short exact sequence

$$0 \rightarrow N \xrightarrow{j} DN \rightarrow CN \rightarrow 0.$$

Iterating the above process, we obtain

$$0 \rightarrow CN \rightarrow DCN \rightarrow C^2N \rightarrow 0$$

and so on. By splicing, we get an exact sequence, or coresolution:

$$0 \rightarrow N \xrightarrow{j} DN \rightarrow DCN \rightarrow \dots \rightarrow DC^{k-1}N \rightarrow DC^kN \rightarrow \dots$$

When N is a $R\Gamma_G$ -module of a finite length, which is the case in our situation, this coresolution has a finite length. To check this, we use the definition of $D_Q(V)$ in Example 4.8 to get

$$D_Q(V)(K) = \text{Hom}_{R[W_G(Q)]}(R[(G/K)^Q], V)$$

for any $R[W_G(Q)]$ -module V . Therefore $D_Q(V)(K)$ is only nonzero for $(Q) \leq (K)$, and at $(Q) = (K)$ the $R[W_G(K)]$ -module $D_Q(V)(K)$ is isomorphic to V , via the isomorphism

$W_G(Q) \cong W_G(K)$ induced by conjugation. This shows that the length of the module $C^k N$ is properly smaller than the length of $C^{k-1} N$ for all $k \geq 1$. \square

We will prove Theorem 5.1 by showing that $H \mapsto \text{Ext}_{R\Gamma_H}^*(M, N)$ has a cohomological Mackey functor structure which is conjugation invariant. First we describe the Mackey functor structure on $\text{Hom}_{R\Gamma_G}(M, N)$.

Theorem 4.11. *For a right $R\Gamma_G$ -module M and a Mackey functor N , let*

$$\text{Hom}_{R\Gamma_G}(M, N): \mathcal{D}(G) \rightarrow R\text{-Mod}$$

denote the function defined by $S \mapsto \text{Hom}_{R\Gamma_G}(M \otimes_R R[S^?], N)$ for any finite G -set S . Then $\text{Hom}_{R\Gamma_G}(M, N)$ inherits a Mackey functor structure.

Proof. We will first define the induction and restriction maps to see that $\text{Hom}_{R\Gamma_G}(M, N)$ is a bifunctor. For $f: S \rightarrow T$ a G -map, the *restriction* map

$$f^*: \text{Hom}_{R\Gamma_G}(M \otimes_R R[T^?], N) \rightarrow \text{Hom}_{R\Gamma_G}(M \otimes_R R[S^?], N)$$

is the composition with $M \otimes_R R[S^?] \xrightarrow{\text{id} \otimes \tilde{f}} M \otimes_R R[T^?]$ where \tilde{f} denotes is the linear extension of the map induced by f . Since the functors $R[S^?]$ satisfy axiom (M2), so does $\text{Hom}_{R\Gamma_G}(M, N)$.

For $f: S \rightarrow T$ a G -map, we define the *induction* map

$$f_*: \text{Hom}_{R\Gamma_G}(M \otimes_R R[S^?], N) \rightarrow \text{Hom}_{R\Gamma_G}(M \otimes_R R[T^?], N)$$

in the following way: let $\varphi_S: M \otimes_R R[S^?] \rightarrow N$ be given. We will describe the homomorphism $\varphi_T = f_*(\varphi_S)$.

$$\varphi_T(V)(x \otimes \alpha) = F_* \left(\varphi_S(U)(F^*(x) \otimes \beta) \right)$$

for $x \in M(V)$ and $\alpha: V \rightarrow T$, where U, β and F are given by the pull-back

$$\begin{array}{ccc} U & \xrightarrow{\beta} & S \\ F \downarrow & & \downarrow f \\ V & \xrightarrow{\alpha} & T \end{array}$$

It is easy to check that this formula for φ_T gives an $R\Gamma_G$ -homomorphism, using the assumption that N is a Mackey functor.

We need to check axiom (M1) for $\text{Hom}_{R\Gamma_G}(M, N)$. For a given pull-back square

$$\begin{array}{ccc} X & \xrightarrow{h} & Y \\ g \downarrow & & \downarrow k \\ S & \xrightarrow{f} & T \end{array}$$

we need to show that $h_* \circ g^* = k^* \circ f_*$. Let $\gamma: V \rightarrow Y$ be any G -map, and consider the extended pull-back diagram

$$\begin{array}{ccccc} U & \xrightarrow{\delta} & X & \xrightarrow{g} & S \\ F \downarrow & & h \downarrow & & \downarrow f \\ V & \xrightarrow{\gamma} & Y & \xrightarrow{k} & T \end{array}$$

The maps $\alpha = k \circ \gamma$ and $\beta = g \circ \delta$ may be used to compute $f_*(\varphi_S)$ as above, and the left-hand square may be used to compute h_* .

For any element $\varphi_S: M \otimes_R R[S^?] \rightarrow N$, we have

$$\begin{aligned} (k^* \circ f_*(\varphi_S))(V)(x \otimes \gamma) &= (f_*(\varphi_S) \circ (\text{id} \otimes k))(V)(x \otimes \gamma) \\ &= f_*(\varphi_S)(V)(x \otimes (k \circ \gamma)) \\ &= F_*(\varphi_S(U)(F^*(x) \otimes (g \circ \delta))) \end{aligned}$$

for any $x \in M(V)$ and $\gamma: V \rightarrow Y$. On the other hand,

$$\begin{aligned} (h_* \circ g^*(\varphi_S))(V)(x \otimes \gamma) &= F_*((g^*\varphi_S)(U)(F^*(x) \otimes \delta)) \\ &= F_*(\varphi_S(U)(F^*(x) \otimes (g \circ \delta))) \end{aligned}$$

for any $x \in M(V)$ and $\gamma: V \rightarrow Y$, so the formula (M1) is verified. \square

As an immediate consequence, for any subgroup $H \leq K$ the G -map $f: G/H \rightarrow G/K$ induces a restriction map

$$\text{Res}_H^K: \text{Hom}_{R\Gamma_K}(M, N) \rightarrow \text{Hom}_{R\Gamma_H}(M, N)$$

defined as the composition of the map

$$f^*: \text{Hom}_{R\Gamma_G}(M \otimes_R R[G/K^?], N) \rightarrow \text{Hom}_{R\Gamma_G}(M \otimes_R R[G/H^?], N)$$

with the ‘Shapiro’ isomorphisms:

$$\text{Hom}_{R\Gamma_G}(M \otimes_R R[G/H^?], N) \cong \text{Hom}_{R\Gamma_H}(M, N)$$

and

$$\text{Hom}_{R\Gamma_G}(M \otimes_R R[G/K^?], N) \cong \text{Hom}_{R\Gamma_K}(M, N)$$

given by [35, Cor. 2.12] and the adjointness property (compare [2, Lemma 2.8.4]). Similarly, we have the induction map

$$\text{Ind}_H^K: \text{Hom}_{R\Gamma_H}(M, N) \rightarrow \text{Hom}_{R\Gamma_K}(M, N)$$

defined by composing the Shapiro isomorphisms with f_* .

Remark 4.12. Since Res_H^G preserves projectives, we see that $P \otimes_R R[G/H^?]$ is projective over $R\Gamma_G$ whenever P is projective over $R\Gamma_G$ (check the categorical lifting property directly or apply Lemma 3.2).

Proposition 4.13. *Let \mathbf{C} be a chain complex of right $R\Gamma_G$ -modules and N be a Mackey functor. Then, the cochain complex*

$$\mathbf{C}^* = \text{Hom}_{R\Gamma_G}(\mathbf{C}, N)$$

with the differential $\delta: \text{Hom}_{R\Gamma_G}(C_i, N) \rightarrow \text{Hom}_{R\Gamma_G}(C_{i+1}, N)$ given by $\delta(\varphi) = \varphi \circ \partial$ is a cochain complex of Mackey functors.

Proof. We have seen that each $C^i = \text{Hom}_{R\Gamma_G}(C_i, N)$ is a Mackey functor by Theorem 4.11. We just need to show that the coboundary maps are Mackey functor maps. Given $f: S \rightarrow T$ we need to show the following diagram commutes:

$$\begin{array}{ccc} \text{Hom}_{R\Gamma_G}(C_i \otimes R[S^?], N) & \xrightarrow{\delta_S} & \text{Hom}_{R\Gamma_G}(C_{i+1} \otimes R[S^?], N) \\ f^* \uparrow \downarrow f_* & & f^* \uparrow \downarrow f_* \\ \text{Hom}_{R\Gamma_G}(C_i \otimes R[T^?], N) & \xrightarrow{\delta_T} & \text{Hom}_{R\Gamma_G}(C_{i+1} \otimes R[T^?], N) \end{array}$$

The proof of commutativity for f^* is easy. In this case, it follows from the commutativity of the following diagram:

$$\begin{array}{ccc} C_i \otimes R[S^?] & \xleftarrow{\partial \otimes \text{id}} & C_{i+1} \otimes R[S^?] \\ \downarrow \text{id} \otimes f & & \downarrow \text{id} \otimes f \\ C_i \otimes R[T^?] & \xleftarrow{\partial \otimes \text{id}} & C_{i+1} \otimes R[T^?] \end{array}$$

For f_* we check the commutativity directly: let $\varphi_S: C_i \otimes R[S^?] \rightarrow N$ be an $R\Gamma_G$ -map. For $x \in C_{i+1}(V)$ and $\alpha: V \rightarrow T$, we have

$$\begin{aligned} [(\delta_T \circ f_*)\varphi_S](x \otimes \alpha) &= (f_*\varphi_S)(\partial x \otimes \alpha) \\ &= F_*[\varphi_S(F^*(\partial x) \otimes \beta)] \end{aligned}$$

where

$$\begin{array}{ccc} U & \xrightarrow{\beta} & S \\ F \downarrow & & \downarrow f \\ V & \xrightarrow{\alpha} & T \end{array}$$

on the other hand,

$$\begin{aligned} [(f_* \circ \delta_S)\varphi_S](x \otimes \alpha) &= F_*[(\delta_S\varphi_S)(F^*(x) \otimes \beta)] \\ &= F_*[\varphi_S \circ (\partial \otimes \text{id})(F^*(x) \otimes \beta)] \\ &= F_*[\varphi_S(\partial F^*(x) \otimes \beta)] \end{aligned}$$

since $\partial F^* = F^*\partial$, we are done. □

Corollary 4.14. *Let M be an $R\Gamma_G$ -module and N be a Mackey functor. Then,*

$$\text{Ext}_{R\Gamma_G}^*(M, N)$$

has a Mackey functor structure. As a Mackey functor $\text{Ext}_{R\Gamma_{\mathcal{F}}}^*(M, N)$ is equal to the homology of the cochain complex of Mackey functors $\text{Hom}_{R\Gamma_{\mathcal{F}}}(\mathbf{P}, N)$ where \mathbf{P} is a projective resolution of M as an $R\Gamma_G$ -module.

Proof. To compute the Ext-groups, note that $S \mapsto \mathbf{P} \otimes_R R[S^?]$ is a projective resolution of the module $S \mapsto M \otimes_R R[S^?]$, for every finite G -set S . \square

Remark 4.15. It follows that a version of the Eckmann-Shapiro isomorphism

$$\text{Ext}_{R\Gamma_G}^*(M \otimes R[G/H^?], N) \cong \text{Ext}_{R\Gamma_H}^*(\text{Res}_H^G M, \text{Res}_H^G N)$$

holds for the Ext-groups over the orbit category (compare [2, 2.8.4]).

Remark 4.16. If N is a Green module over a Green ring \mathcal{G} , then the Mackey functor $\text{Ext}_{R\Gamma_{\mathcal{F}}}^*(M, N)$ also inherits a Green module structure over \mathcal{G} . The basic formula is a pairing

$$\mathcal{G}(S) \times \text{Hom}_{R\Gamma_{\mathcal{F}}}(M \otimes_R R[S^?], N) \rightarrow \text{Hom}_{R\Gamma_{\mathcal{F}}}(M \otimes_R R[S^?], N)$$

induced by the Green module pairing $\mathcal{G} \times N \rightarrow N$. For any $z \in \mathcal{G}(S)$, $x \in M(V)$, and $\alpha: V \rightarrow S$, we define

$$(z \cdot \varphi_S)(V)(x \otimes \alpha) = \alpha^*(z) \cdot \varphi_S(V)(x \otimes \alpha)$$

for any $\varphi_S(V): M(V) \otimes_R R \text{Mor}(S, V) \rightarrow N(V)$. The check that this pairing gives a Green module structure is left to the reader. \square

5. THE PROOF OF THEOREM C

The main purpose of this section to prove the following theorem.

Theorem 5.1. *Let G be a finite group, $R = \mathbf{Z}_{(p)}$, and \mathcal{F} be a family of subgroups in G . Suppose $H \leq G$ controls p -fusion in G . Then,*

$$\text{Res}_H^G: \text{Ext}_{R\Gamma_G}^n(M, N) \rightarrow \text{Ext}_{R\Gamma_H}^n(\text{Res}_H^G M, \text{Res}_H^G N)$$

is an isomorphism for $n \geq 0$, provided that M is an $R\Gamma_G$ -module and N is a cohomological Mackey functor satisfying the condition that $C_G(Q)$ acts trivially on $N(Q)$ and $M(Q)$ for all p -subgroups $Q \leq H$, with $Q \in \mathcal{F}$.

Certain Mackey functors (called *cohomological*) are computable by restriction to the p -Sylow subgroups and the conjugation action of G (see [7, Chap. XII, §10], [19]).

If $H \leq G$ is a subgroup, and $n \in N_G(H)$ then the G -map $f: G/H \rightarrow G/H$ defined by $f(eH) = nH$ has an associated conjugation homomorphism $c_n(h) = n^{-1}hn \in H$, for all $h \in H$. For an arbitrary $R\Gamma_G$ -module M , the induced maps f^* need not be the identity on $M(G/H)$ even if $c_n = \text{id}$ (e.g. if $n \in C_G(H)$).

Definition 5.2. We say a Mackey functor is *cohomological* (over \mathcal{F}) if

$$\text{Ind}_H^K \text{Res}_H^K(u) = |K: H| \cdot u$$

for all $u \in M(K)$, and all $H \leq K$ (for all $K \in \mathcal{F}$). An $R\Gamma_G$ -module M with respect to a family \mathcal{F} is called *conjugation invariant* if $C_G(Q)$ acts trivially on $M(Q)$ for all $Q \in \mathcal{F}$. A Mackey functor is called *conjugation invariant* if it is conjugation invariant as a functor over the corresponding orbit category.

The following lemma will be used in the proof of Theorem 5.9.

Lemma 5.3. *Let $Q \in \mathcal{F}$ and let V be a right $R[W_Q(Q)]$ -module. If \mathcal{F} is a family of p -subgroups, and $R = \mathbb{F}_p$, then $D_Q(V): \mathcal{D}(G) \rightarrow R\text{-Mod}$ is a cohomological Mackey functor over \mathcal{F} . If $C_G(Q)$ acts trivially on V , then $D_Q(V)$ is conjugation invariant.*

Proof. Since all subgroups in \mathcal{F} are p -groups, for the first part we only need to show that the composite $\text{Ind}_H^K \text{Res}_H^K(u) = p \cdot u$, for $K \in \mathcal{F}$ and $H \leq K$ a normal of index p .

Let $f: G/H \rightarrow G/K$ be the G -map given by $gH \mapsto gK$. Consider the induced map $f^Q: (G/H)^Q \rightarrow (G/K)^Q$. Take $t \in (G/K)^Q$. If there is no $s \in (G/H)^Q$ such that $f(s) = t$, then the transfer is trivially zero. Suppose that there is at least one element $s = gH$ which is fixed by Q and maps to $t = gK$. Let k_1, \dots, k_p be coset representatives of H in K . Since k_i normalizes H , the element $gk_iH \in (G/H)^Q$ for each i . Therefore, there are exactly p different $s \in (G/H)^Q$ that map to t . It follows that $f_* \circ f^*$ is multiplication by p , as required. Since we are working here over the finite field \mathbb{F}_p , all the composites $f_* \circ f^* = 0$.

We now show that $D_Q(V)$ is conjugation invariant if $C_G(Q)$ acts trivially on V . In other words, we claim that for all $K \in \mathcal{F}$, the centralizer $C_G(K)$ acts trivially on $\text{Hom}_{R[W_G(Q)]}(R[G/K]^Q, V)$. Consider the way the action is defined: let $c \in C_G(K)$ and $\varphi: R[G/K]^Q \rightarrow V$ be an $R[W_G(Q)]$ -map. Then $(c\varphi)(gK) = \varphi(gcK)$. On the other hand since $gK \in (G/K)^Q$, we have $Q^g \leq K$. So, c centralizes Q^g . This means gcg^{-1} centralizes Q and hence acts trivially on V . This gives

$$\varphi(gcK) = \varphi(gcg^{-1}gK) = gcg^{-1}\varphi(gK) = \varphi(gK)$$

Therefore $(c\varphi)(gK) = \varphi(gK)$ for all gK . This shows that $c \in C_G(K)$ acts as the identity on $\text{Hom}_{R[W_G(Q)]}(R[G/K]^Q, V)$. \square

The cohomological and conjugation properties are inherited by the Ext-functors.

Proposition 5.4. *Let M and N be $R\Gamma_G$ -modules relative to some family \mathcal{F} .*

- (i) *If N is a cohomological Mackey functor over \mathcal{F} , then $\text{Ext}_{R\Gamma_{\mathcal{F}}}^*(M, N)$ is a cohomological Mackey functor over all subgroups $H \leq G$.*
- (ii) *If both M and N are conjugation invariant with respect to \mathcal{F} , then $\text{Ext}_{R\Gamma_{\mathcal{F}}}^*(M, N)$ is conjugation invariant with respect to all subgroups $H \leq G$.*

Proof. We have seen that for $f: S \rightarrow T$, the induced maps

$$\text{Hom}_{R\Gamma_G}(M \otimes R[S^?], N) \begin{array}{c} \xrightarrow{f_*} \\ \xleftarrow{f^*} \end{array} \text{Hom}_{R\Gamma_G}(M \otimes R[T^?], N)$$

satisfy the property that

$$\begin{aligned} [(f_* \circ f^*)\varphi_T](V)(x \otimes \alpha) &= F_*[f^*(\varphi_T)(U)(F^*(x) \otimes \beta)] \\ &= F_*[\varphi_T(U)(F^*(x) \otimes (f \circ \beta))] \\ &= F_*[\varphi_T(U)(F^*(x) \otimes (\alpha \circ F))] \\ &= (F_* \circ F^*)[\varphi_T(V)(x \otimes \alpha)] \end{aligned}$$

for all $x \in M(V)$ and $\alpha: V \rightarrow T$. In the last equality we used the invariance of φ_T with respect to the G -map $F: U \rightarrow V$ (our notation comes from the definition of f_* above). Hence, if $f: G/H \rightarrow G/K$ and $F_* \circ F^*$ is multiplication by $|K: H|$ (this follows from a count of double cosets), then $f_* \circ f^*$ is also multiplication by $|K: H|$.

Let M and N be conjugation invariant right $R\Gamma_G$ -modules, and let \mathbf{P} be a projective resolution of M over $R\Gamma_G$. To show that $\text{Ext}_{R\Gamma_G}^*(M, N)$ is conjugation invariant, it is enough to show that the chain map induced by the conjugation action on $\text{Hom}_{R\Gamma_G}(\mathbf{P}, N)$ is homotopy equivalent to the identity. We remark that the action of an element $c \in C_G(H)$ gives an automorphism $J_c: \text{Or}_{\mathcal{F}} H \rightarrow \text{Or}_{\mathcal{F}} H$, and induces an $R\Gamma_H$ -module chain map $\mathbf{P}(J_c): \text{Res}_H^G(\mathbf{P}) \rightarrow \text{Res}_H^G(\mathbf{P})$.

If $f: G/H \rightarrow G/H$ is given by $eH \mapsto cH$ where $c \in C_G(H)$, then for each degree i ,

$$f_i^*: \text{Hom}_{R\Gamma_G}(P_i \otimes R[G/H^?], N) \rightarrow \text{Hom}_{R\Gamma_G}(P_i \otimes R[G/H^?], N)$$

is given by

$$f_i^*(\varphi_S)(U)(x \otimes \alpha) = \varphi_S(U)(x \otimes f \circ \alpha)$$

where $S = G/H$, $x \in P_i(U)$, and $\alpha: U \rightarrow G/H$ is a G -map. In other words, $f_i^* = \text{Hom}_{R\Gamma_G}(\lambda_i, \text{id})$, where $\lambda_i(x \otimes \alpha) = x \otimes f \circ \alpha$ defines a chain map

$$\lambda: \mathbf{P} \otimes R[G/H^?] \rightarrow \mathbf{P} \otimes R[G/H^?].$$

We may assume that $U = G/K$ with $K \in \mathcal{F}$. Let $\alpha(eK) = gH$. The conjugation action of $c \in C_G(H)$ on $M(U)$ or $N(U)$ is given by the G -map $F: G/K \rightarrow G/K$, where $F(eK) = gcg^{-1}K$ and $f \circ \alpha = \alpha \circ F$. We remark that $z := gcg^{-1} \in C_G(K)$, since $K \subseteq gHg^{-1}$, and that $\mathbf{P}^*(F) = \mathbf{P}(J_z)(K)$. Notice that

$$f_i^*(\varphi_S)(U)(x \otimes \alpha) = (\varphi_S(U)(x \cdot P_i^*(F)^{-1} \otimes \alpha)) \cdot N^*(F),$$

showing that the maps f_i^* are just given by the natural action maps of c on the domain and range of the Hom. Now observe that

$$\mathbf{P}(J_z): \text{Res}_K^G(\mathbf{P}) \rightarrow \text{Res}_K^G(\mathbf{P})$$

is a chain map lifting $M(J_z): \text{Res}_K^G(M) \rightarrow \text{Res}_K^G(M)$. Since M is conjugation invariant, it follows that $\mathbf{P}(J_z) \simeq \text{id}$ by uniqueness (up to chain homotopy) of lifting in projective resolutions. Therefore $\lambda_1 := \lambda \circ (\mathbf{P}^*(F) \otimes \text{id}) \simeq \lambda$, and $f^* \simeq \text{Hom}(\lambda_1, \text{id})$. But for all $x \in P_i(U)$, we have

$$\text{Hom}(\lambda_1, \text{id})(\varphi_S)(U)(x \otimes \alpha) = \varphi_S(U)(x \cdot P_i^*(F) \otimes f \circ \alpha) = (\varphi_S(U)(x \otimes \alpha)) \cdot N^*(F),$$

and hence $f^*(\varphi_S) \simeq \varphi_S$, by the conjugation invariance of N . \square

Definition 5.5. For any subgroup $H \leq G$, and any $R\Gamma_G$ -modules M and N , an element $\alpha \in \text{Ext}_{R\Gamma_H}^n(M, N)$ is called *stable* with respect to G provided that

$$\text{Res}_{H \cap {}^g H}^H(\alpha) = \text{Res}_{H \cap {}^g H}^{{}^g H} c_H^g(\alpha)$$

for any $g \in G$. The map c_H^g is the induced map f_* where $f: G/H \rightarrow G/{}^g H$ is the G -map given by $xH \rightarrow xg^{-1}(gHg^{-1})$.

Theorem 5.6. *Let $R = \mathbf{Z}_{(p)}$ and G be a finite group. For a right $R\Gamma_G$ -module M and a cohomological Mackey functor $N: \mathcal{D}(G) \rightarrow R\text{-Mod}$, the restriction map*

$$\text{Res}_P^G: \text{Ext}_{R\Gamma_G}^n(M, N) \rightarrow \text{Ext}_{R\Gamma_P}^n(M, N)$$

is an isomorphism for $n \geq 0$ onto the stable elements, for any p -Sylow subgroup $P \leq G$.

Proof. By Proposition 5.4(i), $\text{Ext}_{R\Gamma_G}^n(M, N)$ is a cohomological Mackey functor. Now the result follows (as in [33, 2.2]) from the stable element method of Cartan and Eilenberg [7, Chap. XII, 10.1]. \square

Remark 5.7. Since $\text{Ext}_{R\Gamma_G}^*(M, N)$ is a cohomological Mackey functor, it is a Green module over the trivial module \underline{R} , considered as a Green ring by defining $\text{Ind}_H^K: \underline{R}(G/H) \rightarrow \underline{R}(G/K)$ to be multiplication by $|K : H|$ (see [19, Ex. 2.9]). It follows that $\text{Ext}_{R\Gamma_G}^*(M, N)$ is computable in the sense of Dress in terms of the p -Sylow subgroups (see [16, Ex. 5.9]).

The proof of Theorem 5.1. Let $R = \mathbf{Z}_{(p)}$ and G be a finite group. Let $H \leq G$ be a subgroup which controls p -fusion in G . For any cohomological Mackey functor F , the restriction map Res_P^G maps surjectively to the stable elements in $F(P)$, for any p -Sylow subgroup $P \leq G$. If H controls p -fusion in G , and F is conjugation invariant, then all elements in $F(H)$ are stable and

$$\text{Res}_H^G: F(G) \xrightarrow{\sim} F(H)$$

is an isomorphism. This follows by a standard argument used to prove one direction of Mislin's theorem in group cohomology (see, for example, Symonds [33, Theorem 3.5] or Benson [2, Proposition 3.8.4]). We apply Proposition 5.4 and this remark to the cohomological Mackey functor $F = \text{Ext}_{R\Gamma_G}^n(M, N)$, and the proof is complete. \square

In the next section we will need a variation of this result.

Definition 5.8. We say the N is an *atomic* right $R\Gamma_G$ -module of type $Q \in \mathcal{F}$, if $N = I_Q(N(Q))$ where I_Q is the inclusion functor introduced in Section 2.

Theorem 5.9. *Let G be a finite group, $R = \mathbf{Z}_{(p)}$, and let \mathcal{F} be a family of p -subgroups in G . Suppose $H \leq G$ controls p -fusion in G . Then, for $R\Gamma_G$ -modules M and N ,*

$$\text{Res}_H^G: \text{Ext}_{R\Gamma_G}^n(M, N) \rightarrow \text{Ext}_{R\Gamma_H}^n(\text{Res}_H^G M, \text{Res}_H^G N)$$

is an isomorphism for $n \geq 0$, provided that $C_G(Q)$ acts trivially on $M(Q)$ and $N(Q)$ for all $Q \in \mathcal{F}$.

Proof. Without loss of generality, we can assume that N is an atomic $R\Gamma_G$ -module of type Q , with trivial $C_G(Q)$ -action on $N(Q)$. This follows from the 5-lemma (using the filtration of N in [20, 16.8]).

Furthermore, we may also assume that $N(Q)$ is R -torsion free. To see this, observe that as an $N_G(Q)/QC_G(Q)$ -module, $N(Q)$ fits into a short exact sequence $0 \rightarrow L \rightarrow F \rightarrow N(Q) \rightarrow 0$, where F is a free $N_G(Q)/QC_G(Q)$ -module. By taking inflations of these modules, we can consider the sequence as a sequence of $N_G(Q)/Q$ -modules and apply the functor I_Q . This shows that N fits into a sequence $0 \rightarrow N'' \rightarrow N' \rightarrow N \rightarrow 0$, where

both N' and N'' are conjugation invariant and atomic, with an R -torsion free module at Q .

Now let $N_p = N \otimes \mathbb{F}_p = N/pN$. By Lemma 4.10 we have a finite length coresolution

$$(5.10) \quad 0 \rightarrow N_p \rightarrow DN_p \rightarrow DCN_p \rightarrow \cdots \rightarrow DC^m N_p \rightarrow 0$$

for some $m \geq 0$. Since \mathcal{F} is family of p -groups, Lemma 5.3 shows that the Mackey functors $DC^i N_p$ are cohomological over \mathcal{F} and conjugation invariant, for $0 \leq i \leq m$.

We can apply the functors $\text{Ext}_{R\Gamma_i}^*(M, -)$ to the coresolution (5.10). By Proposition 5.4, the Mackey functors $\text{Ext}_{R\Gamma_i}^*(M, DC^i N_p)$ are also cohomological and conjugation invariant. Therefore

$$\text{Res}_H^G: \text{Ext}_{R\Gamma_G}^*(M, N_p) \rightarrow \text{Ext}_{R\Gamma_H}^*(M, N_p)$$

is an isomorphism by Theorem 5.1 and the 5-lemma (using the coresolution). Furthermore, since $N(Q)$ is R -torsion free, we have a short exact sequence

$$0 \rightarrow N/p^{k-1} \rightarrow N/p^k \rightarrow N/p \rightarrow 0,$$

for every $k \geq 1$, and hence by “dévissage” we conclude that

$$(5.11) \quad \text{Res}_H^G: \text{Ext}_{R\Gamma_G}^*(M, N/p^k) \xrightarrow{\cong} \text{Ext}_{R\Gamma_H}^*(M, N/p^k)$$

is an isomorphism, for every $k \geq 1$. To finish the proof it is enough to show that

$$\text{Res}_H^G: \text{Ext}_{R\Gamma_G}^*(M, N) \otimes \widehat{\mathbf{Z}}_p \rightarrow \text{Ext}_{R\Gamma_H}^*(M, N) \otimes \widehat{\mathbf{Z}}_p$$

is an isomorphism. However, for \mathbf{P} any projective resolution of M over $R\Gamma_G$, the complex

$$\text{Hom}_{R\Gamma_G}(\mathbf{P}, N/p^k) = \text{Hom}_{R\Gamma_G}(\mathbf{P}, N) \otimes \mathbf{Z}/p^k$$

is a cochain complex of finitely-generated R -modules. By the universal coefficient theorem in cohomology [30, p. 246], we have an exact sequence

$$0 \rightarrow \text{Ext}_{R\Gamma_G}^n(M, N) \otimes \mathbf{Z}/p^k \rightarrow \text{Ext}_{R\Gamma_G}^n(M, N/p^k) \rightarrow \text{Tor}_1^R(\text{Ext}_{R\Gamma_G}^{n+1}(M, N), \mathbf{Z}/p^k) \rightarrow 0.$$

Since $\widehat{\mathbf{Z}}_p = \varprojlim \mathbf{Z}/p^k$ and the inverse limit functor is left exact, we obtain an exact sequence

$$0 \rightarrow \text{Ext}_{R\Gamma_G}^n(M, N) \otimes \widehat{\mathbf{Z}}_p \rightarrow \varprojlim \text{Ext}_{R\Gamma_G}^n(M, N/p^k) \rightarrow \varprojlim \text{Tor}_1^R(\text{Ext}_{R\Gamma_G}^{n+1}(M, N), \mathbf{Z}/p^k).$$

Now we compare this sequence via Res_H^G to the corresponding sequence for the subgroup H , and use the dévissage isomorphisms (5.11) on the middle term. This shows immediately that Res_H^G is injective on the first term, for all $n \geq 0$. Since the functor Tor_1^R is left exact, we get Res_H^G injective on the third term as well. But now a diagram chase shows that Res_H^G is surjective on the first term. \square

6. CHAIN COMPLEXES OVER ORBIT CATEGORIES

In this section, we prove some theorems about chain complexes over orbit categories. In particular, Proposition 6.8, Proposition 6.4, and Theorem 6.7 will be used in the proof of Theorem A (see Section 9). Most of the results follow from Dold’s theory of algebraic Postnikov systems [11].

As before, G denote a finite group and \mathcal{F} denote a family of subgroups of G . Throughout this section $\Gamma_G = \text{Or}_{\mathcal{F}} G$ and R is a commutative ring. For chain complexes \mathbf{C} and \mathbf{D} ,

the notation $\mathbf{C} \simeq \mathbf{D}$ always means \mathbf{C} is chain homotopy equivalent to \mathbf{D} . For chain isomorphism the standard notation is $\mathbf{C} \cong \mathbf{D}$. When we say \mathbf{C} is a projective chain complex, we mean it is a chain complex of projective modules (which also means that it is projective in the category of chain complexes). A chain complex \mathbf{C} is *positive* if $C_i = 0$ for $i < 0$.

We say that a chain complex \mathbf{C} over $R\Gamma_G$ has *finite homological dimension* (or $\text{hdim } \mathbf{C}$ is finite) if \mathbf{C} is positive, and there exists an integer n such that $H_i(\mathbf{C}) = 0$ for $i > n$. A chain complex \mathbf{C} is *finite* if \mathbf{C} is positive, and there exists an integer n such that $C_i = 0$ for $i > n$. We start with a well known observation about chain complexes.

Lemma 6.1. *Let \mathbf{C} be a projective chain complex of $R\Gamma_G$ -modules which has finite homological dimension. Then, \mathbf{C} is homotopy equivalent to a finite projective chain complex if and only if there is an integer n such that*

$$\text{Ext}_{R\Gamma_G}^i(\mathbf{C}, M) = 0 \text{ for } i > n,$$

for all $R\Gamma_G$ -modules M .

Proof. See Cartan-Eilenberg [7, Chap. XVII, 1.4] for chain complexes over rings. A similar argument as in Proposition 4.5 gives the result over the orbit category. \square

Proposition 6.2. *Let \mathbf{C} be a projective chain complex of $\mathbf{Z}\Gamma_G$ -modules which has a finite homological dimension. Suppose that $\mathbf{Z}_{(p)} \otimes_{\mathbf{Z}} \mathbf{C}$ is chain homotopy equivalent to a finite projective chain complex for all $p \mid |G|$. Then, \mathbf{C} is chain homotopy equivalent to a finite projective complex.*

Proof. Let M be an $R\Gamma_G$ -module. Consider the hypercohomology spectral sequence (see [3, 3.4.3]):

$$E_2^{s,t} = \text{Ext}_{\mathbf{Z}\Gamma_G}^s(H_t(\mathbf{C}), M)$$

which converges to $\text{Ext}_{\mathbf{Z}\Gamma_G}^*(\mathbf{C}, M)$. Since \mathbf{C} has finite homological dimension, for all $i > (l(\Gamma_G) + \text{hdim } \mathbf{C})$, the group $\text{Ext}_{\mathbf{Z}}^i(\mathbf{C}, M)$ is a finite abelian group with exponent dividing a power of $|G|$. Here $l(\Gamma_G)$ is the length of the orbit category, as defined in §2D, and $\text{hdim } \mathbf{C}$ denotes the largest integer n such that $H_n(\mathbf{C}) \neq 0$.

In particular, there is an integer k , independent from M , such that

$$\text{Ext}_{\mathbf{Z}\Gamma_G}^i(\mathbf{C}, M) \cong \bigoplus_{p \mid |G|} \text{Ext}_{\mathbf{Z}_{(p)}\Gamma_G}^i(\mathbf{Z}_{(p)} \otimes_{\mathbf{Z}} \mathbf{C}, M_p)$$

for all $i > k$. Here $M_p = \mathbf{Z}_{(p)} \otimes_{\mathbf{Z}} M$. Now, since $\mathbf{Z}_{(p)} \otimes_{\mathbf{Z}} \mathbf{C}$ is homotopy equivalent to a finite projective complex for all $p \mid |G|$, there is an n such that

$$\text{Ext}_{\mathbf{Z}\Gamma_G}^i(\mathbf{C}, M) = 0$$

for all $i > n$ and for all M . The result follows from the previous lemma. \square

A chain complex version of Rim's theorem also holds.

Proposition 6.3. *Let $R = \mathbf{Z}_{(p)}$ and \mathbf{C} be a projective chain complex over $R\Gamma_G$ with finite homological dimension. Assume that \mathcal{F} is a family of p -subgroups. Then, \mathbf{C} is homotopy equivalent to a finite projective complex if and only if $\text{Res}_P^G \mathbf{C}$ is homotopy equivalent to a finite projective complex for any p -Sylow subgroup P of G .*

Proof. One direction is clear (and holds without assumption on the family \mathcal{F}). Conversely, suppose that $\text{Res}_P^G \mathbf{C}$ is homotopy equivalent to a projective complex with $\text{hdim} = l$. Let n be an integer bigger than both l and $\text{hdim} \mathbf{C}$. Consider

$$\cdots \longrightarrow \text{Res}_P^G C_{n+1} \longrightarrow \text{Res}_P^G C_n \xrightarrow{\partial_n} \text{Res}_P^G C_{n-1} \longrightarrow \cdots \longrightarrow \text{Res}_P^G C_0 \longrightarrow 0.$$

We have

$$\text{Ext}_{R\Gamma_P}^1(\text{Res}_P^G \text{im}(\partial_n), M) \cong \text{Ext}_{R\Gamma_P}^{n+1}(\text{Res}_P^G \mathbf{C}, M) = 0,$$

for every $R\Gamma_P$ -module M . This gives that $\text{Res}_P^G \text{im}(\partial_n)$ is projective. By Theorem 3.9, we obtain that $\text{im}(\partial_n)$ has finite projective resolution. Thus, \mathbf{C} is chain homotopy equivalent to a finite projective complex. \square

We also prove a chain complex version of Theorem 5.9. Recall the definition of *conjugation invariant* $R\Gamma_G$ -modules given in (5.2).

Proposition 6.4. *Let G be a finite group, and \mathcal{F} be a family of p -subgroups in G . Suppose $H \leq G$ controls p -fusion in G and $R = \mathbf{Z}_{(p)}$. Let \mathbf{C}^H be a positive projective chain complex of $R\Gamma_H$ -modules such that the homology groups $H_i(\mathbf{C}^H)$ are conjugation invariant right $R\Gamma_H$ -modules, for every $i \geq 0$. Then, the following holds:*

- (i) *There exists a positive projective chain complex \mathbf{C}^G of $R\Gamma_G$ -modules such that $\text{Res}_H^G \mathbf{C}^G$ is homotopy equivalent to \mathbf{C}^H .*
- (ii) *If \mathbf{C}^H is homotopy equivalent to a finite projective complex, then \mathbf{C}^G is also homotopy equivalent to a finite complex.*

For the proof we will need the theory of algebraic Postnikov systems due to Dold [11, §7]. According to this theory, given a positive projective chain complex \mathbf{C} , there is a sequence of positive projective chain complexes $\mathbf{C}(i)$ indexed by positive integers such that $f: \mathbf{C} \rightarrow \mathbf{C}(i)$ induces a homology isomorphism for dimensions $\leq i$. Moreover, there is a tower of maps

$$\begin{array}{ccccc} & & \mathbf{C}(i) & & \\ & \nearrow & \downarrow & & \\ & & \mathbf{C}(i-1) & \xrightarrow{\alpha_i} & \Sigma^{i+1} \mathbf{P}(H_i) \\ & \nearrow & \vdots & & \\ \mathbf{C} & \longrightarrow & \mathbf{C}(1) & \xrightarrow{\alpha_2} & \Sigma^3 \mathbf{P}(H_2) \\ & \searrow & \downarrow & & \\ & & \mathbf{C}(0) & \xrightarrow{\alpha_1} & \Sigma^2 \mathbf{P}(H_1) \end{array}$$

such that $\mathbf{C}(i) = \Sigma^{-1} \mathbf{C}(\alpha_i)$, where $\mathbf{C}(\alpha_i)$ denotes the algebraic mapping cone of α_i , and $\mathbf{P}(H_i)$ denotes a projective resolution of the homology module H_i .

Recall that the algebraic mapping cone of a chain map $f: \mathbf{C} \rightarrow \mathbf{D}$ is defined as the chain complex $\mathbf{C}(f) = \mathbf{D} \oplus \Sigma \mathbf{C}$ with boundary map given by $\partial(x, y) = (\partial x + f(x), \partial y)$.

Note that Σ^n is the shift operator for chain complexes which is defined by $(\Sigma^n \mathbf{C})_i = C_{i-n}$ for every integer n .

The algebraic Postnikov system has similar properties to the Postnikov system in homotopy theory. The maps $\alpha_i: \mathbf{C}(i-1) \rightarrow \Sigma^{i+1} \mathbf{P}(H_i)$ are called *k-invariants* and they are well defined up to chain homotopy equivalence. We can consider the *k-invariants* as classes in $\text{Ext}_{R\Gamma_G}^{i+1}(\mathbf{C}(i-1), H_i)$, since there is an isomorphism

$$[\mathbf{C}(i-1), \Sigma^{i+1} \mathbf{P}(H_i)] \cong \text{Ext}_{R\Gamma_G}^{i+1}(\mathbf{C}(i-1), H_i)$$

between chain homotopy classes of chain maps and the Ext-groups of chain complexes (see Dold [11] for details). The *k-invariants* $\alpha_i \in \text{Ext}_{R\Gamma_G}^{i+1}(\mathbf{C}(i-1), H_i)$ are defined inductively and they uniquely specify \mathbf{C} up to chain homotopy equivalence.

We also need a lifting result for $R\Gamma_H$ -modules.

Lemma 6.5. *Let G be a finite group, and \mathcal{F} be a family of p -subgroups in G . Suppose $H \leq G$ controls p -fusion in G . Then the restriction map $M \mapsto \text{Res}_H^G(M)$ gives a bijection between the isomorphism classes of conjugation invariant right $R\Gamma_G$ -modules and conjugation invariant right $R\Gamma_H$ -modules.*

Proof. A conjugation invariant right $R\Gamma_G$ -module M is a functor $\text{Or}_{\mathcal{F}} G \rightarrow R\text{-Mod}$ which factors through the quotient category $\text{Or}_{\mathcal{F}} G \rightarrow \text{Sub}_{\mathcal{F}} G$. Here $\text{Sub}_{\mathcal{F}} G$ has objects $K \in \mathcal{F}$ and morphisms $\text{Mor}_{\text{Sub}_{\mathcal{F}} G}(K, L) = \text{Mor}_{\text{Or}_{\mathcal{F}} G}(G/K, G/L)/C_G(K)$, where an element $c \in C_G(K)$ acts on a G -map defined by $f(eK) = gL$ via the composition $eK \mapsto cgL$ (see [21, p. 206]).

Consider the functor $F: \text{Or}_{\mathcal{F}} H \rightarrow \text{Or}_{\mathcal{F}} G$ given on objects by $H/K \mapsto G/K$ (see Section 3), and on morphisms by induced maps. First note that every object of $\text{Sub}_{\mathcal{F}} G$ is isomorphic to an object of $\text{Sub}_{\mathcal{F}} H$, since every p -subgroup of G is conjugate to a subgroup of H . In addition, F induces a bijection of morphism sets

$$\text{Mor}_{\text{Sub}_{\mathcal{F}} H}(K, L) \rightarrow \text{Mor}_{\text{Sub}_{\mathcal{F}} G}(K, L)$$

since H controls p -fusion in G . Suppose that $F(f_1) \approx F(f_2)$, where $f_1(eK) = h_1L$ and $f_2(eK) = h_2L$, for some $h_1, h_2 \in H$. By assumption, there exists $c \in C_G(K)$ such that $ch_2L = h_1L$, or $h_1^{-1}ch_2 \in L \leq H$. But this implies $c \in C_H(K)$ so $f_1 \approx f_2$ and F is injective on morphisms. Given $f: G/K \rightarrow G/L$ with $K \leq H$, $f(eK) = gL$ and $g^{-1}Kg \subseteq L \leq H$, we have $g = ch$ for some $c \in C_G(K)$ and $h \in H$, because H controls p -fusion in G . Hence $f \approx F(f_1)$, where $f_1(eK) = hL$ and F is surjective on morphisms.

Therefore the functor $F: \text{Or}_{\mathcal{F}} H \rightarrow \text{Or}_{\mathcal{F}} G$ induces an equivalence of categories

$$\bar{F}: \text{Sub}_{\mathcal{F}} H \approx \text{Sub}_{\mathcal{F}} G$$

by [24, IV.4, Theorem 1, p. 91]. □

Proof of Proposition 6.4. Part (ii) follows from Proposition 6.3, so it is enough to prove the existence of \mathbf{C}^G . By Lemma 6.5, for each $i \geq 0$ there exists a conjugation invariant right $R\Gamma_G$ -module H_i^G such that $\text{Res}_H^G(H_i^G) = H_i(\mathbf{C}^H)$.

Consider the Postnikov tower for \mathbf{C}^H . Since $\mathbf{C}^H(0) = \mathbf{P}(H_0(\mathbf{C}^H))$ there is a complex $\mathbf{C}^G(0)$ such that $\text{Res}_H^G \mathbf{C}^G(0) \simeq \mathbf{C}^H(0)$. In this case, the complex $\mathbf{C}^G(0)$ can be taken as a projective resolution of H_0^G . Now, we will show that such a lifting exists for $\mathbf{C}^H(i)$ for all

i. For this we prove a slightly stronger statement so that we can carry out an induction. We claim that the following holds for all $n \geq 0$.

- (i) $\mathbf{C}^H(n)$ lifts to a chain complex $\mathbf{C}^G(n)$
- (ii) The restriction map

$$\text{Res}_H^G: \text{Ext}_{R\Gamma_G}^*(\mathbf{C}^G(n), N) \rightarrow \text{Ext}_{R\Gamma_H}^*(\mathbf{C}^H(n), \text{Res}_H^G N)$$

is an isomorphism for all $* \geq 0$ and for every $R\Gamma_G$ -module N which is conjugation invariant.

We have already shown that $\mathbf{C}^H(0)$ lifts to $\mathbf{C}^G(0)$. For the second property, first observe that $\mathbf{C}^G(0)$ is chain homotopy equivalent to a chain complex with single module H_0^G and similarly, $\mathbf{C}^H(0) \simeq H_0^H$. So, we need to show that

$$\text{Res}_H^G: \text{Ext}_{R\Gamma_G}^*(H_0^G, H_1^G) \rightarrow \text{Ext}_{R\Gamma_H}^*(H_0^H, H_1^H)$$

is an isomorphism. This follows from Theorem 5.9, because of our assumption on homology groups.

Now, assume that both (i) and (ii) hold for $n = i - 1$. Then, take

$$\alpha_i^G \in \text{Ext}_{R\Gamma_G}^{i+1}(\mathbf{C}^G(i-1), H_i^G)$$

which corresponds to the class $\alpha_i^H \in \text{Ext}_{R\Gamma_H}^{i+1}(\mathbf{C}^H(i-1), H_i^H)$ under the isomorphism given in (ii). Let $\mathbf{C}^G(i) = \Sigma^{-1}\mathbf{C}(\alpha_i^G)$. Then, we have a short exact sequence of chain complexes

$$0 \longrightarrow \mathbf{C}^G(i) \longrightarrow \mathbf{C}^G(i-1) \xrightarrow{\alpha_i^G} \Sigma^{i+1}\mathbf{P}(H_i^G) \longrightarrow 0$$

Since $\text{Res}_H^G \alpha_i^G = \alpha_i^H$, we have $\text{Res}_H^G \mathbf{C}^G(i) \simeq \mathbf{C}^H(i)$. Now, we will show that (ii) holds for $\mathbf{C}^G(i)$. By the Five Lemma, it is enough to show that

$$\text{Res}_H^G: \text{Ext}_{R\Gamma_G}^*(\Sigma^{i+1}\mathbf{P}(H_i^G), N) \rightarrow \text{Ext}_{R\Gamma_H}^*(\Sigma^{i+1}\mathbf{P}(H_i^H), N)$$

is an isomorphism for all $* \geq 0$, and for every $R\Gamma_G$ -module N which is conjugation invariant. But, this follows from Theorem 5.9. \square

Now, we prove one of the main results of this section which allows us to glue p -local chain complexes. We first give a definition.

Definition 6.6. Let \mathbf{C} be a chain complex over $R\Gamma_G$. We say that \mathbf{C} has *homology gaps of length n* , if $H_{i+k}(\mathbf{C}) = 0$ for $0 < k < n$, whenever $H_i(\mathbf{C}) \neq 0$.

Theorem 6.7. *Let G be a finite group of order m . For each prime $p \mid m$, let $\mathbf{C}^{(p)}$ be a positive projective chain complex of $\mathbf{Z}_{(p)}\Gamma_G$ -modules. Suppose that*

- (i) $\mathbf{C}^{(p)}$ has homology gaps of length $> l(\Gamma_G)$, for all $p \mid m$.
- (ii) there exists a graded $\mathbf{Z}\Gamma_G$ -module \mathbf{H} such that $H_i(\mathbf{C}^{(p)}) \cong \mathbf{Z}_{(p)} \otimes_{\mathbf{Z}} \mathbf{H}_i$ for all $i \geq 0$, and for all $p \mid m$.

Then, there is a projective chain complex \mathbf{C} of $\mathbf{Z}\Gamma_G$ -modules such that $\mathbf{Z}_{(p)} \otimes_{\mathbf{Z}} \mathbf{C} \simeq \mathbf{C}^{(p)}$, for each prime $p \mid m$, and $H_i(\mathbf{C}) = \mathbf{H}_i$ for $i \geq 0$.

Proof. We will construct \mathbf{C} inductively. The case $i = 0$ is trivial, because in this case we can take $\mathbf{C}(0) = \mathbf{P}(\mathbf{H}_0)$. Assume now that $\mathbf{C}(i - 1)$ has been constructed in such a way that $\mathbf{Z}_{(p)} \otimes_{\mathbf{Z}} \mathbf{C}(i - 1) \simeq \mathbf{C}_{i-1}^{(p)}$ for all $p \mid m$. If $\mathbf{H}_i = 0$, then we can take $\mathbf{C}(i) = \mathbf{C}(i - 1)$ and it will satisfy the condition that $\mathbf{Z}_{(p)} \otimes_{\mathbf{Z}} \mathbf{C}(i) \simeq \mathbf{C}_i^{(p)}$. So, assume \mathbf{H}_i is nonzero. If $i + 1 > (l(\Gamma_G) + \text{hdim } \mathbf{C}(i - 1))$, then we have

$$\begin{aligned} \text{Ext}_{\mathbf{Z}\Gamma_G}^{i+1}(\mathbf{C}(i - 1), \mathbf{P}(\mathbf{H}_i)) &\cong \bigoplus_{p \mid m} \text{Ext}_{\mathbf{Z}_{(p)}\Gamma_G}^{i+1}(\mathbf{Z}_{(p)} \otimes_{\mathbf{Z}} \mathbf{C}(i - 1), \mathbf{H}_i^{(p)}) \\ &\cong \bigoplus_{p \mid m} \text{Ext}_{\mathbf{Z}_{(p)}\Gamma_G}^{i+1}(\mathbf{C}^{(p)}(i - 1), \mathbf{H}_i^{(p)}) \end{aligned}$$

where $\mathbf{H}_i^{(p)} = \mathbf{Z}_{(p)} \otimes_{\mathbf{Z}} \mathbf{H}_i$. Note that the above condition on $(i + 1)$ is satisfied since the distance between nonzero homology groups of $\mathbf{C}^{(p)}$ is bigger than $l(\Gamma_G)$. Choose $\alpha_i \in \text{Ext}_{\mathbf{Z}\Gamma_G}^{i+1}(\mathbf{C}(i - 1), \mathbf{P}(\mathbf{H}_i))$ so that under the p -localization map, α_i is mapped to the i -th k -invariant $\alpha_i^{(p)}$ of the p -local complex $\mathbf{C}^{(p)}$, for every $p \mid m$. Let $\mathbf{C}(i) = \Sigma^{-1}\mathbf{C}(\alpha_i)$. For each prime $p \mid m$, we have a diagram of the form

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbf{C}(i) & \longrightarrow & \mathbf{C}(i - 1) & \xrightarrow{\alpha_i} & \mathbf{P}(\mathbf{H}_i) & \longrightarrow & 0 \\ & & & & \downarrow \varphi_p & & \downarrow & & \\ 0 & \longrightarrow & \mathbf{C}^{(p)}(i) & \longrightarrow & \mathbf{C}^{(p)}(i - 1) & \xrightarrow{\alpha_i^{(p)}} & \mathbf{P}(\mathbf{H}_i^{(p)}) & \longrightarrow & 0 \end{array}$$

where the vertical map φ_p is given by the composition

$$\varphi_p: \mathbf{C}(i - 1) \rightarrow \mathbf{Z}_{(p)} \otimes_{\mathbf{Z}} \mathbf{C}(i - 1) \cong \mathbf{C}^{(p)}(i - 1).$$

The first map in the above composition is induced by the usual inclusion of integers into p -local integers. From this diagram, it is clear that there is a map $\mathbf{C}(i) \rightarrow \mathbf{C}^{(p)}(i)$ which induces an isomorphism on homology when it is localized at p . Thus, it gives a chain homotopy equivalence $\mathbf{Z}_{(p)} \otimes_{\mathbf{Z}} \mathbf{C}(i) \simeq \mathbf{C}^{(p)}(i)$, for $p \mid m$. This completes the proof. \square

We conclude this section with a technique (used in the proof of Theorem A) for modifying the homology of a given (finite, projective) chain complex \mathbf{C} over the orbit category. A projective resolution $\mathbf{P} \rightarrow M$ has *length* $\leq \ell$, provided that $P_i = 0$ for $i > \ell$.

Proposition 6.8. *Let Γ be an EI-category. Let $\varphi: \mathbf{H}_k \rightarrow \mathbf{H}'_k$ be an $R\Gamma$ -module homomorphism, where $\mathbf{H}_k = H_k(\mathbf{C})$. Suppose that both kernel and cokernel of φ admit finite projective resolutions of length $\leq \ell$, and that $H_{k+j} = 0$ for $1 \leq j < \ell$. Then there is a $R\Gamma$ -chain complex \mathbf{C}' such that $H_i(\mathbf{C}') = H_i(\mathbf{C})$, for $i \neq k$, and $H_k(\mathbf{C}') = \mathbf{H}'_k$.*

Proof. First suppose that φ is surjective. Let

$$0 \rightarrow P_{k+\ell} \rightarrow \cdots \rightarrow P_k \rightarrow \ker \varphi \rightarrow 0$$

be a projective resolution for $\ker \varphi$. Since \mathbf{C} is exact in the range $[k+1, k+\ell)$, we have a chain map

$$\begin{array}{ccccccccccccccc} \cdots & \longrightarrow & 0 & \longrightarrow & P_{k+\ell} & \longrightarrow & \cdots & \longrightarrow & P_{k+1} & \longrightarrow & P_k & \longrightarrow & \ker \varphi & \longrightarrow & 0 \\ & & \downarrow & & \downarrow f_{k+\ell} & & & & \downarrow f_{k+1} & & \downarrow f_k & & \downarrow \text{hook} & & \\ \cdots & \longrightarrow & C_{k+\ell+1} & \longrightarrow & C_{k+\ell} & \longrightarrow & \cdots & \longrightarrow & C_{k+1} & \longrightarrow & Z_k & \longrightarrow & \mathbf{H}_k & \longrightarrow & 0 \end{array}$$

This gives a chain map $f: \mathbf{P} \rightarrow \mathbf{C}$, where $f_k: P_k \rightarrow C_k$ is the composition of f_k with the inclusion $Z_k \subset C_k$. Let $\mathbf{C}' = \mathbf{C}(f)$ denote the mapping cone of f . The induced map

$$\ker \varphi = H_k(\mathbf{P}) \rightarrow H_k(\mathbf{C}) = \mathbf{H}_k$$

on homology is given by the inclusion, and hence $H_k(\mathbf{C}') = \mathbf{H}'_k$, with $H_i(\mathbf{C}') = H_i(\mathbf{C})$ for $i \neq k$.

Now suppose that φ is an injective map, so that

$$(6.9) \quad 0 \rightarrow \mathbf{H}_k \xrightarrow{\varphi} \mathbf{H}'_k \rightarrow \text{coker } \varphi \rightarrow 0$$

is exact. Let $\epsilon: \mathbf{P} \rightarrow \text{coker } \varphi$ be a projective resolution of $\text{coker } \varphi$ of length $\leq \ell$, indexed so that $\epsilon: P_k \rightarrow \text{coker } \varphi \rightarrow 0$. We form the pull-back

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbf{H}_k & \longrightarrow & \widehat{\mathbf{H}}_k & \longrightarrow & P_k \longrightarrow 0 \\ & & \parallel & & \downarrow \widehat{\varphi} & & \downarrow \epsilon \\ 0 & \longrightarrow & \mathbf{H}_k & \longrightarrow & \mathbf{H}'_k & \longrightarrow & \text{coker } \varphi \longrightarrow 0 \end{array}$$

of the sequence (6.9) by ϵ , and note that $\widehat{\mathbf{H}}_k \cong \mathbf{H}_k \oplus P_k$. The chain complex

$$\cdots \rightarrow C_{k+1} \rightarrow C_k \oplus P_k \rightarrow C_{k-1} \rightarrow \cdots \rightarrow C_0 \rightarrow 0$$

has homology $\widehat{\mathbf{H}}_k$ at $i = k$, and $\widehat{\varphi}: \widehat{\mathbf{H}}_k \rightarrow \mathbf{H}'_k$ is surjective. By the pull-back diagram,

$$\ker \widehat{\varphi} \cong \ker(\epsilon: P_k \rightarrow \text{coker } \varphi).$$

Since $\text{coker } \varphi$ has a projective resolution of length $\leq \ell$, it follows that $\ker \widehat{\varphi}$ has a projective resolution of length $< \ell$. Hence the assumptions needed for the surjective case hold for $\widehat{\varphi}: \widehat{\mathbf{H}}_k \rightarrow \mathbf{H}'_k$, and we are done by the argument above.

The general case is done by expressing the map $\varphi: \mathbf{H}_k \rightarrow \mathbf{H}'_k$ as the composition of a surjection and an injection. \square

7. THE FINITENESS OBSTRUCTION

Let G be a finite group and \mathcal{F} be a family of subgroups of G . The main result of this section is Theorem 7.6: given a finite projective chain complex \mathbf{C} of $\mathbf{Z}\Gamma_G$ -modules, for $\Gamma_G = \text{Or}_{\mathcal{F}} G$, we can obtain a finite free complex by taking join tensor of \mathbf{C} with itself sufficiently many times. This result is an adaptation to the orbit category of the fundamental work of Swan [32]. We first introduce some definitions, based on the material in Lück [20, §10-11].

Let Γ be an EI-category. We denote by $K_0(\mathbf{Z}\Gamma)$ the Grothendieck ring of isomorphism classes of projective $\mathbf{Z}\Gamma$ -modules and $K_0(\mathbf{Z}\Gamma, \text{free})$ denote the Grothendieck ring

of isomorphism classes of free $\mathbf{Z}\Gamma$ -modules (under direct sum $M \oplus N$ and tensor product $M \otimes_{\mathbf{Z}} N$). We have an exact sequence of abelian groups

$$0 \rightarrow K_0(\mathbf{Z}\Gamma, \text{free}) \rightarrow K_0(\mathbf{Z}\Gamma) \xrightarrow{q} \tilde{K}_0(\mathbf{Z}\Gamma) \rightarrow 0$$

defining the quotient group $\tilde{K}_0(\mathbf{Z}\Gamma)$.

Note that $K_0(\mathbf{Z}\Gamma, \text{free})$ is a subring, but not an ideal in general. This is because the tensor product of a free module with a projective module is not free in $\mathbf{Z}\Gamma$. For example, if P is a projective module which is not free, then $P \otimes_{\mathbf{Z}} \underline{\mathbf{Z}} \cong P$ is not a free $\mathbf{Z}\Gamma_G$ -module although $\underline{\mathbf{Z}}$ is free when $G \in \mathcal{F}$.

Given a finite projective chain complex of $\mathbf{Z}\Gamma$ -modules

$$\mathbf{C}: 0 \rightarrow C_n \rightarrow C_{n-1} \rightarrow \cdots \rightarrow C_1 \rightarrow C_0 \rightarrow 0$$

we define

$$\sigma(\mathbf{C}) = \sum_{i=0}^n (-1)^i [C_i] \in K_0(\mathbf{Z}\Gamma)$$

and

$$\tilde{\sigma}(\mathbf{C}) = q(\sigma(\mathbf{C})) \in \tilde{K}_0(\mathbf{Z}\Gamma).$$

The class $\tilde{\sigma}(\mathbf{C})$ is called the *finiteness obstruction* since it is the only obstruction for \mathbf{C} to be chain homotopy equivalent to a finite free chain complex.

From now on, we assume that all the chain complexes are positive and projective. As always, we assume all modules are finitely generated.

The following are standard results which show that $\tilde{\sigma}(\mathbf{C})$ is an invariant, and that it is an obstruction for finiteness.

Lemma 7.1. *If \mathbf{C} and \mathbf{D} are chain homotopy equivalent, then $\sigma(\mathbf{C}) = \sigma(\mathbf{D})$.*

Proof. See [20, 11.2]. □

Lemma 7.2. *Let \mathbf{C} and \mathbf{D} be finite chain complexes of projective $\mathbf{Z}\Gamma$ -modules. Then, $\sigma(\mathbf{C} \otimes_{\mathbf{Z}} \mathbf{D}) = \sigma(\mathbf{C}) \cdot \sigma(\mathbf{D})$.*

Proof. See [20, 11.18] and the sharper result in [20, 11.24]. □

Lemma 7.3. *Let \mathbf{C} be a finite chain complex with $\tilde{\sigma}(\mathbf{C}) = 0$. Then \mathbf{C} is chain homotopy equivalent to a finite chain complex of free $\mathbf{Z}\Gamma$ -modules.*

Proof. See Swan [32, Proposition 5.1]. □

Given two chain complexes of $R\Gamma$ -modules \mathbf{C} and \mathbf{D} , consider the corresponding augmented complexes

$$\tilde{\mathbf{C}}: \cdots \rightarrow C_2 \rightarrow C_1 \rightarrow C_0 \rightarrow \underline{R} \rightarrow 0$$

$$\tilde{\mathbf{D}}: \cdots \rightarrow D_2 \rightarrow D_1 \rightarrow D_0 \rightarrow \underline{R} \rightarrow 0$$

Taking their tensor product, we obtain a complex of the form

$$\tilde{\mathbf{C}} \otimes_R \tilde{\mathbf{D}}: \cdots \rightarrow C_1 \oplus D_1 \oplus C_0 \otimes D_0 \rightarrow C_0 \oplus D_0 \rightarrow \underline{R} \rightarrow 0.$$

Definition 7.4. We define the **join tensor**, denoted $\mathbf{C} * \mathbf{D}$, of two positive augmented chain complexes C and D by the formula

$$\widetilde{\mathbf{C} * \mathbf{D}} = \Sigma \left(\widetilde{\mathbf{C}} \otimes_R \widetilde{\mathbf{D}} \right),$$

where Σ denote the suspension of a chain complex defined by $(\Sigma \mathbf{C})_i = C_{i-1}$ for all i .

Lemma 7.5. *Let \mathbf{C} and \mathbf{D} be finite chain complexes of projective $\mathbf{Z}\Gamma$ -modules. Then, $\sigma(\mathbf{C} * \mathbf{D}) = \sigma(\mathbf{C}) + \sigma(\mathbf{D}) - \sigma(\mathbf{C}) \cdot \sigma(\mathbf{D})$.*

Proof. Note that $(\mathbf{C} * \mathbf{D})_k = C_k \oplus D_k \oplus \bigoplus_{i+j=k-1} C_i \otimes_{\mathbf{Z}} D_j$, for each $k \geq 0$. Therefore,

$$\sigma(\mathbf{C} * \mathbf{D}) = \sum_k (-1)^k [C_k] + \sum_k (-1)^k [D_k] - \sum_{i+j=k-1} (-1)^k [C_i \otimes D_j]$$

and the result follows. \square

We often express the above formula by writing

$$(1 - \sigma(\mathbf{C} * \mathbf{D})) = (1 - \sigma(\mathbf{C}))(1 - \sigma(\mathbf{D})).$$

Whenever it is written in this way, one should understand it as a formal expression of the formula given in Lemma 7.5. The main theorem of this section is the following:

Theorem 7.6. *Let $\Gamma_G = \text{Or}_{\mathcal{F}} G$ where G is a finite group and \mathcal{F} is a family of subgroups in G . Given a finite chain complex \mathbf{C} of projective $\mathbf{Z}\Gamma_G$ -modules, there exists an integer n such that n -fold join tensor $*_n \mathbf{C}$ of the complex \mathbf{C} is chain equivalent to a finite complex of free $\mathbf{Z}\Gamma_G$ -modules.*

We need to show that the finiteness obstruction $\tilde{\sigma}(*_n \mathbf{C})$ vanishes for some n . In the proof we will use a result by Oliver and Segev [26].

Proposition 7.7. *Let G be a finite group and let P and P' be any two finitely generated projective $\mathbf{Z}G$ -modules. Then, $P \otimes_{\mathbf{Z}} P'$ is stably free as a $\mathbf{Z}G$ -module.*

Proof. See [26, Proposition C.3]. \square

We also need the following splitting theorem for $K_0(\mathbf{Z}\Gamma)$.

Theorem 7.8. *Let Γ be a EI-category. Then, the map*

$$K_0(S) : K_0(\mathbf{Z}\Gamma) \rightarrow \bigoplus_{x \in \text{Iso}(\Gamma)} K_0(\mathbf{Z}[x]),$$

defined by $[P] \rightarrow [S_x(P)]$ on each $x \in \text{Iso}(\Gamma)$, is an isomorphism. The same holds when K_0 is replaced by \tilde{K}_0 .

Proof. See Lück [10, Proposition 11.29]. \square

As a consequence of this theorem, if Γ is finite then $\tilde{K}_0(\mathbf{Z}\Gamma)$ is finite: in this case Γ has finitely many isomorphism classes of objects $x \in \text{Ob}(\Gamma)$, and $\text{Aut}[x]$ is a finite group (apply Swan [31, Prop. 9.1]). In particular, if $\Gamma = \text{Or}_{\mathcal{F}} G$, then the group $\tilde{K}_0(\Gamma)$ is finite.

From now on we assume $\Gamma_G = \text{Or}_{\mathcal{F}} G$ for some finite group G , relative to some family \mathcal{F} . The splitting theorem above can also be used to give a filtration of $\tilde{K}_0(\Gamma_G)$. Recall that every projective $\mathbf{Z}\Gamma_G$ -module is of the form

$$P \cong \bigoplus_{H \in T} E_H S_H P$$

where T is a set of representatives of conjugacy classes of elements in \mathcal{F} . So, another way to express the above splitting theorem is to write

$$K_0(\mathbf{Z}\Gamma_G) \cong \bigoplus_{H \in T} K_0(\mathbf{Z}\Gamma_G)_H$$

where $K_0(\mathbf{Z}\Gamma_G)_H = \{[P] \mid E_H S_H P \cong P\}$. Note that this is only a splitting as abelian groups, but using this we can give a filtration for the ring structure of $K_0(\mathbf{Z}\Gamma_G)$. Let

$$\emptyset = T_0 \subseteq T_1 \subseteq \cdots \subseteq T_m = T$$

be a filtration of T such that if $H \in T_i$ and $K \in T_j$ and ${}^g H \leq K$ for some $g \in G$, then $i \leq j$. This gives a filtration

$$0 = K_0(\mathbf{Z}\Gamma_G)_0 \subseteq K_0(\mathbf{Z}\Gamma_G)_1 \subseteq \cdots \subseteq K_0(\mathbf{Z}\Gamma_G)_m = K_0(\mathbf{Z}\Gamma_G)$$

where

$$K_0(\mathbf{Z}\Gamma_G)_i = \{[P] \mid P = \bigoplus_{H \in T_i} E_H S_H P\}.$$

Lemma 7.9. *Let V be a $\mathbf{Z}[N_G(H)/H]$ -module and W be a $\mathbf{Z}[N_G(K)/K]$ -module. Then,*

$$E_H V \otimes_{\mathbf{Z}} E_K W \cong \bigoplus_{HgK \in H \backslash G / K} E_{H \cap {}^g K} (\text{Res}_{H \cap {}^g K} E_H V \otimes_{\mathbf{Z}} \text{Res}_{H \cap {}^g K} E_K W).$$

Proof. Applying the definition, we get

$$E_H V \otimes_{\mathbf{Z}} E_K W = (V \otimes W) \otimes_{\mathbf{Z}[\text{Aut}(G/H) \times \text{Aut}(G/K)]} \mathbf{Z} \text{Map}_G(?, G/H \times G/K)$$

where $\text{Map}_G(X, Y)$ denotes the set G -sets from X to Y (see [10, 11.30] for a similar computation). Since

$$G/H \times G/K = \coprod_{HgK \in H \backslash G / K} G/(H \cap {}^g K),$$

the module $E_H V \otimes_{\mathbf{Z}} E_K W$ decomposes as

$$\bigoplus_{HgK \in H \backslash G / K} E_{H \cap {}^g K} U_{H \cap {}^g K}$$

where $U_{H \cap {}^g K}$ are $N_G(H \cap {}^g K)/(H \cap {}^g K)$ -modules. Applying $S_{H \cap {}^g K}$, we find

$$\begin{aligned} U_{H \cap {}^g K} &= S_{H \cap {}^g K}(E_H V \otimes_{\mathbf{Z}} E_K W) = \text{Res}_{H \cap {}^g K}(E_H V \otimes_{\mathbf{Z}} E_K W) \\ &= \text{Res}_{H \cap {}^g K} E_H V \otimes_{\mathbf{Z}} \text{Res}_{H \cap {}^g K} E_K W. \end{aligned}$$

□

Lemma 7.10. $K_0(\mathbf{Z}\Gamma_G)_i$ is an ideal of $K_0(\mathbf{Z}\Gamma_G)$.

Proof. For $E_H S_H P$ and $E_K S_K Q$, we have

$$E_H S_H P \otimes_{\mathbf{Z}} E_K S_K Q = \bigoplus_L E_L V_L$$

where $L = H \cap {}^g K$ for some $g \in G$. So, if $H \in T_i$, $K \in T_j$, and $L \in T_k$, then $k \leq i, j$. \square

Now, Theorem 7.6 follows by induction from the following proposition.

Proposition 7.11. *Let \mathbf{C} be a finite chain complex of projective $\mathbf{Z}\Gamma_G$ -modules. If $\tilde{\sigma}(S_H \mathbf{C}) = 0$ for all $H \in T \setminus T_i$, then there is an n such that $\tilde{\sigma}(S_H(\ast_n \mathbf{C})) = 0$ for all $H \in T \setminus T_{i-1}$.*

Proof. An element in $\sigma(\mathbf{C})$ can be expressed as a sum $u + \sum_j v_j + w$ where

$$u = \sum_{H \in T_{i-1}} \sigma(E_H S_H \mathbf{C}), \quad \sum_j v_j = \sum_{H \in T_i \setminus T_{i-1}} \sigma(E_H S_H \mathbf{C}), \quad w = \sum_{H \in T \setminus T_i} \sigma(E_H S_H \mathbf{C}).$$

By Lemma 7.5, we have

$$1 - \sigma(\ast_n \mathbf{C}) = (1 - \sigma(\mathbf{C}))^n = (1 - (u + \sum_j v_j + w))^n \in K_0(\mathbf{Z}\Gamma_G)$$

So,

$$1 - \sigma(\ast_n \mathbf{C}) \equiv \left(1 - \left(\sum_j v_j + w\right)\right)^n \pmod{K_0(\mathbf{Z}\Gamma_G)_{i-1}}$$

By Lemma 7.9, it is easy to see that $v_j \cdot v_k \equiv 0 \pmod{K_0(\mathbf{Z}\Gamma_G)_{i-1}}$, for $j \neq k$. Note that

$$v_j \cdot v_j \equiv \text{stably free} \pmod{K_0(\mathbf{Z}\Gamma_G)_{i-1}}$$

by Proposition 7.7. To complete the proof, observe that modulo $K_0(\mathbf{Z}\Gamma_G)_{i-1}$,

$$\begin{aligned} 1 - \sigma(\ast_n \mathbf{C}) &\equiv \left(1 - \left(\sum_j v_j + w\right)\right)^n \\ &\equiv 1 + \sum_{k=1}^n \binom{n}{k} (-1)^k \left(\sum_j v_j + w\right)^k \\ &= 1 + \sum_{k=1}^n \binom{n}{k} (-1)^k \sum_j k v_j w^{k-1} + \text{stably free} \\ &= 1 + n \sum_{k=1}^n \sum_j \binom{n-1}{k-1} (-1)^k v_j w^{k-1} + \text{stably free}. \end{aligned}$$

This shows that $\sigma(\ast_n \mathbf{C})$ is stably free for some n , since $\tilde{K}_0(\mathbf{Z}\Gamma_G)$ is a finite group. \square

8. REALIZATION OF FREE CHAIN COMPLEXES

Let X be G -CW complex, and let \mathcal{F} be a family of subgroups of G . Throughout this section, R denotes a commutative ring and Γ_G denotes the orbit category $\text{Or}_{\mathcal{F}} G$.

Definition 8.1. We say that a G -CW complex X has *isotropy in \mathcal{F}* , provided that $X^H \neq \emptyset$ implies $H \in \mathcal{F}$, for all $H \leq G$.

The main result of this section is Theorem 8.10, which shows that under certain conditions a finite free chain complex over the orbit category can be realized by a finite G -CW complex with isotropy in \mathcal{F} . This is a generalization of Swan [32, Theorem A], which is based on a construction of Milnor [32, 3.1].

Associated to a G -CW complex X with isotropy in \mathcal{F} , there is a chain complex of $R\Gamma_G$ -modules defined by

$$\mathbf{C}(X^?; R): \quad \cdots \xrightarrow{\partial_{n+1}} R[X_n^?] \xrightarrow{\partial_n} R[X_{n-1}^?] \rightarrow \cdots \xrightarrow{\partial_1} R[X_0^?] \rightarrow 0$$

where X_i denotes the set of i -dimensional cells in X and $R[X_i^?]$ is the coefficient system with $R[X_i^?](H) = R[X_i^H]$. We denote the homology of this complex by $H_*(X^?; R)$, and in particular

$$H_*(X^?; R)(H) = H_*(X^H; R).$$

Given a chain complex \mathbf{C} of $R\Gamma_G$ -modules, there is a *dimension function* $\text{Dim } \mathbf{C}: \mathcal{F} \rightarrow \mathbf{Z}$, constant on conjugacy classes of subgroups, defined by

$$(\text{Dim } \mathbf{C})(H) = \dim \mathbf{C}(H),$$

for all $H \in \mathcal{F}$, where the dimension of a chain complex of R -modules is defined in the usual way as the largest integer d such $C_d \neq 0$.

It will be convenient to write $(H) \leq (K)$ whenever $H^g \leq K$ for some $g \in G$. Here (H) denotes the set of subgroups conjugate to H in G .

Definition 8.2. We call a function $\underline{d}: \mathcal{F} \rightarrow \mathbf{Z}$ *monotone* if it satisfies the property that $\underline{d}(K) \leq \underline{d}(H)$ whenever $(H) \leq (K)$. We say that a monotone function \underline{d} is *strictly monotone* if $\underline{d}(K) < \underline{d}(H)$, whenever $(H) \leq (K)$ and $(H) \neq (K)$. \square

Note that \underline{d} monotone implies that \underline{d} is constant on conjugacy classes (such functions are usually called *super class functions*). We remark that the dimension function of a projective chain complex is always monotone: if $(E_H P)(K) \neq 0$, then $(E_H P)(L) \neq 0$ for every $L \leq K$.

A chain complex \mathbf{C} of $R\Gamma_G$ -modules is *connected* if \mathbf{C} is positive and $H_0(\mathbf{C}) = \underline{R}$.

Definition 8.3. Let $\underline{n}: \mathcal{F} \rightarrow \mathbf{Z}$ be a monotone, non-negative function. A complex \mathbf{C} of $R\Gamma_G$ -modules is called an *\underline{n} -Moore complex* if it is connected, and for all $H \in \mathcal{F}$, the reduced homology $\tilde{H}_i(\mathbf{C}(H)) = 0$, for $i \neq \underline{n}(H)$. \square

A special case of an \underline{n} -Moore complex is a homology \underline{n} -sphere.

Definition 8.4. We say that a complex \mathbf{C} of $R\Gamma_G$ -modules is an *R -homology \underline{n} -sphere* if it is an \underline{n} -Moore complex, and for all $H \in \mathcal{F}$, we have $\tilde{H}_i(\mathbf{C}(H)) \cong R$, for $i = \underline{n}(H)$. A

homology \underline{n} -sphere is called *oriented* if the $N_G(H)/H$ -action is trivial on the homology of $\mathbf{C}(H)$ for all $H \in \mathcal{F}$.

The chain complex associated to the unit sphere $X = S(V)$ of a real or complex representation V of G is an example of a \mathbf{Z} -homology \underline{n} -sphere, where $\underline{n}(H) = \dim X^H$. A G -CW complex X with this property is a *homotopy representation* in the sense of tom Dieck (see [10, Chap. II, Def. 10.1]), provided that its dimension function is strictly monotone. We will not use this terminology further.

We now introduce a technique to remove free modules above the homological dimension from a chain complex, without changing its chain homotopy type. For this delicate process we first need some algebraic lemmas.

Definition 8.5. Let Γ be an EI-category. A free $R\Gamma$ -module F is called *isotypic* of type $x \in \text{Ob}(\Gamma)$ if it is isomorphic to a direct sum of copies of the free module $E_x R[x]$.

For extensions involving isotypic modules we have a splitting property.

Lemma 8.6. *Let*

$$\mathcal{E} : 0 \rightarrow F \rightarrow F' \rightarrow M \rightarrow 0$$

be a short exact sequence of $R\Gamma$ -modules over an EI-category Γ , such that both F and F' are isotypic free modules of the same type $x \in \text{Ob}(\Gamma)$. If $M(x)$ is R -torsion free, then \mathcal{E} splits and M is stably free.

Proof. It is enough to prove the result in case $F = E_x R[x]$, where $x \in \text{Ob} \Gamma$. The general case follows from this by an easy induction. Consider the extension

$$\mathcal{E} : 0 \rightarrow E_x R[x] \xrightarrow{j} F \rightarrow M \rightarrow 0 .$$

By the adjointness property

$$\text{Hom}_{R\Gamma}(E_x R[x], N) \cong \text{Hom}_{R[x]}(R[x], N(x))$$

for any $R\Gamma$ -module N . We apply this to the given injection $j: E_x R[x] \rightarrow F' = (E_x R[x])^m$. Since

$$\mathcal{E}(x) : 0 \rightarrow R[x] \xrightarrow{j} R[x]^m \rightarrow M(x) \rightarrow 0$$

has R -torsion free cokernel $M(x)$, this sequence splits over $R[x]$. By the naturality of the adjointness property, we get a splitting of j over $R\Gamma$. \square

Recall that $\text{hdim } \mathbf{C}(H)$ denotes the homological dimension of the chain complex $\mathbf{C}(H)$.

Proposition 8.7. *Let \mathbf{C} be a finite free chain complex of $R\Gamma_G$ -modules, and let $H \in \mathcal{F}$ have the property that $\text{hdim } \mathbf{C}(H) < d := \dim \mathbf{C}(H)$. Suppose that $\dim \mathbf{C}(K) \leq (d - 2)$ for all $(H) \leq (K)$, $(H) \neq (K)$. Then $\mathbf{C} \simeq \mathbf{D}$, where \mathbf{D} is a finite free complex with $\dim \mathbf{D}(H) = d - 1$, and $\dim \mathbf{D}(K) = \dim \mathbf{C}(K)$ for all $(K) \neq (H)$.*

Proof. Consider the subcomplex \mathbf{C}' of \mathbf{C} formed by free summands of \mathbf{C} isomorphic to $\mathbf{Z}[G/K^?]$, with $(G/K)^H \neq 0$ or equivalently $(H) \leq (K)$. The boundary maps of \mathbf{C}' are the restrictions of the usual boundary maps to these submodules. Since $\dim \mathbf{C}(K) \leq (d - 2)$

for all $(H) \leq (K)$ such that $(H) \neq (K)$, the free modules C'_d and C'_{d-1} are isotypic of type G/H . We have

$$\mathbf{C}' : 0 \rightarrow C'_d \rightarrow C'_{d-1} \rightarrow \cdots \rightarrow C'_1 \rightarrow C'_0 \rightarrow 0$$

where $d = \dim \mathbf{C}(H)$. Note that $\mathbf{C}(H) = \mathbf{C}'(H)$, so the map $\partial_d : C'_d \rightarrow C'_{d-1}$ is injective by the condition that $\text{hdim } \mathbf{C}(H) < \dim \mathbf{C}(H)$. Now we can apply Lemma 8.6 to the extension

$$0 \rightarrow C'_d \xrightarrow{\partial_d} C'_{d-1} \rightarrow \text{coker } \partial_d \rightarrow 0$$

and conclude that $\text{coker}(\partial_d)$ is a stably free $R\Gamma_G$ -module. By adding elementary chain complexes to \mathbf{C} of the form $\mathbf{Z}[G/H^?] \xrightarrow{\text{id}} \mathbf{Z}[G/H^?]$ in the adjacent dimensions $(d-1)$ and $(d-2)$, we can assume that $\text{coker}(\partial_d)$ is free. Consider the diagram

$$\begin{array}{cccccccccccc} \cdots & \longrightarrow & 0 & \longrightarrow & C'_d & \xrightarrow{\text{id}} & C'_d & \longrightarrow & 0 & \longrightarrow & \cdots & \longrightarrow & 0 & \longrightarrow & 0 \\ & & \downarrow & & \downarrow \text{id} & & \downarrow \partial_d & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \mathbf{C}' : & \cdots & \longrightarrow & 0 & \longrightarrow & C'_d & \longrightarrow & C'_{d-1} & \longrightarrow & C'_{d-2} & \longrightarrow & \cdots & \longrightarrow & C'_0 & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \parallel & & \parallel & & \parallel & & \parallel \\ \mathbf{D}' : & \cdots & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \text{coker } \partial_d & \longrightarrow & C'_{d-2} & \longrightarrow & \cdots & \longrightarrow & C'_0 & \longrightarrow & 0 \end{array}$$

The chain complex \mathbf{D}' is a chain complex of free modules and it is chain homotopy equivalent to \mathbf{C}' . Now define \mathbf{D} as the push-out in the the following diagram:

$$\begin{array}{ccccc} \ker \cong \ker & & & & \\ \downarrow & & \downarrow & & \\ \mathbf{C}' & \longrightarrow & \mathbf{C} & \longrightarrow & \mathbf{C}/\mathbf{C}' \\ \downarrow & & \downarrow & & \parallel \\ \mathbf{D}' & \longrightarrow & \mathbf{D} & \longrightarrow & \mathbf{C}/\mathbf{C}' \end{array}$$

Since, \mathbf{C}' and \mathbf{D}' are chain homotopy equivalent, then \mathbf{C} and \mathbf{D} are chain homotopy equivalent. Also, note that $\dim \mathbf{D}(H) = \dim \mathbf{D}'(H) = (d-1)$, and $\dim \mathbf{D}(K) = \dim \mathbf{C}(K)$ for all $(K) \neq (H)$. \square

This immediately gives the following.

Corollary 8.8. *Let \mathbf{C} be a finite free chain complex of $R\Gamma_G$ -modules. Suppose that \mathbf{C} is a homology \underline{n} -sphere, with \underline{n} strictly monotone. Then \mathbf{C} is chain homotopy equivalent to a complex \mathbf{D} with $\text{Dim } \mathbf{D} = \underline{n}$.*

Proof. Since \mathbf{C} is a homology \underline{n} -sphere, $\underline{n}(K) = \text{hdim } \mathbf{C}(K)$, for all $K \in \mathcal{F}$. We apply the previous result to a subgroup H , which is maximal with respect to the property that $\text{hdim } \mathbf{C}(H) < d := \dim \mathbf{C}(H)$. Then $\underline{n}(K) = \dim \mathbf{C}(K)$ for all $K \in \mathcal{F}$ larger than H . Since \underline{n} is strictly monotone, $\dim \mathbf{C}(K) \leq (d-2)$ for all $(H) \leq (K)$, $(H) \neq (K)$. This process can be repeated until $\text{Dim } \mathbf{D} = \underline{n}$. \square

When the dimension function of \mathbf{C} is not strictly monotone, we get a weaker result. Following Section 2, we define $l(H, K)$ as the maximum length of a chain of conjugacy classes of subgroups

$$(H) = (H_0) \not\leq (H_1) \not\leq \dots \not\leq (H_l) = (K)$$

where all $H_i \in \mathcal{F}$, $0 \leq i \leq l$.

Corollary 8.9. *Let \mathbf{C} be a finite free chain complex of $R\Gamma_G$ -modules, and let $\underline{n}: \mathcal{F} \rightarrow \mathbf{Z}$ be a monotone function such that $\text{hdim } \mathbf{C}(H) \leq \underline{n}(H)$ for all $H \in \mathcal{F}$. Assume that $l(H, K) \leq k$ whenever $\underline{n}(H) = \underline{n}(K)$. Then, \mathbf{C} is chain homotopy equivalent to a complex \mathbf{D} which satisfies $D_i(H) = 0$ for all $i > \underline{n}(H) + k$.*

Proof. Let

$$(H) = (H_0) \not\leq (H_1) \not\leq \dots \not\leq (H_l) = (K)$$

be a maximal length chain of subgroups in \mathcal{F} with $\underline{n}(H) = \underline{n}(K)$. Since \underline{n} is monotone, $\underline{n}(H_i) = \underline{n}(H)$ for $0 \leq i \leq l$. By repeated application of Proposition 8.7, working down from the maximal element K , we can obtain $\text{dim } \mathbf{C}(H_{l-i}) = \underline{n}(H) + i$, for $0 \leq i \leq l$. Since $l = l(H, K) \leq k$, we have $\text{dim } \mathbf{C}(H) \leq \underline{n}(H) + k$ as required. \square

The main purpose of this section is to prove the following theorem:

Theorem 8.10 (Pamuk [27]). *Let \mathbf{C} be a finite free chain complex of $\mathbf{Z}\Gamma_G$ -modules. Suppose \mathbf{C} is an \underline{n} -Moore complex such that $\underline{n}(H) \geq 3$ for all $H \in \mathcal{F}$. Suppose further that $C_i(H) = 0$ for all $i > \underline{n}(H) + 1$, and all $H \in \mathcal{F}$. Then there is a finite G -CW complex X , such that $\mathbf{C}(X^\?; \mathbf{Z})$ is chain homotopy equivalent to \mathbf{C} , as chain complexes of $\mathbf{Z}\Gamma_G$ -modules.*

Note that the resulting complex X will have isotropy in \mathcal{F} . We first prove a lemma (compare [20, Thm. 13.19]).

Lemma 8.11. *Let X be a finite G -CW complex. Suppose that we are given a free $\mathbf{Z}\Gamma_G$ -module F , and a $\mathbf{Z}\Gamma_G$ -module homomorphism $\varphi: F \rightarrow H_n(X^\?; \mathbf{Z})$, for some $n \geq 2$. Assume further that X^H is $(n-1)$ -connected for every $H \in \mathcal{F}$ such that $\mathbf{Z}[G/H^\?]$ is a summand of F . Then, by attaching $(n+1)$ -cells to X , we can obtain a G -CW complex Y such that*

$$H_i(X^\?; \mathbf{Z}) \cong H_i(Y^\?; \mathbf{Z}) \text{ for } i \neq n, n+1,$$

and

$$0 \rightarrow H_{n+1}(X^\?; \mathbf{Z}) \rightarrow H_{n+1}(Y^\?; \mathbf{Z}) \rightarrow F \xrightarrow{\varphi} H_n(X^\?; \mathbf{Z}) \rightarrow H_n(Y^\?; \mathbf{Z}) \rightarrow 0$$

is exact.

Proof. Let Z be a wedge of n -spheres with a G action on them such that $\tilde{H}_n(Z^\?; \mathbf{Z}) \cong F$ as $\mathbf{Z}\Gamma_G$ -modules. We want to construct a map $f: Z \rightarrow X$ realizing φ . But $H_n(X^H; \mathbf{Z}) \cong \pi_n(X^H)$, for every $H \in \mathcal{F}$ such that $\mathbf{Z}[G/H^\?]$ is a summand of F , since X^H is assumed to be $(n-1)$ -connected. Therefore, we can represent the images of an $\mathbf{Z}[N_G(H)/H]$ -basis under φ for the isotypic summand in F of type G/H by maps $f_i: S^n \rightarrow X^H$. We extend these maps equivariantly to maps $\tilde{f}_i: S^n \times G/H \rightarrow X$. By repeating this construction for each type G/H in F , we obtain an equivariant map $f: Z \rightarrow X$ realizing φ . Take Y to be the mapping cone of f . Then, it is easy to see that Y satisfies the desired conditions. \square

We also need the following lemma:

Lemma 8.12. *Let \mathbf{C} be a finite free chain complex of $\mathbf{Z}\Gamma_G$ -modules. Suppose that \mathbf{C} is connected, and $H_i(\mathbf{C}) = 0$, for $i = 1, 2$. Then, \mathbf{C} is chain homotopy equivalent to a complex of the form*

$$\cdots \rightarrow C_n \rightarrow C_{n-1} \rightarrow \cdots \rightarrow C_3 \rightarrow C_2(X) \rightarrow C_1(X) \rightarrow C_0(X) \rightarrow 0$$

where $C_2(X) \rightarrow C_1(X) \rightarrow C_0(X) \rightarrow 0$ is the initial part of the chain complex $\mathbf{C}(X^?; \mathbf{Z})$, for some G -CW complex X with isotropy in \mathcal{F} , and X^H simply-connected for all $H \in \mathcal{F}$.

Proof. There is a G -CW complex $E_{\mathcal{F}}G$ satisfying the properties

- (i) All isotropy subgroups of $E_{\mathcal{F}}G$ are in \mathcal{F} ,
- (ii) For every $H \in \mathcal{F}$, the fixed point set $(E_{\mathcal{F}}G)^H$ is contractible [22, Theorem 1.9].

The chain complex $\mathbf{D} := \mathbf{C}((E_{\mathcal{F}}G)^?; \mathbf{Z})$ of this space gives a free resolution of \mathbf{Z} as a $\mathbf{Z}\Gamma_G$ -module. Since $H_i(\mathbf{C}) = 0$, for $i = 1, 2$, the following sequences are both exact

$$(8.13) \quad \begin{array}{ccccccccccc} 0 & \longrightarrow & A & \longrightarrow & C_2 & \xrightarrow{\partial_2^C} & C_1 & \xrightarrow{\partial_1^C} & C_0 & \longrightarrow & \mathbf{Z} & \longrightarrow & 0 \\ & & & & & & & & & & \parallel & & \\ 0 & \longrightarrow & B & \longrightarrow & D_2 & \xrightarrow{\partial_2^D} & D_1 & \xrightarrow{\partial_1^D} & D_0 & \longrightarrow & \mathbf{Z} & \longrightarrow & 0 \end{array}$$

where $A = \ker \partial_2^C$ and $B = \ker \partial_2^D$.

By an *elementary operation* on a sequence $A \rightarrow C_2 \rightarrow C_1 \rightarrow C_0$ we mean adding or removing trivial free summands $F \xrightarrow{id} F$ in adjacent dimensions. It is clear that elementary operations don't change the chain homotopy type of the upper and lower sequences in diagram (8.13).

Then, by Schanuel's Lemma [32, 1.1], there exist free modules F and F' such that $A \oplus F \cong B \oplus F'$. In fact, the argument in Schanuel's lemma can be extended to say that the induced isomorphism $\gamma: A \oplus F \cong B \oplus F'$ comes from a chain isomorphism after a sequence of elementary operations (compare [20, p. 279]).

In other words, there exists a chain isomorphism

$$(8.14) \quad \begin{array}{ccccccccccc} 0 & \longrightarrow & A \oplus F & \longrightarrow & C_2 \oplus F_2 & \longrightarrow & C_1 \oplus F_1 & \longrightarrow & C_0 \oplus F_0 & \longrightarrow & \mathbf{Z} & \longrightarrow & 0 \\ & & \gamma \downarrow \cong & & \cong \downarrow f_2 & & \cong \downarrow f_1 & & \cong \downarrow f_0 & & \parallel & & \\ 0 & \longrightarrow & B \oplus F' & \longrightarrow & D_2 \oplus F'_2 & \longrightarrow & D_1 \oplus F'_1 & \longrightarrow & D_0 \oplus F'_0 & \longrightarrow & \mathbf{Z} & \longrightarrow & 0 \end{array}$$

for some suitable choices of free modules, where the upper and lower sequences in diagram (8.14) are obtained from those in diagram (8.13) by elementary operations (see Proposition 3.3.3 in [27]).

In the first step, we stabilize $(A \rightarrow C_2) \mapsto (A \oplus F \rightarrow C_2 \oplus F)$, by adding the identity on F , and similarly $(B \rightarrow D_2) \mapsto (B \oplus F' \rightarrow D_2 \oplus F')$. We therefore have a chain map

$$\begin{array}{ccccccccccc} 0 & \longrightarrow & A \oplus F & \longrightarrow & C_2 \oplus F & \longrightarrow & C_1 & \longrightarrow & C_0 & \longrightarrow & \underline{\mathbf{Z}} & \longrightarrow & 0 \\ & & \gamma \downarrow \cong & & \downarrow & & \downarrow & & \downarrow & & \parallel & & \\ 0 & \longrightarrow & B \oplus F' & \longrightarrow & D_2 \oplus F' & \longrightarrow & D_1 & \longrightarrow & D_0 & \longrightarrow & \underline{\mathbf{Z}} & \longrightarrow & 0 \end{array}$$

which is a chain homotopy equivalence (by composition with the chain map in (8.14)). After an elementary operation on \mathbf{C} , we can use the isomorphism $\gamma: A \oplus F \cong B \oplus F'$ to splice the bottom sequence to \mathbf{C} , and obtain a chain homotopy equivalence

$$\begin{array}{ccccccccccc} \cdots & \longrightarrow & C_4 & \longrightarrow & C_3 \oplus F & \longrightarrow & C_2 \oplus F & \longrightarrow & C_1 & \longrightarrow & C_0 & \longrightarrow & \underline{\mathbf{Z}} & \longrightarrow & 0 \\ & & \parallel & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \parallel & & \\ \cdots & \longrightarrow & C_4 & \longrightarrow & C_3 \oplus F & \longrightarrow & D_2 \oplus F' & \longrightarrow & D_1 & \longrightarrow & D_0 & \longrightarrow & \underline{\mathbf{Z}} & \longrightarrow & 0 \end{array}$$

The top sequence is chain homotopy equivalent to \mathbf{C} , so to complete the proof we need to show that the sequence $D_2 \oplus F' \rightarrow D_1 \rightarrow D_0 \rightarrow 0$ can be realized as the first three terms of a chain complex of a G -CW complex X , with isotropy in \mathcal{F} , such that X^H is simply connected for all $H \in \mathcal{F}$: since $E_{\mathcal{F}}G$ is contractible, using Lemma 8.11, we can attach free 2-cells to its two skeleton $E_{\mathcal{F}}G^{(2)}$. The resulting complex X will have the desired properties. \square

Now, we are ready to prove Theorem 8.10.

Proof of Theorem 8.10. We can assume that the complex \mathbf{C} is of the form given in Lemma 8.12. We obtain a map $\varphi: C_3 \rightarrow C_2(X^{(2)})$ which induces an isomorphism $Z_2(\mathbf{C}) \rightarrow Z_2(X^{(2)})$ between 2-cycles of these chain complexes. This is the starting point for an inductive argument based on applying Lemma 8.11 at each step.

Fix $n \geq 2$, and assume by induction that there is an n -dimensional G -CW complex $X^{(n)}$, and a chain map

$$\begin{array}{ccccccccccc} \cdots & \longrightarrow & C_{n+2} & \longrightarrow & C_{n+1} & \longrightarrow & C_n & \longrightarrow & \cdots & \longrightarrow & C_1 & \longrightarrow & C_0 & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \\ \cdots & \longrightarrow & 0 & \longrightarrow & Z_n(X^{(n)}) & \longrightarrow & C_n(X^{(n)}) & \longrightarrow & \cdots & \longrightarrow & C_1(X^{(n)}) & \longrightarrow & C_0(X^{(n)}) & \longrightarrow & 0 \end{array}$$

which induces an homology isomorphism for dimensions less than or equal to $(n-1)$, and at dimension n the induced map $Z_n(\mathbf{C}) \rightarrow Z_n(X^{(n)})$ is an isomorphism.

Note that $\dim \mathbf{C}(H) \leq \underline{n}(H) + 1$ by assumption. If $\mathbf{Z}[G/H^?]$ is a summand of C_{n+1} , then $(n+1) \leq \dim \mathbf{C}(H) \leq \underline{n}(H) + 1$ implies $\underline{n}(H) \geq n$, and hence the H -fixed set of $X^{(n)}$ is $(n-1)$ -connected. We can now apply Lemma 8.11 to the map $\varphi: C_{n+1} \rightarrow H_n(X^{(n)}; \mathbf{Z})$ defined by the composition

$$\phi: C_{n+1} \rightarrow Z_n(\mathbf{C}) \cong Z_n(X^{(n)}) \rightarrow H_n(X^{(n)}; \mathbf{Z}).$$

Let us call the resulting complex $X^{(n+1)}$. Note that there is a chain map $\mathbf{C} \rightarrow \mathbf{C}(X^{(n+1)})$ which induces an isomorphism on homology for dimensions $\leq n$, and at dimension $n+1$

we have an isomorphism $Z_{n+1}(\mathbf{C}) \rightarrow Z_{n+1}(X^{(n+1)})$. Since \mathbf{C} is finite dimensional, after finitely many steps, we will obtain a finite dimensional G -CW complex X and a chain map $f: \mathbf{C} \rightarrow \mathbf{C}(X)$ which induces isomorphism on homology for all dimensions. Since both \mathbf{C} and $\mathbf{C}(X)$ are free $\mathbf{Z}\Gamma_G$ -chain complexes, f is a chain homotopy equivalence as desired. \square

9. THE PROOF OF THEOREM A

Let $G = S_5$, the symmetric group of order 120 permuting $\{1, 2, 3, 4, 5\}$, and let $S_4 \leq G$ denote the permutations fixing $\{5\}$. We work relative to the family \mathcal{F} of rank 1 subgroups of 2-power order. Let $\Gamma_G = \text{Or}_{\mathcal{F}}G$. Our family \mathcal{F} consists of the subgroups of G which are conjugate to one the subgroups in the set

$$\{1, C_2^A, C_2^B, C_4\}$$

where $C_2^A = \langle (12)(34) \rangle$, $C_2^B = \langle (12) \rangle$, and $C_4 = \langle (1234) \rangle$. In addition we will consider the Sylow subgroups $C_3 = \langle (123) \rangle$ and $C_5 = \langle (12345) \rangle$. It is convenient to note that for $H = S_4 \leq G$, we have

$$N_H(C_4) = D_8 = N_H(C_2^A) = N_G(C_4),$$

while $N_H(C_2^B) = E = \langle (12), (34) \rangle$, and $N_H(C_3) = S_{\{123\}}$. On the other hand, $N_G(C_2^B) = \langle (12), S_{\{345\}} \rangle$ and $N_G(C_3) = S_{\{123\}} \times \langle (45) \rangle$.

Our strategy will be to construct finite projective complexes $\mathbf{C}^{(p)}$ with isotropy in \mathcal{F} over $\mathbf{Z}_{(p)}\Gamma_G$, for each prime p dividing the order of $|G|$, which are R -homology \underline{n} -spheres with respect to the the same homology dimension function \underline{n} . The gluing theory of Section 6, Theorem 6.7, will be used to construct a finite projective \mathbf{Z} -homology \underline{n} -sphere over $\mathbf{Z}\Gamma_G$ from this data. Then the join construction from Section 7 will allow us to find a finite free complex, to which the realization theorem of Section 8 will apply.

We introduce the notation \underline{R}_0 for the $R\Gamma_G$ -module defined by $\underline{R}_0(K) = 0$, for $K \neq 1$, and $\underline{R}_0(1) = R$ with trivial G -action. In other words, $\underline{R}_0 = I_1(R)$ as defined in Section 2.

9A. The case $p = 2$. Let $H = S_4 \leq G$, $R = \mathbf{Z}_{(2)}$ and consider the standard H -action on the 2-sphere given by the rotational symmetries of the octahedron. Let X denote the H -CW complex associated to the first barycentric subdivision of the octahedron. Then X has isotropy in the family consisting of the cyclic subgroups of H of orders ≤ 4 .

Let $\Gamma_H = \text{Or}_{\mathcal{F}}H$ denote the orbit category for H with respect to the family $\mathcal{F}_H = \mathcal{F} \cap H$. Consider the chain complex $\mathbf{C}(X^?; R)$ as a chain complex of $R\Gamma_H$ -modules, by restricting this functor to the full subcategory Γ_H of the orbit category $\text{Or}(H)$. This gives an exact sequence of the form

$$0 \rightarrow \underline{R}_0 \rightarrow 2R[H/1^?] \rightarrow 3R[H/1^?] \rightarrow R[H/C_4^?] \oplus R[H/C_2^B^?] \oplus R[H/C_3^?] \rightarrow \mathbf{H}_0 \rightarrow 0,$$

where all the modules in the extension (excluding the ends) except $R[H/C_3^?]$ are free $R\Gamma_H$ -modules, and $\mathbf{H}_0 = H_0(\mathbf{C}(X^?; R))$.

Since $R[H/C_3]$ is a projective RH -module (it is induced up from R , which is projective over $R[C_3]$), we see that $R[H/C_3^?] = I_1R[H/C_3]$ as $R\Gamma_H$ -modules. Therefore $\mathbf{C}(X^?; R)$ is a finite projective chain complex over $R\Gamma_H$.

It is useful to represent an $R\Gamma_H$ -module M by a labelled tree diagram:

$$M = \begin{array}{c} M(C_4) \\ | \\ M(C_2^A) \quad M(C_2^B) \\ | \quad / \\ M(1) \end{array}$$

with one vertex for each isomorphism class of objects, and edges given by the partial ordering of the subgroups in \mathcal{F} up to conjugacy. The labels are given by the $R[N_H(K)/K]$ -modules $M(K)$, for $K \in \mathcal{F}$.

For the homology module $H_0(X^?; R)$ over the orbit category Γ_H we have the diagram

$$\mathbf{H}_0 = \begin{array}{c} R[D_8/C_4] \\ | \\ R[D_8/C_4] \quad R[E/C_2^B] \\ | \quad / \\ R \end{array}$$

The $(k+1)$ -fold join of $\mathbf{C}(X^?; R)$ with itself (see Section 7) is a finite projective complex of the form

$$\mathbf{C} : 0 \rightarrow \underline{R}_0 \rightarrow C_n \rightarrow \cdots \rightarrow C_k \rightarrow \cdots \rightarrow C_0 \rightarrow \underline{R} \rightarrow 0$$

over $R\Gamma_H$ with $(n+1) = 3(k+1)$. In case $X = S(V)$, where $V \cong R^3$ is a real orthogonal H -representation, then the join construction on spheres just produces the unit sphere $S(V \oplus \cdots \oplus V)$ in the direct sum of $(k+1)$ copies of V . This sphere has real dimension $n = 3(k+1) - 1$. The purpose of the join construction is to produce a complex with dimension gaps between the non-zero homology groups, as required by Theorem 6.7 for glueing the different primes together.

We have $H_0(\mathbf{C}) = \underline{R}$ and $H_n = \underline{R}_0$. If $(k+1)$ is even, then $H_k(\mathbf{C}(Q)) = R$, with trivial $N_H(Q)/Q$ -action, and $\tilde{H}_i(\mathbf{C}(Q)) = 0$, for $i \neq k$, for each non-trivial $Q \in \mathcal{F}$. By Proposition 6.4, we obtain a chain complex $\mathbf{C}^{(2)}$ of projective $R\Gamma_G$ -modules, having homology isomorphic to R , with trivial $N_G(Q)/Q$ -action. By construction, the homology dimension function \underline{n} for $\mathbf{C}^{(2)}$ is the same as for \mathbf{C} . Notice that \underline{n} is monotone, but not strictly monotone.

9B. The case $p = 3$. Let $R = \mathbf{Z}_{(3)}$ and $K = C_2^B$. The 3-period of $G = S_5$ is four [7, Chap. XII, Ex. 11], so by Swan [32] there exists a periodic projective resolution \mathbf{P} with

$$0 \rightarrow R \rightarrow P_n \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow R \rightarrow 0$$

over the group ring RG , for any n such that $4 \mid (n+1)$. We will assume that $12 \mid (n+1)$, and let k be defined by the equation $(n+1) = 3(k+1)$. Similarly, since $N_G(K)/K \cong S_3$

also has 3-period 4, we have a chain complex \mathbf{D} yielding a periodic projective resolution

$$0 \rightarrow R \rightarrow D_k \rightarrow \cdots \rightarrow D_0 \rightarrow R \rightarrow 0$$

over RS_3 . In the rest of this section we let $W_K = N_G(K)/K$ to simplify the notation.

We want a finite projective chain complex \mathbf{C} over RG which fits into an extension of chain complexes

$$0 \rightarrow E_1\mathbf{P} \rightarrow \mathbf{C} \rightarrow I_K\mathbf{D} \rightarrow 0$$

where the induced exact sequence on the 0-th homology

$$0 \rightarrow \underline{R}_0 \rightarrow H_0(\mathbf{C}) \rightarrow I_K R \rightarrow 0$$

is the non-trivial extension of tree diagrams (with vertices at $\{1, K\}$)

$$0 \rightarrow \begin{array}{c} 0 \\ | \\ R \end{array} \rightarrow \begin{array}{c} R \\ | \\ R \end{array} \xrightarrow{id} \begin{array}{c} R \\ | \\ 0 \end{array} \rightarrow 0.$$

For a projective $R[W_K]$ -module D , the module $I_K D$ has a finite projective resolution of the form

$$0 \longrightarrow E_1 \operatorname{Res}_1 E_K D \longrightarrow E_K D \longrightarrow I_K D \longrightarrow 0.$$

By definition of the functors E_x and I_x (see Section 2), the canonical map

$$E_x M \rightarrow I_x M \rightarrow 0$$

is always surjective for any $R[x]$ -module M . We have $E_K R[W_K] = R[G/K?]$ and hence $E_K D$ is projective. Also $\operatorname{Res}_1 E_K D$ is projective, because it is a summand of $R[G/K]$ which is projective as a $\mathbf{Z}_{(3)}G$ -module. This shows that, once constructed, \mathbf{C} will be homotopy equivalent to a finite projective chain complex by Lemma 6.2.

Associated to every RG -chain map $f: \operatorname{Res}_1 E_K \mathbf{D} \rightarrow \mathbf{P}$, there is a chain complex \mathbf{C} which fits into the push-out diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & E_1 \operatorname{Res}_1 E_K \mathbf{D} & \longrightarrow & E_K \mathbf{D} & \longrightarrow & I_K \mathbf{D} \longrightarrow 0 \\ & & \downarrow E_1 f & & \downarrow & & \parallel \\ 0 & \longrightarrow & E_1 \mathbf{P} & \longrightarrow & \mathbf{C} & \longrightarrow & I_K \mathbf{D} \longrightarrow 0. \end{array}$$

We want to choose f so that \mathbf{C} satisfies the condition on homology. Note that

$$H_0(\operatorname{Res}_1 E_K \mathbf{D}) = \operatorname{Res}_1 E_K R = R[G/N_G(K)].$$

Since the modules $\operatorname{Res}_1 E_K D_i = (E_K D_i)(1)$ are projective for all i and \mathbf{P} is exact, there exists a chain map $f: \operatorname{Res}_1 E_K \mathbf{D} \rightarrow \mathbf{P}$

$$\begin{array}{ccccccc} \cdots & \longrightarrow & (E_K D_1)(1) & \longrightarrow & (E_K D_0)(1) & \longrightarrow & R[G/N_G(K)] \longrightarrow 0 \\ & & \downarrow f_1 & & \downarrow f_0 & & \downarrow \varepsilon \\ \cdots & \longrightarrow & P_1 & \longrightarrow & P_0 & \longrightarrow & R \longrightarrow 0 \end{array}$$

lifting the augmentation map $R[G/N_G(K)] \xrightarrow{\varepsilon} R$. To see that the resulting push-out complex \mathbf{C} has the desired properties, consider the homology at zero for the diagram of

chain complexes given above. Since I_K is an exact functor, $H_1(I_K \mathbf{D}) = I_K H_1(\mathbf{D}) = 0$, and we get

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_0(E_1 \operatorname{Res}_1 E_K \mathbf{D}) & \longrightarrow & H_0(E_K \mathbf{D}) & \longrightarrow & H_0(I_K \mathbf{D}) \longrightarrow 0 \\ & & \downarrow H_0(E_1 f) & & \downarrow & & \parallel \\ 0 & \longrightarrow & H_0(E_1 \mathbf{P}) & \longrightarrow & H_0(\mathbf{C}) & \longrightarrow & H_0(I_K \mathbf{D}) \longrightarrow 0 \end{array}$$

where $H_0(E_1 \mathbf{P}) = E_1 R$. Note that

$$H_0(E_1 \operatorname{Res}_1 E_K \mathbf{D}) = E_1 \operatorname{Res}_1 H_0(E_K \mathbf{D}) = E_1 \operatorname{Res}_1 E_K R = E_1 R[G/N_G(K)].$$

This gives a diagram of the form

$$\begin{array}{ccccccc} & & E_1 \ker \varepsilon & \xlongequal{\quad} & E_1 \ker \varepsilon & & \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & E_1 R[G/N_G(K)] & \longrightarrow & E_K R & \longrightarrow & I_K R \longrightarrow 0 \\ & & \downarrow E_1(\varepsilon) & & \downarrow & & \parallel \\ 0 & \longrightarrow & E_1 R & \longrightarrow & H_0(\mathbf{C}) & \longrightarrow & I_K R \longrightarrow 0 \end{array}$$

where the middle vertical sequence of $R\Gamma_G$ -modules is given by

$$0 \rightarrow \begin{array}{c} 0 \\ \downarrow \\ \ker \varepsilon \end{array} \rightarrow \begin{array}{c} R \\ \downarrow \\ R[G/N_G(K)] \end{array} \rightarrow \begin{array}{c} R \\ \downarrow \text{id} \\ R \end{array} \rightarrow 0$$

This shows that $H_0(\mathbf{C})$ has the desired form.

Now, to obtain the same homology dimension function as for the complex $\mathbf{C}^{(2)}$, more homology must be added to the complex \mathbf{C} . We need to extend \mathbf{H}_k and \mathbf{H}_0 via the non-split extensions

$$0 \rightarrow \mathbf{H}_0 \rightarrow \hat{\mathbf{H}}_0 \rightarrow N \rightarrow 0 \quad \text{and} \quad 0 \rightarrow H_k \rightarrow \hat{H}_k \rightarrow N \rightarrow 0$$

where

$$N = \begin{array}{ccc} & R & \\ & \downarrow \text{id} & \\ & R & \searrow 0 \\ & \downarrow & \\ & 0 & \end{array}$$

The module N has a finite projective resolution of the form

$$0 \rightarrow E_1 R[G/D_8] \rightarrow E_{C_4} R \rightarrow N \rightarrow 0.$$

Note that $\operatorname{Res}_1 E_{C_4} R = R[G/N_G(C_4)] = R[G/D_8]$, and for $Q = C_2^A$ we have

$$\operatorname{Res}_Q E_{C_4} R = R \otimes_{R[D_8/C_4]} R[(G/C_4)^Q] = R \otimes_{R[D_8/C_4]} R[N_G(Q)/N_{C_4}(Q)] = R$$

where the equality in the middle comes from Lemma 3.6. Since R is projective as an $R[D_8/C_4]$ -module, $E_{C_4}R$ is projective. It is easy to see that $E_1R[G/D_8]$ is also projective. So, by Proposition 6.8, we can replace \mathbf{C} with a finite projective chain complex $\mathbf{C}^{(3)}$ over $R\Gamma_G$ which has the desired homology.

9C. The case $p = 5$. For $p = 5$, the situation is easier than the case $p = 3$. Let $R = \mathbf{Z}_{(5)}$. The 5-period of S_5 equals 8, so by Swan [32] there exists a periodic projective resolution \mathbf{P} over the group ring RG , giving an exact sequence

$$0 \rightarrow R \rightarrow P_n \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow R \rightarrow 0$$

for any positive integer n such that $n + 1 = 3(k + 1)$ for some integer k , with $8 \mid (k + 1)$. We start with the $R\Gamma_G$ -complex $\mathbf{C} = E_1\mathbf{P}$ obtained by the extension functor from \mathbf{P} . Since \mathbf{C} has no homology at the non-trivial 2-subgroups in \mathcal{F} , we need to change the homology at \mathbf{H}_0 and at \mathbf{H}_k to match the homology we have for $p = 2$ and $p = 3$. Note that we need to extend H_k and \mathbf{H}_0 via the non-split extensions

$$0 \rightarrow \mathbf{H}_0 \rightarrow \hat{\mathbf{H}}_0 \rightarrow M \rightarrow 0 \quad \text{and} \quad 0 \rightarrow \mathbf{H}_k \rightarrow \hat{\mathbf{H}}_k \rightarrow M \rightarrow 0$$

where

$$M = \begin{array}{ccc} & R & \\ & \downarrow \text{id} & \\ & R & R \\ & \downarrow & \swarrow \\ & 0 & \end{array}$$

Let $K = C_2^B$. The module M is the direct sum of L (which has the same form as N) and $I_K R$. We claim that each of these modules have finite projective resolutions over $R\Gamma_G$. For $I_K R$ we have a resolution of the form

$$0 \rightarrow E_1R[G/(K \times S_3)] \rightarrow E_K R \rightarrow I_K R \rightarrow 0.$$

Note that

$$\text{Res}_1 E_K R = R[G/N_G(K)] = R[G/(K \times S_3)]$$

where S_3 denotes the subgroup of S_5 generated by symmetries of $\{3, 4, 5\}$. Since R is projective as an $R[N_G(K)/K]$ -module, $E_K R$ is projective. It is clear that $E_1R[G/(K \times S_3)]$ is also projective. So, the above resolution is a projective resolution of $I_K R$. We can also write a finite projective resolution for L (similar to the resolution given for N). So, by Proposition 6.8, we can replace \mathbf{C} with a finite projective chain complex $\mathbf{C}^{(5)}$ which has the desired homology.

The proof of Theorem A. We will first construct a projective chain complex \mathbf{C} over $\mathbf{Z}\Gamma_G$ with isotropy in \mathcal{F} , by applying Theorem 6.7 to glue the p -local complexes $\mathbf{C}^{(p)}$, for $p = 2, 3, 5$. Note that in the constructions of $\mathbf{C}^{(p)}$ above, we may choose any integer k such that k odd, $n + 1 = 3(k + 1)$, $12 \mid (n + 1)$ and $8 \mid (k + 1)$. To satisfy the first condition in Theorem 6.7, that the distance between non-zero homology groups of the $\mathbf{C}^{(p)}$ is larger than $l(\Gamma_G) = 2$, we will also need $k \geq 3$ and $n - k \geq 3$.

Remark 9.1. The minimum value for k satisfying the requirements used above is $k = 7$, which gives $n = \dim \mathbf{C} = 23$.

The $\mathbf{Z}\Gamma_G$ -module \mathbf{H} needed to satisfy the second condition in Theorem 6.7 is given by $\mathbf{H}_i(K) = \mathbf{Z}$, for $i = 0, \underline{n}(K)$ with $K \in \mathcal{F}$, and zero otherwise. By Lemma 6.2, \mathbf{C} is chain homotopy equivalent to a finite projective complex. To obtain a finite free complex, we can apply Theorem 7.6, which (possibly after some joins) produces a finite free $\mathbf{Z}\Gamma_G$ -chain complex \mathbf{C} with the \mathbf{Z} -homology of an \underline{n} -sphere, and $\underline{n}(K) \geq 3$ for all $K \in \mathcal{F}$.

Note that our homology dimension function \underline{n} is not strictly monotone, since $\underline{n}(C_2^A) = \underline{n}(C_4)$, but by Corollary 8.9 we can modify our complex to satisfy the conditions for geometric realization in Theorem 8.10, since $l(C_2^A, C_4) = 1$. Applying Theorem 8.10, we conclude that $G = S_5$ acts on a finite G -CW complex X with isotropy in \mathcal{F} . \square

Remark 9.2. For this particular example we needed to apply Theorem 7.6 with one join tensor operation, because $\tilde{K}_0(\mathbf{Z}\Gamma_G) = \mathbf{Z}/2$. This follows from Theorem 7.8, Lemma 7.5 and well-known calculations showing that $\tilde{K}_0(\mathbf{Z}[N_G(Q)/Q]) = 0$, for $1 \neq Q \in \mathcal{F}$, but $\tilde{K}_0(\mathbf{Z}[G]) = \mathbf{Z}/2$. Note that, by Dress induction, it is enough to consider the projective class groups of p -hypercyclic subgroups of G (see [9, §50], [28]). We therefore obtain a finite G -CW complex $X \simeq S^{47}$ with isotropy in \mathcal{F} .

REFERENCES

- [1] A. Adem and J. H. Smith, *Periodic complexes and group actions*, Ann. of Math. (2) **154** (2001), 407–435.
- [2] D. J. Benson, *Representations and cohomology. I*, second ed., Cambridge Studies in Advanced Mathematics, vol. 30, Cambridge University Press, Cambridge, 1998, Basic representation theory of finite groups and associative algebras.
- [3] ———, *Representations and cohomology. II*, second ed., Cambridge Studies in Advanced Mathematics, vol. 31, Cambridge University Press, Cambridge, 1998, Cohomology of groups and modules.
- [4] S. Bouc, *Résolutions de foncteurs de Mackey*, Group representations: cohomology, group actions and topology (Seattle, WA, 1996), Proc. Sympos. Pure Math., vol. 63, Amer. Math. Soc., Providence, RI, 1998, pp. 31–83.
- [5] N. Brady, I. J. Leary, and B. E. A. Nucinkis, *On algebraic and geometric dimensions for groups with torsion*, J. London Math. Soc. (2) **64** (2001), 489–500.
- [6] G. E. Bredon, *Equivariant cohomology theories*, Lecture Notes in Mathematics, No. 34, Springer-Verlag, Berlin, 1967.
- [7] H. Cartan and S. Eilenberg, *Homological algebra*, Princeton Landmarks in Mathematics, Princeton University Press, Princeton, NJ, 1999, With an appendix by David A. Buchsbaum, Reprint of the 1956 original.
- [8] O. Coşkun and E. Yalçın, *A Tate cohomology sequence for generalized Burnside rings*, J. Pure Appl. Algebra **213** (2009), 1306–1315.
- [9] C. W. Curtis and I. Reiner, *Methods of representation theory. Vol. II*, John Wiley & Sons Inc., New York, 1987, With applications to finite groups and orders, A Wiley-Interscience Publication.
- [10] T. t. Dieck, *Transformation groups*, de Gruyter Studies in Mathematics **8**, Walter de Gruyter & Co., Berlin, 1987.
- [11] A. Dold, *Zur Homotopietheorie der Kettenkomplexe*, Math. Ann. **140** (1960), 278–298.
- [12] R. M. Dotzel and G. C. Hamrick, *p -group actions on homology spheres*, Invent. Math. **62** (1981), 437–442.

- [13] A. W. M. Dress, *Induction and structure theorems for orthogonal representations of finite groups*, Ann. of Math. (2) **102** (1975), 291–325.
- [14] J. Grodal, *Higher limits via subgroup complexes*, Ann. of Math. (2) **155** (2002), 405–457.
- [15] J. Grodal and J. H. Smith, *Algebraic models for finite G -spaces*, Oberwolfach Reports, vol. 4, Eur. Math. Soc., Zürich, 2007, Abstracts from the Homotopy Theory workshop, September 16–22, 2007, organized by Paul Goerss, John Greenlees and Stefan Schwede, pp. 2690–2692.
- [16] I. Hambleton, L. R. Taylor, and E. B. Williams, *Dress Induction and the Burnside quotient Green ring*, Algebra & Number Theory **3** (2009), 511–541.
- [17] S. Jackowski, J. McClure, and B. Oliver, *Homotopy classification of self-maps of BG via G -actions. II*, Ann. of Math. (2) **135** (1992), 227–270.
- [18] M. A. Jackson, *$Qd(p)$ -free rank two finite groups act freely on a homotopy product of two spheres*, J. Pure Appl. Algebra **208** (2007), 821–831.
- [19] T.-Y. Lam, *Induction theorems for Grothendieck groups and Whitehead groups of finite groups*, Ann. Sci. École Norm. Sup. (4) **1** (1968), 91–148.
- [20] W. Lück, *Transformation groups and algebraic K -theory*, Lecture Notes in Mathematics, vol. 1408, Springer-Verlag, Berlin, 1989, Mathematica Gottingensis.
- [21] ———, *Chern characters for proper equivariant homology theories and applications to K - and L -theory*, J. Reine Angew. Math. **543** (2002), 193–234.
- [22] ———, *Survey on classifying spaces for families of subgroups*, Infinite groups: geometric, combinatorial and dynamical aspects, Progr. Math., vol. 248, Birkhäuser, Basel, 2005, pp. 269–322.
- [23] S. Mac Lane, *Homology*, Classics in Mathematics, Springer-Verlag, Berlin, 1995, Reprint of the 1975 edition.
- [24] ———, *Categories for the working mathematician*, second ed., Graduate Texts in Mathematics, vol. 5, Springer-Verlag, New York, 1998.
- [25] G. Mislin, *On group homomorphisms inducing mod- p cohomology isomorphisms*, Comment. Math. Helv. **65** (1990), 454–461.
- [26] B. Oliver and Y. Segev, *Fixed point free actions on \mathbf{Z} -acyclic 2-complexes*, Acta Math. **189** (2002), 203–285.
- [27] S. Pamuk, *Periodic resolutions and finite group actions*, Ph.D. thesis, McMaster University, 2008.
- [28] I. Reiner, *Projective class groups of symmetric and alternating groups*, Linear and Multilinear Algebra **3** (1975/76), 147–153, Collection of articles dedicated to Olga Taussky-Todd.
- [29] D. S. Rim, *Modules over finite groups*, Ann. of Math. (2) **69** (1959), 700–712.
- [30] E. H. Spanier, *Algebraic topology*, McGraw-Hill Book Co., New York, 1966.
- [31] R. G. Swan, *Induced representations and projective modules*, Ann. of Math. (2) **71** (1960), 552–578.
- [32] ———, *Periodic resolutions for finite groups*, Ann. of Math. (2) **72** (1960), 267–291.
- [33] P. Symonds, *Mackey functors and control of fusion*, Bull. London Math. Soc. **36** (2004), 623–632.
- [34] ———, *Smith theory for algebraic varieties*, Algebr. Geom. Topol. **4** (2004), 121–131 (electronic).
- [35] ———, *The Bredon cohomology of subgroup complexes*, J. Pure Appl. Algebra **199** (2005), 261–298.

DEPARTMENT OF MATHEMATICS, MCMASTER UNIVERSITY, CANADA
E-mail address: hambleton@mcmaster.ca

DEPARTMENT OF MATHEMATICS, MCMASTER UNIVERSITY, CANADA
E-mail address: pamuks@math.mcmaster.ca

DEPARTMENT OF MATHEMATICS, BILKENT UNIVERSITY, 06800 BILKENT, ANKARA, TURKEY
E-mail address: yalcine@fen.bilkent.edu.tr