

$$\left(-\frac{1}{3}, \pm \frac{4}{3}\sqrt{2}\right)$$

## Antiderivatives

Given  $f'(x)$ , what could  $f(x)$  be?

**Example:** A particle is accelerated along a line with acceleration  $a(t) = t$ . What is its velocity  $v(t)$  at time  $t$ ?

**One possible answer:**  $v(t) = \frac{t^2}{2}$  (particle starts from rest).

**Another possible answer:**  $v(t) = \frac{t^2}{2} - 37$  (particle starts with velocity  $-37$ ).

$$v(t) = \frac{t^2}{2} + c \quad c \text{ any constant}$$

**Definition:**  $f$  is an antiderivative of  $g$  if  $f' = g$  (i.e.  $f'(x) = g(x)$  for all  $x$ ).

**Uniqueness:** "Antiderivatives are unique up to addition of a constant".

If  $f_1$  and  $f_2$  are antiderivatives of  $g$ , then for some constant  $c$ ,  $f_2(x) = f_1(x) + c$ .

**Proof:**

$$(f_2(x) - f_1(x))' = f_2'(x) - f_1'(x) = g(x) - g(x) = 0$$

so (by MVT)  $f_2(x) - f_1(x)$  is constant, say  $f_2(x) - f_1(x) = c$  for all  $x$ .


So knowing the value of an antiderivative at a single point determines the antiderivative uniquely.

**Example:** If  $f(x)$  is an antiderivative of  $\sin(x)$ , and  $f(\pi) = 0$ , what is  $f(x)$ ?

$$\frac{d}{dx}(-\cos(x)) = \sin(x) \quad \text{but} \quad -\cos(\pi) = 1$$

$$\text{so let } f(x) = -1 - \cos(x)$$

$$\text{then } f'(x) = \sin(x), \quad f(\pi) = -1 + 1 = 0$$



**Solution:** We know that  $-\cos(x)$  is an antiderivative of  $\sin(x)$ , so

$$f(x) = -\cos(x) + c$$

for some real number  $c$ .

$$\text{Now } 0 = f(\pi) = -\cos(\pi) + c = -(-1) + c = 1 + c, \text{ so } c = -1.$$

$$\text{So } f(x) = -\cos(x) - 1.$$

**Example:** A lunar astronaut throws a ball; it experiences vertical acceleration  $y''(t) = -1.62$ , but no air resistance. If the initial vertical velocity is  $y'(0) = 1$  and the initial horizontal velocity is  $x'(0) = 2$ , and its initial position is  $(x(0), y(0)) = (0, 1)$ , and the ground is at  $y = 0$ , what is the ball's trajectory? How high does it go, and when and where does it hit the ground?

**Solution:**  $y''(t) = -1.62$ , so  $y'(t) = -1.62t + c_1$ . But  $y'(0) = 1$ , so  $c_1 = 1$  and  $y'(t) = 1 - 1.62t$ . So  $y(t) = t - 0.81t^2 + c_2$ , and since  $y(0) = 1$  we have  $y(t) = 1 + t - 0.81t^2$ .

Meanwhile,  $x''(t) = 0$ , so  $x'(t) = x'(0) = 2$ , so  $x(t) = 2t + c_3$ , and  $x(0) = 0$  so  $x(t) = 2t$ . So

$$(x(t), y(t)) = (2t, 1 + t - 0.81t^2)$$

Max height:  $y'(t) = 1 - 1.62t = 0 \Leftrightarrow t = \frac{1}{1.62} = 0.62$ , so at  $t = 0.62$  the ball achieves its maximum height  $y(0.62) = 1 + 0.62 - 0.81(0.62)^2 = 1.31$ . Hits ground:  $y(t) = 0 \Leftrightarrow t = \frac{-1 \pm \sqrt{1^2 + 4(0.81)}}{2(-0.81)} \Leftrightarrow t = -0.65$  or  $t = 1.89$ ; negative time is irrelevant to the problem; the ball hits the ground at  $t = 1.89$ ,  $x = 2(1.89) = 3.78$ .

Some antiderivatives:

g(x) antiderivatives of g(x)

$x$	$x$
$x^2$	$x^2$
$x^3$	$x^3$
$x^{n+1}$	$x^{n+1}$
$\frac{1}{x}$	$\ln x  + c$
$\frac{1}{x^2}$	$-\frac{1}{x} + c$
$\frac{1}{x^3}$	$-\frac{1}{2x^2} + c$
$\frac{1}{x^4}$	$-\frac{1}{3x^3} + c$
$\frac{1}{x^5}$	$-\frac{1}{4x^4} + c$
$\frac{1}{x^6}$	$-\frac{1}{5x^5} + c$
$\frac{1}{x^7}$	$-\frac{1}{6x^6} + c$
$\frac{1}{x^8}$	$-\frac{1}{7x^7} + c$
$\frac{1}{x^9}$	$-\frac{1}{8x^8} + c$
$\frac{1}{x^{10}}$	$-\frac{1}{9x^9} + c$
$\frac{1}{x^{11}}$	$-\frac{1}{10x^{10}} + c$
$\frac{1}{x^{12}}$	$-\frac{1}{11x^{11}} + c$
$\frac{1}{x^{13}}$	$-\frac{1}{12x^{12}} + c$
$\frac{1}{x^{14}}$	$-\frac{1}{13x^{13}} + c$
$\frac{1}{x^{15}}$	$-\frac{1}{14x^{14}} + c$
$\frac{1}{x^{16}}$	$-\frac{1}{15x^{15}} + c$
$\frac{1}{x^{17}}$	$-\frac{1}{16x^{16}} + c$
$\frac{1}{x^{18}}$	$-\frac{1}{17x^{17}} + c$
$\frac{1}{x^{19}}$	$-\frac{1}{18x^{18}} + c$
$\frac{1}{x^{20}}$	$-\frac{1}{19x^{19}} + c$