

Fraïssé Theory for C^* -algebras

Alessandro Vignati
York University, Toronto

ASL Winter Meeting, Boise, Idaho
March, 21st, 2017

- 1 Introduction
- 2 \mathcal{Z} and \mathcal{W}
- 3 Connections with quantifier elimination

1 Introduction

2 \mathcal{Z} and \mathcal{W}

3 Connections with quantifier elimination

The first two slides of this talk can be taken from Bradd's talk. In particular, we will work in the setting of C^* -algebras.

Definition

A trace τ on a C^* -algebra A is a linear functional $\tau: A \rightarrow \mathbb{C}$ with $\|\tau\| = 1$, $\tau(aa^*) \geq 0$ and $\tau(ab) = \tau(ba)$ for all $a, b \in A$. τ is faithful if $\tau(aa^*) = 0$ implies $a = 0$.

If A and B are C^* -algebras, σ, τ traces on A and B resp., and ϕ is such that $\tau(\phi(a)) = \sigma(a)$ for all $a \in A$, ϕ is said trace preserving, denoted $\phi: (A, \sigma) \rightarrow (B, \tau)$. If A and B are unital the pullback of a trace is a trace, so every unital embedding is trace preserving for some traces.

If we consider the language of C^* -algebra together with a trace, trace preserving injections are exactly our embeddings.

The first two slides of this talk can be taken from Bradd's talk. In particular, we will work in the setting of C^* -algebras.

Definition

A trace τ on a C^* -algebra A is a linear functional $\tau: A \rightarrow \mathbb{C}$ with $\|\tau\| = 1$, $\tau(aa^*) \geq 0$ and $\tau(ab) = \tau(ba)$ for all $a, b \in A$. τ is faithful if $\tau(aa^*) = 0$ implies $a = 0$.

If A and B are C^* -algebras, σ, τ traces on A and B resp., and ϕ is such that $\tau(\phi(a)) = \sigma(a)$ for all $a \in A$, ϕ is said trace preserving, denoted $\phi: (A, \sigma) \rightarrow (B, \tau)$. If A and B are unital the pullback of a trace is a trace, so every unital embedding is trace preserving for some traces.

If we consider the language of C^* -algebra together with a trace, trace preserving injections are exactly our embeddings.

The first two slides of this talk can be taken from Bradd's talk. In particular, we will work in the setting of C^* -algebras.

Definition

A trace τ on a C^* -algebra A is a linear functional $\tau: A \rightarrow \mathbb{C}$ with $\|\tau\| = 1$, $\tau(aa^*) \geq 0$ and $\tau(ab) = \tau(ba)$ for all $a, b \in A$. τ is faithful if $\tau(aa^*) = 0$ implies $a = 0$.

If A and B are C^* -algebras, σ, τ traces on A and B resp., and ϕ is such that $\tau(\phi(a)) = \sigma(a)$ for all $a \in A$, ϕ is said trace preserving, denoted $\phi: (A, \sigma) \rightarrow (B, \tau)$. If A and B are unital the pullback of a trace is a trace, so every unital embedding is trace preserving for some traces.

If we consider the language of C^* -algebra together with a trace, trace preserving injections are exactly our embeddings.

The first two slides of this talk can be taken from Bradd's talk. In particular, we will work in the setting of C^* -algebras.

Definition

A trace τ on a C^* -algebra A is a linear functional $\tau: A \rightarrow \mathbb{C}$ with $\|\tau\| = 1$, $\tau(aa^*) \geq 0$ and $\tau(ab) = \tau(ba)$ for all $a, b \in A$. τ is faithful if $\tau(aa^*) = 0$ implies $a = 0$.

If A and B are C^* -algebras, σ, τ traces on A and B resp., and ϕ is such that $\tau(\phi(a)) = \sigma(a)$ for all $a \in A$, ϕ is said trace preserving, denoted $\phi: (A, \sigma) \rightarrow (B, \tau)$. If A and B are unital the pullback of a trace is a trace, so every unital embedding is trace preserving for some traces.

If we consider the language of C^* -algebra together with a trace, trace preserving injections are exactly our embeddings.

Let \mathcal{L} be a language including the language of C^* -algebras, and \mathcal{K} be a class of finitely generated \mathcal{L} -structures considered with distinguished generators. \mathcal{K} is said a Fraïssé class if it satisfies:

- the JEP: for all $A, B \in \mathcal{K}$ there is $C \in \mathcal{K}$ in which A and B both embed;
- the NAP: if A, B_1, B_2 are in \mathcal{K} and $\phi_i: A \rightarrow B_i$ are \mathcal{L} -embeddings, $F \subseteq A$ is finite and $\epsilon > 0$ then there are $C \in \mathcal{K}$ and $\psi_i: B_i \rightarrow C$ such that

$$\|\psi_1 \circ \phi_1(a) - \psi_2 \circ \phi_2(a)\| < \epsilon, \text{ whenever } a \in F$$

Let \mathcal{K}_n be the space of all n -generated elements of \mathcal{K} , $A, B \in \mathcal{K}_n$ with distinguished generators \bar{a} and \bar{b} . Consider the pseudo-metric

$$d_n(A, B) = \inf_{C, \phi, \psi} \sup_{i \leq n} \|\phi(\bar{a}_i) - \psi(\bar{b}_i)\|_C$$

where C is quantified in \mathcal{K} and ψ, ϕ quantify over all embeddings $\phi: A \rightarrow C, \psi: B \rightarrow C$.

- the WPP: the \mathcal{K}_n considered with the pseudo-metric d_n is separable, for all n .

Let \mathcal{L} be a language including the language of C^* -algebras, and \mathcal{K} be a class of finitely generated \mathcal{L} -structures considered with distinguished generators. \mathcal{K} is said a Fraïssé class if it satisfies:

- the JEP: for all $A, B \in \mathcal{K}$ there is $C \in \mathcal{K}$ in which A and B both embed;
- the NAP: if A, B_1, B_2 are in \mathcal{K} and $\phi_i: A \rightarrow B_i$ are \mathcal{L} -embeddings, $F \subseteq A$ is finite and $\epsilon > 0$ then there are $C \in \mathcal{K}$ and $\psi_i: B_i \rightarrow C$ such that

$$\|\psi_1 \circ \phi_1(a) - \psi_2 \circ \phi_2(a)\| < \epsilon, \text{ whenever } a \in F$$

Let \mathcal{K}_n be the space of all n -generated elements of \mathcal{K} , $A, B \in \mathcal{K}_n$ with distinguished generators \bar{a} and \bar{b} . Consider the pseudo-metric

$$d_n(A, B) = \inf_{C, \phi, \psi} \sup_{i \leq n} \|\phi(\bar{a}_i) - \psi(\bar{b}_i)\|_C$$

where C is quantified in \mathcal{K} and ψ, ϕ quantify over all embeddings $\phi: A \rightarrow C, \psi: B \rightarrow C$.

- the WPP: the \mathcal{K}_n considered with the pseudo-metric d_n is separable, for all n .

Let \mathcal{L} be a language including the language of C^* -algebras, and \mathcal{K} be a class of finitely generated \mathcal{L} -structures considered with distinguished generators. \mathcal{K} is said a Fraïssé class if it satisfies:

- the JEP: for all $A, B \in \mathcal{K}$ there is $C \in \mathcal{K}$ in which A and B both embed;
- the NAP: if A, B_1, B_2 are in \mathcal{K} and $\phi_i: A \rightarrow B_i$ are \mathcal{L} -embeddings, $F \subseteq A$ is finite and $\epsilon > 0$ then there are $C \in \mathcal{K}$ and $\psi_i: B_i \rightarrow C$ such that

$$\|\psi_1 \circ \phi_1(a) - \psi_2 \circ \phi_2(a)\| < \epsilon, \text{ whenever } a \in F$$

Let \mathcal{K}_n be the space of all n -generated elements of \mathcal{K} , $A, B \in \mathcal{K}_n$ with distinguished generators \bar{a} and \bar{b} . Consider the pseudo-metric

$$d_n(A, B) = \inf_{C, \phi, \psi} \sup_{i \leq n} \|\phi(\bar{a}_i) - \psi(\bar{b}_i)\|_C$$

where C is quantified in \mathcal{K} and ψ, ϕ quantify over all embeddings $\phi: A \rightarrow C, \psi: B \rightarrow C$.

- the WPP: the \mathcal{K}_n considered with the pseudo-metric d_n is separable, for all n .

If \mathcal{K} is a Fraïssé class, and M is a limit of structures in \mathcal{K} , M is called a \mathcal{K} -structure. A \mathcal{K} -structure is

- \mathcal{K} -universal if every element of \mathcal{K} embeds in M
- approximately \mathcal{K} -homogeneous if for all $A \in \mathcal{K}$, $F \subseteq A$ finite, $\epsilon > 0$ and $\phi_1, \phi_2: A \rightarrow M$ embeddings there is an automorphism ρ of M with

$$\|\rho(\phi_1(a)) - \phi_2(a)\| < \epsilon, a \in F$$

If \mathcal{K} is a Fraïssé class and M is a separable \mathcal{K} -structure which is both \mathcal{K} -universal and approximately \mathcal{K} -homogeneous, M is said a Fraïssé limit.

Theorem (Ben Yaacov)

If \mathcal{K} is a Fraïssé class and its Fraïssé limit exists, it is unique up to isomorphism.

If \mathcal{K} is a Fraïssé class, and M is a limit of structures in \mathcal{K} , M is called a \mathcal{K} -structure. A \mathcal{K} -structure is

- \mathcal{K} -universal if every element of \mathcal{K} embeds in M
- approximately \mathcal{K} -homogeneous if for all $A \in \mathcal{K}$, $F \subseteq A$ finite, $\epsilon > 0$ and $\phi_1, \phi_2: A \rightarrow M$ embeddings there is an automorphism ρ of M with

$$\|\rho(\phi_1(a)) - \phi_2(a)\| < \epsilon, a \in F$$

If \mathcal{K} is a Fraïssé class and M is a separable \mathcal{K} -structure which is both \mathcal{K} -universal and approximately \mathcal{K} -homogeneous, M is said a Fraïssé limit.

Theorem (Ben Yaacov)

If \mathcal{K} is a Fraïssé class and its Fraïssé limit exists, it is unique up to isomorphism.

- Introduction
- \mathcal{Z} and \mathcal{W}
- Connections with quantifier elimination

If p and q are coprime, let

$$Z_{p,q} = \{f \in C([0, 1], M_p \otimes M_q) \mid f(0) \in M_p \otimes 1, f(1) \in 1 \otimes M_q\}.$$

$Z_{p,q}$ are called dimension drop algebras. Traces of $Z_{p,q}$ correspond to probability measures on $[0, 1]$ by

$$\tau_\mu(f) = \int_0^1 \tau(f(t)) d\mu(t).$$

All traces are of this form. If μ is diffuse, τ is said diffuse. Every (nonzero) $*$ -homomorphism $\phi: Z_{p,q} \rightarrow Z_{p',q'}$ is trace preserving for some traces σ and τ .

Proposition

- pq divides $p'q'$ if and only if there is an embedding $Z_{p,q} \rightarrow Z_{p',q'}$.
- Let σ, τ be faithful traces on $Z_{p,q}$. If τ is diffuse, there is an embedding $(Z_{p,q}, \sigma) \rightarrow (Z_{p,q}, \tau)$.
- If also σ is diffuse, ϕ can be chosen to be an isomorphism.

If p and q are coprime, let

$$Z_{p,q} = \{f \in C([0, 1], M_p \otimes M_q) \mid f(0) \in M_p \otimes 1, f(1) \in 1 \otimes M_q\}.$$

$Z_{p,q}$ are called dimension drop algebras. Traces of $Z_{p,q}$ correspond to probability measures on $[0, 1]$ by

$$\tau_\mu(f) = \int_0^1 \tau(f(t)) d\mu(t).$$

All traces are of this form. If μ is diffuse, τ is said diffuse. Every (nonzero) $*$ -homomorphism $\phi: Z_{p,q} \rightarrow Z_{p',q'}$ is trace preserving for some traces σ and τ .

Proposition

- pq divides $p'q'$ if and only if there is an embedding $Z_{p,q} \rightarrow Z_{p',q'}$.
- Let σ, τ be faithful traces on $Z_{p,q}$. If τ is diffuse, there is an embedding $(Z_{p,q}, \sigma) \rightarrow (Z_{p,q}, \tau)$.
- If also σ is diffuse, ϕ can be chosen to be an isomorphism.

If p and q are coprime, let

$$Z_{p,q} = \{f \in C([0, 1], M_p \otimes M_q) \mid f(0) \in M_p \otimes 1, f(1) \in 1 \otimes M_q\}.$$

$Z_{p,q}$ are called dimension drop algebras. Traces of $Z_{p,q}$ correspond to probability measures on $[0, 1]$ by

$$\tau_\mu(f) = \int_0^1 \tau(f(t)) d\mu(t).$$

All traces are of this form. If μ is diffuse, τ is said diffuse. Every (nonzero) $*$ -homomorphism $\phi: Z_{p,q} \rightarrow Z_{p',q'}$ is trace preserving for some traces σ and τ .

Proposition

- pq divides $p'q'$ if and only if there is an embedding $Z_{p,q} \rightarrow Z_{p',q'}$.
- Let σ, τ be faithful traces on $Z_{p,q}$. If τ is diffuse, there is an embedding $(Z_{p,q}, \sigma) \rightarrow (Z_{p,q}, \tau)$.
- If also σ is diffuse, ϕ can be chosen to be an isomorphism.

Theorem (Jiang-Su)

There are increasing sequences of coprimes p_i, q_i and maps $\phi_i: Z_{p_i, q_i} \rightarrow Z_{p_{i+1}, q_{i+1}}$ such that $\mathcal{Z} = \lim_i (Z_{p_i, q_i}, \phi_i)$ is simple, monotracial and has the same K -theory as \mathbb{C} . Let p_i, q_i and ψ_i and $\mathcal{A} = \lim (Z_{p_i, q_i}, \psi_i)$. If \mathcal{A} is simple monotracial and has the same K -theory as \mathcal{Z} , then $\mathcal{A} \cong \mathcal{Z}$.

Jiang and Su's \mathcal{Z} is pivotal in the classification programme of C^* -algebras. \mathcal{Z} is unique and universal in many senses. It is self-absorbing ($\mathcal{Z} \otimes \mathcal{Z} \cong \mathcal{Z}$) in a very strong sense. The (amenable) algebras that have the property of absorbing \mathcal{Z} (i.e., $A \otimes \mathcal{Z} \cong A$) are the ones for which there are hopes of obtaining classification.

Theorem (Jiang-Su)

There are increasing sequences of coprimes p_i, q_i and maps $\phi_i: Z_{p_i, q_i} \rightarrow Z_{p_{i+1}, q_{i+1}}$ such that $\mathcal{Z} = \lim_i (Z_{p_i, q_i}, \phi_i)$ is simple, monotracial and has the same K -theory as \mathbb{C} . Let p_i, q_i and ψ_i and $\mathcal{A} = \lim (Z_{p_i, q_i}, \psi_i)$. If \mathcal{A} is simple monotracial and has the same K -theory as \mathcal{Z} , then $\mathcal{A} \cong \mathcal{Z}$.

Jiang and Su's \mathcal{Z} is pivotal in the classification programme of C^* -algebras. \mathcal{Z} is unique and universal in many senses. It is self-absorbing ($\mathcal{Z} \otimes \mathcal{Z} \cong \mathcal{Z}$) in a very strong sense. The (amenable) algebras that have the property of absorbing \mathcal{Z} (i.e., $A \otimes \mathcal{Z} \cong A$) are the ones for which there are hopes of obtaining classification.

Theorem (Jiang-Su)

There are increasing sequences of coprimes p_i, q_i and maps $\phi_i: Z_{p_i, q_i} \rightarrow Z_{p_{i+1}, q_{i+1}}$ such that $\mathcal{Z} = \lim_i (Z_{p_i, q_i}, \phi_i)$ is simple, monotracial and has the same K -theory as \mathbb{C} . Let p_i, q_i and ψ_i and $\mathcal{A} = \lim (Z_{p_i, q_i}, \psi_i)$. If \mathcal{A} is simple monotracial and has the same K -theory as \mathcal{Z} , then $\mathcal{A} \cong \mathcal{Z}$.

Jiang and Su's \mathcal{Z} is pivotal in the classification programme of C^* -algebras. \mathcal{Z} is unique and universal in many senses. It is self-absorbing ($\mathcal{Z} \otimes \mathcal{Z} \cong \mathcal{Z}$) in a very strong sense. The (amenable) algebras that have the property of absorbing \mathcal{Z} (i.e., $A \otimes \mathcal{Z} \cong A$) are the ones for which there are hopes of obtaining classification.

Theorem (Jiang-Su)

There are increasing sequences of coprimes p_i, q_i and maps $\phi_i: Z_{p_i, q_i} \rightarrow Z_{p_{i+1}, q_{i+1}}$ such that $\mathcal{Z} = \lim_i (Z_{p_i, q_i}, \phi_i)$ is simple, monotracial and has the same K -theory as \mathbb{C} . Let p_i, q_i and ψ_i and $\mathcal{A} = \lim (Z_{p_i, q_i}, \psi_i)$. If \mathcal{A} is simple monotracial and has the same K -theory as \mathcal{Z} , then $\mathcal{A} \cong \mathcal{Z}$.

Jiang and Su's \mathcal{Z} is pivotal in the classification programme of C^* -algebras. \mathcal{Z} is unique and universal in many senses. It is self-absorbing ($\mathcal{Z} \otimes \mathcal{Z} \cong \mathcal{Z}$) in a very strong sense. The (amenable) algebras that have the property of absorbing \mathcal{Z} (i.e., $A \otimes \mathcal{Z} \cong A$) are the ones for which there are hopes of obtaining classification.

Theorem (EFHKKL, Masumoto)

The class $\{(Z_{p,q}, \tau) \mid p, q \text{ coprimes}, \tau \text{ faithful trace}\}$ is a Fraïssé class. \mathcal{Z} is its Fraïssé limit.

So for all p, q coprimes, $\phi_1, \phi_2: Z_{p,q} \rightarrow \mathcal{Z}$, $F \subseteq Z_{p,q}$ and $\epsilon > 0$ there is an automorphism ρ of \mathcal{Z} such that

$$\|\rho(\phi_1(a)) - \phi_2(a)\| < \epsilon$$

for all $a \in F$.

EFHKKL's Proof is based on some known facts about \mathcal{Z} . Masumoto's one on a careful study of what the maps between dimension drop algebras can be.

Theorem (EFHKKL, Masumoto)

The class $\{(Z_{p,q}, \tau) \mid p, q \text{ coprimes}, \tau \text{ faithful trace}\}$ is a Fraïssé class. \mathcal{Z} is its Fraïssé limit.

So for all p, q coprimes, $\phi_1, \phi_2: Z_{p,q} \rightarrow \mathcal{Z}$, $F \subseteq Z_{p,q}$ and $\epsilon > 0$ there is an automorphism ρ of \mathcal{Z} such that

$$\|\rho(\phi_1(a)) - \phi_2(a)\| < \epsilon$$

for all $a \in F$.

EFHKKL's Proof is based on some known facts about \mathcal{Z} . Masumoto's one on a careful study of what the maps between dimension drop algebras can be.

Theorem (EFHKKL, Masumoto)

The class $\{(Z_{p,q}, \tau) \mid p, q \text{ coprimes}, \tau \text{ faithful trace}\}$ is a Fraïssé class. \mathcal{Z} is its Fraïssé limit.

So for all p, q coprimes, $\phi_1, \phi_2: Z_{p,q} \rightarrow \mathcal{Z}$, $F \subseteq Z_{p,q}$ and $\epsilon > 0$ there is an automorphism ρ of \mathcal{Z} such that

$$\|\rho(\phi_1(a)) - \phi_2(a)\| < \epsilon$$

for all $a \in F$.

EFHKKL's Proof is based on some known facts about \mathcal{Z} . Masumoto's one on a careful study of what the maps between dimension drop algebras can be.

Question

As we know that $\mathcal{Z} \cong \mathcal{Z} \otimes \mathcal{Z}$, what can we say about maps $\phi_1, \phi_2: Z_{p,q} \otimes Z_{p',q'} \rightarrow \mathcal{Z}$? In other terms, can we prove that \mathcal{Z} and $\mathcal{Z} \otimes \mathcal{Z}$ are the Fraïssé limit of the same class?

This is more difficult than one can think. In fact, maps $Z_{p,q} \otimes Z_{p',q'} \rightarrow Z_{p'',q''}$ are more complicated than one can think. Despite that, maps $Z_{p,q} \otimes Z_{p',q'} \rightarrow Z_{p'',q''}$ are always well behaved.

Theorem (Jacelon-V.)

The class $\{(Z_{p,q}, \tau), (Z_{p,q} \otimes Z_{p',q'}, \sigma)\}$ is a Fraïssé class. Both \mathcal{Z} and $\mathcal{Z} \otimes \mathcal{Z}$ are its Fraïssé limits, so $\mathcal{Z} \cong \mathcal{Z} \otimes \mathcal{Z}$.

Question

As we know that $\mathcal{Z} \cong \mathcal{Z} \otimes \mathcal{Z}$, what can we say about maps $\phi_1, \phi_2: Z_{p,q} \otimes Z_{p',q'} \rightarrow \mathcal{Z}$? In other terms, can we prove that \mathcal{Z} and $\mathcal{Z} \otimes \mathcal{Z}$ are the Fraïssé limit of the same class?

This is more difficult than one can think. In fact, maps $Z_{p,q} \otimes Z_{p',q'} \rightarrow Z_{p'',q''}$ are more complicated than one can think. Despite that, maps $Z_{p,q} \otimes Z_{p',q'} \rightarrow Z_{p'',q''}$ are always well behaved.

Theorem (Jacelon-V.)

The class $\{(Z_{p,q}, \tau), (Z_{p,q} \otimes Z_{p',q'}, \sigma)\}$ is a Fraïssé class. Both \mathcal{Z} and $\mathcal{Z} \otimes \mathcal{Z}$ are its Fraïssé limits, so $\mathcal{Z} \cong \mathcal{Z} \otimes \mathcal{Z}$.

Question

As we know that $\mathcal{Z} \cong \mathcal{Z} \otimes \mathcal{Z}$, what can we say about maps $\phi_1, \phi_2: Z_{p,q} \otimes Z_{p',q'} \rightarrow \mathcal{Z}$? In other terms, can we prove that \mathcal{Z} and $\mathcal{Z} \otimes \mathcal{Z}$ are the Fraïssé limit of the same class?

This is more difficult than one can think. In fact, maps $Z_{p,q} \otimes Z_{p',q'} \rightarrow Z_{p'',q''}$ are more complicated than one can think. Despite that, maps $Z_{p,q} \otimes Z_{p',q'} \rightarrow Z_{p'',q''}$ are always well behaved.

Theorem (Jacelon-V.)

The class $\{(Z_{p,q}, \tau), (Z_{p,q} \otimes Z_{p',q'}, \sigma)\}$ is a Fraïssé class. Both \mathcal{Z} and $\mathcal{Z} \otimes \mathcal{Z}$ are its Fraïssé limits, so $\mathcal{Z} \cong \mathcal{Z} \otimes \mathcal{Z}$.

Let

$$A_{n,k} = C([0, 1], M_n \otimes M_k) \mid f(0) = 1 \otimes c, f(1) = 1_{n-1} \otimes c\}$$

These are called Razak's blocks. Traces are, as before, given by probability measures on the interval. The absence of the unit doesn't allow to say that every embedding of $A_{n,k} \rightarrow A_{n',k'}$ is trace preserving.

Proposition

- If σ, τ are faithful diffuse traces on $A_{n,k}$ then there is an isomorphism $(A_{n,k}, \sigma) \rightarrow (A_{n,k}, \tau)$
- If $p \geq 2$ and τ, σ are faithful traces on $A_{n,k}$ and $A_{pn, (pn-1)k}$, τ being diffuse, there is a trace preserving embedding $\phi: (A_{n,k}, \sigma) \rightarrow (A_{pn, (pn-1)k}, \tau)$.

Let

$$A_{n,k} = C([0, 1], M_n \otimes M_k) \mid f(0) = 1 \otimes c, f(1) = 1_{n-1} \otimes c\}$$

These are called Razak's blocks. Traces are, as before, given by probability measures on the interval. The absence of the unit doesn't allow to say that every embedding of $A_{n,k} \rightarrow A_{n',k'}$ is trace preserving.

Proposition

- If σ, τ are faithful diffuse traces on $A_{n,k}$ then there is an isomorphism $(A_{n,k}, \sigma) \rightarrow (A_{n,k}, \tau)$
- If $p \geq 2$ and τ, σ are faithful traces on $A_{n,k}$ and $A_{pn, (pn-1)k}$, τ being diffuse, there is a trace preserving embedding $\phi: (A_{n,k}, \sigma) \rightarrow (A_{pn, (pn-1)k}, \tau)$.

Let

$$A_{n,k} = C([0, 1], M_n \otimes M_k) \mid f(0) = 1 \otimes c, f(1) = 1_{n-1} \otimes c\}$$

These are called Razak's blocks. Traces are, as before, given by probability measures on the interval. The absence of the unit doesn't allow to say that every embedding of $A_{n,k} \rightarrow A_{n',k'}$ is trace preserving.

Proposition

- If σ, τ are faithful diffuse traces on $A_{n,k}$ then there is an isomorphism $(A_{n,k}, \sigma) \rightarrow (A_{n,k}, \tau)$
- If $p \geq 2$ and τ, σ are faithful traces on $A_{n,k}$ and $A_{pn, (pn-1)k}$, τ being diffuse, there is a trace preserving embedding $\phi: (A_{n,k}, \sigma) \rightarrow (A_{pn, (pn-1)k}, \tau)$.

Theorem (Jacelon)

There are increasing sequences n_i, k_i and trace preserving (for some traces) maps $\phi_i: A_{n_i, k_i} \rightarrow A_{n_{i+1}, k_{i+1}}$ such that $\mathcal{W} = \lim_i (A_{n_i, k_i}, \phi_i)$ is simple, stably projectionless monotracial and with trivial K -theory.

Let n_i, k_i and ψ_i and $A = \lim (A_{n_i, k_i}, \psi_i)$. If A is simple monotracial stably projectionless and with trivial K -theory, then $A \cong \mathcal{W}$.

Recent work of Elliott-Niu and Gong-Lin showed the first evidences that \mathcal{W} plays the same role in the classification of nonunital algebras as \mathcal{Z} does for the unital case. \mathcal{W} is a universal objects in many ways. On the other hand, that $\mathcal{W} \otimes \mathcal{W} \cong \mathcal{W}$ was only recently proved, involving a long and complicated proof in classification theory.

Theorem (Jacelon)

There are increasing sequences n_i, k_i and trace preserving (for some traces) maps $\phi_i: A_{n_i, k_i} \rightarrow A_{n_{i+1}, k_{i+1}}$ such that $\mathcal{W} = \lim_i (A_{n_i, k_i}, \phi_i)$ is simple, stably projectionless monotracial and with trivial K -theory. Let n_i, k_i and ψ_i and $A = \lim (A_{n_i, k_i}, \psi_i)$. If A is simple monotracial stably projectionless and with trivial K -theory, then $A \cong \mathcal{W}$.

Recent work of Elliott-Niu and Gong-Lin showed the first evidences that \mathcal{W} plays the same role in the classification of nonunital algebras as \mathcal{Z} does for the unital case. \mathcal{W} is a universal objects in many ways. On the other hand, that $\mathcal{W} \otimes \mathcal{W} \cong \mathcal{W}$ was only recently proved, involving a long and complicated proof in classification theory.

Theorem (Jacelon)

There are increasing sequences n_i, k_i and trace preserving (for some traces) maps $\phi_i: A_{n_i, k_i} \rightarrow A_{n_{i+1}, k_{i+1}}$ such that $\mathcal{W} = \lim_i (A_{n_i, k_i}, \phi_i)$ is simple, stably projectionless monotracial and with trivial K -theory. Let n_i, k_i and ψ_i and $A = \lim (A_{n_i, k_i}, \psi_i)$. If A is simple monotracial stably projectionless and with trivial K -theory, then $A \cong \mathcal{W}$.

Recent work of Elliott-Niu and Gong-Lin showed the first evidences that \mathcal{W} plays the same role in the classification of nonunital algebras as \mathcal{Z} does for the unital case. \mathcal{W} is a universal objects in many ways. On the other hand, that $\mathcal{W} \otimes \mathcal{W} \cong \mathcal{W}$ was only recently proved, involving a long and complicated proof in classification theory.

Theorem (Jacelon-V.)

The class $\{(A_{n,k}, \sigma) \mid \sigma \text{ is a faithful trace}\}$ is a Fraïssé class. \mathcal{W} is its limit.

Whenever $\phi_1, \phi_2: (A_{n,k}, \sigma) \rightarrow (\mathcal{W}, \tau)$ are embeddings for some faithful diffuse σ and $F \subseteq A_{n,k}$ and $\epsilon > 0$ are given there is an automorphism ρ of \mathcal{W} such that

$$\|\rho(\phi_1(a)) - \phi_2(a)\| < \epsilon$$

for $a \in F$.

The techniques used are similar to the ones of Masumoto's proof that \mathcal{Z} is a Fraïssé limit.

Theorem (Jacelon-V.)

The class $\{(A_{n,k}, \sigma) \mid \sigma \text{ is a faithful trace}\}$ is a Fraïssé class. \mathcal{W} is its limit.

Whenever $\phi_1, \phi_2: (A_{n,k}, \sigma) \rightarrow (\mathcal{W}, \tau)$ are embeddings for some faithful diffuse σ and $F \subseteq A_{n,k}$ and $\epsilon > 0$ are given there is an automorphism ρ of \mathcal{W} such that

$$\|\rho(\phi_1(a)) - \phi_2(a)\| < \epsilon$$

for $a \in F$.

The techniques used are similar to the ones of Masumoto's proof that \mathcal{Z} is a Fraïssé limit.

Theorem (Jacelon-V.)

The class $\{(A_{n,k}, \sigma) \mid \sigma \text{ is a faithful trace}\}$ is a Fraïssé class. \mathcal{W} is its limit.

Whenever $\phi_1, \phi_2: (A_{n,k}, \sigma) \rightarrow (\mathcal{W}, \tau)$ are embeddings for some faithful diffuse σ and $F \subseteq A_{n,k}$ and $\epsilon > 0$ are given there is an automorphism ρ of \mathcal{W} such that

$$\|\rho(\phi_1(a)) - \phi_2(a)\| < \epsilon$$

for $a \in F$.

The techniques used are similar to the ones of Masumoto's proof that \mathcal{Z} is a Fraïssé limit.

The goal is to reproduce the argument used for $\{(Z_{p,q}, \tau), (Z_{p,q} \otimes Z_{p',q'}, \sigma)\}$ to show that $\{(A_{n,k}, \sigma), (A_{n,k} \otimes A_{n',k'}, \tau)\}$ is a Fraïssé class.

Proposition

Fix n, k .

- There are well-behaved maps $A_{n,k} \rightarrow A_{n',k'}$ for some n', k' .
- There are well-behaved maps $A_{n,k} \otimes A_{n,k} \rightarrow A_{n',k'}$ for some n', k' .

Problem

Not all maps $A_{n,k} \otimes A_{n,k} \rightarrow A_{n',k'}$ are well-behaved. Proving NAP seems difficult. Also, there is no well-behaved trace preserving map $A_{n,k} \rightarrow A_{n',k'} \otimes A_{n'',k''}$

Question

*Is there any trace preserving map $A_{n,k} \rightarrow A_{n',k'} \otimes A_{n',k'}$?
Do we need to add more structure and remove the trace?*

The goal is to reproduce the argument used for $\{(Z_{p,q}, \tau), (Z_{p,q} \otimes Z_{p',q'}, \sigma)\}$ to show that $\{(A_{n,k}, \sigma), (A_{n,k} \otimes A_{n',k'}, \tau)\}$ is a Fraïssé class.

Proposition

Fix n, k .

- There are well-behaved maps $A_{n,k} \rightarrow A_{n',k'}$ for some n', k' .
- There are well-behaved maps $A_{n,k} \otimes A_{n,k} \rightarrow A_{n',k'}$ for some n', k' .

Problem

Not all maps $A_{n,k} \otimes A_{n,k} \rightarrow A_{n',k'}$ are well-behaved. Proving NAP seems difficult. Also, there is no well-behaved trace preserving map $A_{n,k} \rightarrow A_{n',k'} \otimes A_{n',k'}$

Question

Is there any trace preserving map $A_{n,k} \rightarrow A_{n',k'} \otimes A_{n',k'}$?
Do we need to add more structure and remove the trace?

The goal is to reproduce the argument used for $\{(Z_{p,q}, \tau), (Z_{p,q} \otimes Z_{p',q'}, \sigma)\}$ to show that $\{(A_{n,k}, \sigma), (A_{n,k} \otimes A_{n',k'}, \tau)\}$ is a Fraïssé class.

Proposition

Fix n, k .

- There are well-behaved maps $A_{n,k} \rightarrow A_{n',k'}$ for some n', k' .
- There are well-behaved maps $A_{n,k} \otimes A_{n,k} \rightarrow A_{n',k'}$ for some n', k' .

Problem

Not all maps $A_{n,k} \otimes A_{n,k} \rightarrow A_{n',k'}$ are well-behaved. Proving NAP seems difficult. Also, there is no well-behaved trace preserving map $A_{n,k} \rightarrow A_{n',k'} \otimes A_{n'',k''}$

Question

Is there any trace preserving map $A_{n,k} \rightarrow A_{n',k'} \otimes A_{n',k'}$?
Do we need to add more structure and remove the trace?

The goal is to reproduce the argument used for $\{(Z_{p,q}, \tau), (Z_{p,q} \otimes Z_{p',q'}, \sigma)\}$ to show that $\{(A_{n,k}, \sigma), (A_{n,k} \otimes A_{n',k'}, \tau)\}$ is a Fraïssé class.

Proposition

Fix n, k .

- There are well-behaved maps $A_{n,k} \rightarrow A_{n',k'}$ for some n', k' .
- There are well-behaved maps $A_{n,k} \otimes A_{n,k} \rightarrow A_{n',k'}$ for some n', k' .

Problem

Not all maps $A_{n,k} \otimes A_{n,k} \rightarrow A_{n',k'}$ are well-behaved. Proving NAP seems difficult. Also, there is no well-behaved trace preserving map $A_{n,k} \rightarrow A_{n',k'} \otimes A_{n'',k''}$

Question

Is there any trace preserving map $A_{n,k} \rightarrow A_{n',k'} \otimes A_{n',k'}$?
Do we need to add more structure and remove the trace?

- Introduction
- \mathcal{Z} and \mathcal{W}
- **8** Connections with quantifier elimination

A separable C^* algebra A has quantifier elimination if, for all separable $B \equiv A$, all F finitely generated C^* -algebras, $\mathcal{U} \in \beta\mathbb{N} \setminus \mathbb{N}$ and embeddings $\phi: F \rightarrow B^{\mathcal{U}}$, $\iota: F \rightarrow B$ there is $\kappa: B \rightarrow B^{\mathcal{U}}$ such that the following commutes:

$$\begin{array}{ccc} F & \xrightarrow{\phi} & B^{\mathcal{U}} \\ & \searrow \iota & \nearrow \kappa \\ & B & \end{array}$$

A has property (\star) if for all F finitely generated C^* -algebras, $\mathcal{U} \in \beta\mathbb{N} \setminus \mathbb{N}$ and embeddings $\phi: F \rightarrow A^{\mathcal{U}}$, $\iota: F \rightarrow A$ there is $\kappa: A \rightarrow A^{\mathcal{U}}$ such that the following commutes:

$$\begin{array}{ccc} F & \xrightarrow{\phi} & A^{\mathcal{U}} \\ & \searrow \iota & \nearrow \kappa \\ & A & \end{array}$$

This property, for example, holds if $A = \mathcal{O}_2$ and F is assumed to be nuclear.

A has property (\star) if for all F finitely generated C^* -algebras, $\mathcal{U} \in \beta\mathbb{N} \setminus \mathbb{N}$ and embeddings $\phi: F \rightarrow A^{\mathcal{U}}$, $\iota: F \rightarrow A$ there is $\kappa: A \rightarrow A^{\mathcal{U}}$ such that the following commutes:

$$\begin{array}{ccc} F & \xrightarrow{\phi} & A^{\mathcal{U}} \\ & \searrow \iota & \nearrow \kappa \\ & A & \end{array}$$

This property, for example, holds if $A = \mathcal{O}_2$ and F is assumed to be nuclear.

Theorem (Eagle-Farah-Kirchberg-V., Eagle-Goldbring-V.)

- *In the language of unital C^* -algebras, the only C^* -algebras with QE are \mathbb{C} , \mathbb{C}^2 , $M_2(\mathbb{C})$ and $C(2^{\mathbb{N}})$.*
- *The only nonabelian C^* -algebra with (\star) is $M_2(\mathbb{C})$.*

One is tempted, therefore, to add a predicate such as the trace. If one considers the language of tracial unital C^* -algebras, then \mathcal{Z} has property (\star) if one restricts itself to those F of the form $F = \bigotimes_{i \leq n} Z_{p_i, q_i}$, where p_i and q_i are coprime for all i .

If A is a Fraïssé limit for the Fraïssé class \mathcal{K} , for all the examples we have, A satisfies (\star) whenever $F \in \mathcal{K}$. This is where the absence of HP kicks in.

Question

Does \mathcal{Z} satisfy (\star) in the language of tracial unital C^ -algebras? Does it have QE?*

Question

What is the largest class of algebras for which \mathcal{W} satisfies (\star) in the language of tracial C^ -algebras?*

One is tempted, therefore, to add a predicate such as the trace. If one considers the language of tracial unital C^* -algebras, then \mathcal{Z} has property (\star) if one restricts itself to those F of the form $F = \bigotimes_{i \leq n} Z_{p_i, q_i}$, where p_i and q_i are coprime for all i .

If A is a Fraïssé limit for the Fraïssé class \mathcal{K} , for all the examples we have, A satisfies (\star) whenever $F \in \mathcal{K}$. This is where the absence of HP kicks in.

Question

Does \mathcal{Z} satisfy (\star) in the language of tracial unital C^ -algebras? Does it have QE?*

Question

What is the largest class of algebras for which \mathcal{W} satisfies (\star) in the language of tracial C^ -algebras?*

One is tempted, therefore, to add a predicate such as the trace. If one considers the language of tracial unital C^* -algebras, then \mathcal{Z} has property (\star) if one restricts itself to those F of the form $F = \bigotimes_{i \leq n} Z_{p_i, q_i}$, where p_i and q_i are coprime for all i .

If A is a Fraïssé limit for the Fraïssé class \mathcal{K} , for all the examples we have, A satisfies (\star) whenever $F \in \mathcal{K}$. This is where the absence of HP kicks in.

Question

Does \mathcal{Z} satisfy (\star) in the language of tracial unital C^ -algebras? Does it have QE?*

Question

What is the largest class of algebras for which \mathcal{W} satisfies (\star) in the language of tracial C^ -algebras?*

One is tempted, therefore, to add a predicate such as the trace. If one considers the language of tracial unital C^* -algebras, then \mathcal{Z} has property (\star) if one restricts itself to those F of the form $F = \bigotimes_{i \leq n} Z_{p_i, q_i}$, where p_i and q_i are coprime for all i .

If A is a Fraïssé limit for the Fraïssé class \mathcal{K} , for all the examples we have, A satisfies (\star) whenever $F \in \mathcal{K}$. This is where the absence of HP kicks in.

Question

Does \mathcal{Z} satisfy (\star) in the language of tracial unital C^ -algebras? Does it have QE?*

Question

What is the largest class of algebras for which \mathcal{W} satisfies (\star) in the language of tracial C^ -algebras?*

Thank you!