

# ASSIGNMENT 6

1(a)  $M = k \cdot \frac{1}{p}$ , where  $k$  is a real number

(b) A function of the form  $f(x) = mx + b$ , where  $m$  and  $b$  are real numbers

(c) No, because of the  $x^{-1} = \frac{1}{x}$  term

(d) When  $b = 0$  (proportional = line through origin)

(e) linear:  $y = ax + b, b \neq 0$     proportional:  $y = ax$



ratio  $\frac{\text{output}}{\text{input}}$  not constant    ratio  $\frac{\text{output}}{\text{input}}$  is constant


(f)

x	y
0	5
1	7
2	9
3	11
4	13

← slope is constant, but ratios  $\frac{7}{1}, \frac{9}{2}, \frac{11}{3}, \frac{13}{4}$  are not equal


2(a)  $r < 0$





$x^{-4}$  decreases faster

3. 1986 ... 25,309 ...  $t=0$  (population in thousands)  
 1996 ... 28,847 ...  $t=10$




$$\text{slope} = \frac{28,847 - 25,309}{10} = 353.8$$

$$P(t) - 25,309 = 353.8(t - 0)$$

$$\text{so } P(t) = 25,309 + 353.8t$$

prediction for 2006:  $P(20) = 32,385$   
 (higher than actual population)

4. switch axes



$$\text{slope} = \frac{100 - 0}{212 - 32} = \frac{100}{180} = \frac{5}{9}$$

$$C - 0 = \frac{5}{9}(F - 32)$$

$$C = \frac{5}{9}(F - 32)$$

point-slope equation  
 using  $(32, 0)$

$$5. \quad \text{BMI} = \frac{m}{h^2} \frac{[\text{kg}]}{[\text{m}^2]} = \frac{2,20426 [\text{lb}]}{\left(\frac{100}{2,54}\right)^2 [\text{in}^2]}$$

$$= 0,0014221 \frac{[\text{lb}]}{[\text{in}^2]}$$

So if we want to have the same value in BMI numbers, then we need to multiply BMI by

$$1/0,0014221 = 703,18$$

$$\text{Thus:} \quad \text{BMI} = \frac{m}{h^2} \frac{[\text{kg}]}{[\text{m}^2]}$$

$$= 703,18 \cdot \frac{m}{h^2} \frac{[\text{lb}]}{[\text{in}^2]}$$

$$6. \quad \text{BMI}_A = \frac{m}{h^2}$$

$$\text{BMI}_B = \frac{m}{(1,05h)^2} = \frac{1}{1,05^2} \cdot \frac{m}{h^2} = \underline{\underline{0,907}} \text{ BMI}_A$$

$$\text{BMI}_C = \frac{0,95m}{h^2} = \underline{\underline{0,95}} \cdot \text{BMI}_A$$

So B has lower BMI

$$7. \quad (a) \quad T(B) = a \cdot \sqrt[3]{B} \quad \text{so}$$

$$T(2B) = a \cdot \sqrt[3]{2B} = \sqrt[3]{2} \cdot a \sqrt[3]{B} \approx 1,26 T(B)$$

So if the body mass doubles, the blood circulation time increases by a factor of 1,26 (ie, by 26%)

$$(b) \quad T(B) = a \cdot \sqrt[3]{B} \rightarrow 152 = a \sqrt[3]{5400}$$


$$\text{so } a = 152 / \sqrt[3]{5400} \approx 8.66$$

$$\text{so } T(B) = 8.66 \cdot \sqrt[3]{B}$$

$$(c) \quad T(0.1) = 8.66 \cdot \sqrt[3]{0.1} \approx 4.02 \text{ s} \rightarrow \text{smaller}$$

$$T(38,000) = 8.66 \cdot \sqrt[3]{38,000} \approx 291.15 \text{ s} \rightarrow \text{larger}$$

compared to  
example 1.1.13



8. (a) Surface area is proportional to volume raised to the power of  $2/3$  ( $S \propto V^{2/3}$ )

(b) When a baby grows to twice her size then the volume of her body (hence the mass) increases eight-fold. The strength of a bone is proportional to the cross-sectional area, and thus quadruples as the baby grows to twice her size. To compensate for the increase in mass, the bone thickness increases by more than a factor of 2 (precisely by a factor of 2.83)

(c) Radiocarbon dating can be used to date objects that are not older than about 57,000 years

(d)  $C(t) = C(0) e^{kt}$   $C(0)$  = initial amount of  $^{40}\text{K}$

half life:  $0.5 C(0) = C(0) e^{k \cdot 1.248 \cdot 10^9}$

$1.248 \cdot 10^9 k = \ln 0.5$

$k = \frac{\ln 0.5}{1.248 \cdot 10^9} \approx -0.555406 \cdot 10^{-9}$

so  $C(t) = C(0) e^{-0.555406 \cdot 10^{-9} \cdot t}$


$0.9645 C(0) = C(0) e^{-0.555406 \cdot 10^{-9} \cdot t}$

$t = \frac{\ln(0.9645)}{-0.555406 \cdot 10^{-9}} \approx 6.5079 \cdot 10^7$

$\approx 65,079,000$  years

9.  $\ln S(t) = \ln 24 + 1.8t \approx 1.8t + 3.18$

$\ln T(t) = \ln 24 - 0.8t \approx -0.8t + 3.18$



(negative  $t$  might, or might not make sense, depending on context)

10.  $\min = -1$

amplitude = 5

$\max = 9$

period = 2

average = 4

phase = 0

11.(a)  $y = \sin\left(4\left(t + \frac{\pi}{4}\right)\right)$  ... sine graph compressed by a factor of 4 then shifted  $\pi/4$  units to the left

min = -1  
 max = 1  
 average = 0  
 amplitude = 1






period =  $\frac{2\pi}{4} = \frac{\pi}{2}$   
 shift  $\frac{\pi}{4}$  to the left (or: phase =  $-\frac{\pi}{4}$ )

(b)  $y = \cos\left(\frac{t}{2}\right) + 5$  ... cosine graph stretch by a factor of 2 then up 5 units

min = 4  
 max = 6  
 average = 5  
 amplitude = 1


period =  $2\pi / \frac{1}{2} = 4\pi$   
 shift (phase) = 0

(c)  $y = -2\sin(3t) + 4$  ... sine graph, compressed horizontally by a factor of 3, expanded vertically by a factor of 2, reflected across X-axis, moved up 4 units



min = 2, max = 6, average = 4, amplitude = 2  
 phase = 0, period =  $2\pi/3$



12.



period is 12

$$\frac{2\pi}{a} = 12 \rightarrow a = \frac{2\pi}{12} = \frac{\pi}{6}$$

use cosine:  $\cos(at) \rightarrow \cos\left(\frac{\pi}{6}t\right)$



(average is  $\frac{2+4.6}{2} = 3.3$   
amplitude is 1.3)

$$\boxed{-1.3 \cos\left(\frac{\pi}{6}t\right) + 3.3}$$

