



Research

Cite this article: Molina C, Earn DJD. 2015 Game theory of pre-emptive vaccination before bioterrorism or accidental release of smallpox. *J. R. Soc. Interface* **12**: 20141387. <http://dx.doi.org/10.1098/rsif.2014.1387>

Received: 19 December 2014
Accepted: 8 April 2015

Subject Areas:
biomathematics

Keywords:
vaccination, smallpox, game theory, bioterrorism, epidemiology, cost-effectiveness

Author for correspondence:
Chai Molina
e-mail: cmgraff@gmail.com

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rsif.2014.1387> or via <http://rsif.royalsocietypublishing.org>.

Game theory of pre-emptive vaccination before bioterrorism or accidental release of smallpox

Chai Molina¹ and David J. D. Earn^{1,2}

¹Department of Mathematics and Statistics, and ²M.G. deGroot Institute for Infectious Disease Research, McMaster University, Hamilton, Ontario, Canada L8S 4K1

Smallpox was eradicated in the 1970s, but new outbreaks could be seeded by bioterrorism or accidental release. Substantial vaccine-induced morbidity and mortality make pre-emptive mass vaccination controversial, and if vaccination is voluntary, then there is a conflict between self- and group interests. This conflict can be framed as a tragedy of the commons, in which herd immunity plays the role of the commons, and free-riding (i.e. not vaccinating pre-emptively) is analogous to exploiting the commons. This game has been analysed previously for a particular post-outbreak vaccination scenario. We consider several post-outbreak vaccination scenarios and compare the expected increase in mortality that results from voluntary versus imposed vaccination. Below a threshold level of post-outbreak vaccination effort, expected mortality is independent of the level of response effort. A lag between an outbreak starting and a response being initiated increases the post-outbreak vaccination effort necessary to reduce mortality. For some post-outbreak vaccination scenarios, even modest response lags make it impractical to reduce mortality by increasing post-outbreak vaccination effort. In such situations, if decreasing the response lag is impossible, the only practical way to reduce mortality is to make the vaccine safer (greater post-outbreak vaccination effort leads only to fewer people vaccinating pre-emptively).

1. Introduction

The number of annual cases of smallpox in the early 1950s, just prior to the World Health Organization's global eradication programme, is estimated at 50 million [1]. The eradication campaign was successful [1], but samples of the variola virus are still kept in at least two known laboratories in Russia and the USA [2]. In a worrying incident in July 2014, previously forgotten vials containing samples of smallpox, some of which were viable, were found in a laboratory at the National Institutes of Health campus in Bethesda, MD, USA [3]. Thus, the threat of the reintroduction of smallpox, whether inadvertently or in a bioterrorist attack, is still present.

Consequently, some countries—notably the USA—are interested in measures to protect their populations from potential smallpox infection. Prophylactic vaccination for smallpox carries a high cost (relative to other vaccines in use today), as the probability of death following vaccination—or 'risk from being vaccinated'—is $r_v \simeq 10^{-6}$ and serious side-effects occur with probability $\sim 10^{-3}$ [1]. Of course, infection with smallpox carries a much greater risk, because the case fatality proportion—the 'risk from infection'—is $r_i \simeq 0.3$ [1]. (See table 1 for a summary of parameter estimates.)

The substantial vaccine-induced morbidity and mortality associated with smallpox vaccination make pre-emptive mass vaccination controversial. If vaccination is voluntary, then there is a conflict between self- and group interests. This conflict can be framed as a tragedy of the commons, in which herd immunity plays the role of the commons, and free-riding (i.e. not vaccinating pre-emptively) is analogous to exploiting the commons. A previous game-theoretical study by Bauch *et al.* [4] examined this conflict of interest, and focused on the trade-off between prophylactic vaccination and post-outbreak mass vaccination (which has been shown to outperform contact-traced vaccination in a bioterrorism setting

Table 1. Summary of the fundamental (i.e. not derived) numerical parameters in our analysis, together with estimated values. Note that in [4] the probability of an outbreak was denoted r rather than a . Here, we use r for the relative risk, as in [5]. The proportion of the population infected initially by a bioterrorist attack or accidental release, α , corresponds to infection of 5000 individuals in a population of 290 million (after [4]).

quantity	interpretation	value	source
r_v	mortality risk from vaccination (probability)	10^{-6}	[4]
r_i	mortality risk from infection (probability)	0.3	[4]
\mathcal{R}_0	basic reproductive ratio	5	[6–8]
t_{ser}	mean serial interval	22 days	[9, p. 141]
$1/\sigma$	mean latent period (SEIRV)	15 days	[1, p. 188] and [4]
$\phi_{(\text{model})}$	vaccination effort parameter (exact interpretation depends on model)	see table 4	
t_{lag}	response lag before initiation of post-outbreak vaccination	0 days, except in §7.5	
a	probability of attack or accidental release per lifetime	0.01	[4]
α	proportion of susceptibles initially infected in an outbreak	$5000/290 \times 10^6 \simeq 1.72 \times 10^{-5}$	[4]

Table 2. Summary of derived parameters.

quantity	interpretation	value
$r = r_v/r_i$	relative risk (from being vaccinated compared with natural infection)	$10^{-6}/0.3 \simeq 3.33 \times 10^{-6}$
$1/\gamma$	mean time infectious (SIRV)	$t_{\text{ser}} = 22$ days
$1/\gamma$	mean time infectious (SEIRV)	$t_{\text{ser}} - (1/\sigma) = 7$ days
β	transmission rate	$\gamma\mathcal{R}_0$
π_p	probability that an un-vaccinated individual will eventually be infected if the vaccine coverage level in the population is p	derived from epidemic model in §5
ψ_p	probability of an individual un-vaccinated at the beginning of the epidemic becoming vaccinated, given vaccine coverage level p	derived from epidemic model in §5

[10]). In particular, they showed that if the decision regarding pre-emptive vaccination is left to the individual, then the vaccine coverage achieved will be suboptimal from the group perspective. Bauch *et al.* [4] assumed that, once a post-outbreak vaccination campaign begins, individuals will be vaccinated at a constant rate determined by existing infrastructure.

Various mechanisms might drive the rate of vaccination. Vaccination at a constant rate might be achieved if vaccination centres are flooded by individuals seeking the vaccine, and are operating at peak capacity. However, public responsiveness to such a campaign is hard to predict. If demand for the vaccine does not exceed the maximal rate of distribution by public health services, the post-outbreak dynamics might play out differently, depending on the public's reaction patterns. For example, media reports on the number of new cases might influence individuals to obtain the vaccine; in that case, it is reasonable to model the vaccination rate as proportional to smallpox incidence.

In this paper, we return to the problem posed by Bauch *et al.* [4], but compare a variety of possible post-outbreak vaccination scenarios (described intuitively in §2 and in precise mathematical terms in §5). Whereas the scenario considered in [4] could only be analysed numerically, several of the vaccination scenarios that we consider here can be addressed analytically to obtain exact results. To this end, in §3, we make some adjustments to the game-theoretical framework of Bauch & Earn [5], so that it can be applied to the scenarios we investigate here.

Throughout this paper, we use smallpox as an illustrative example. However, our analyses can be applied to any

vaccine-preventable infectious disease that could be used for bioterrorism or released accidentally, and for which the susceptible–infectious–removed (SIR) or susceptible–exposed–infectious–removed (SEIR) models are applicable (see §5). Our qualitative results appear to be robust to which post-outbreak vaccination scenario is considered and the specific parameter values (we prove this in some cases), but the precise numerical values will vary.

We calculate the vaccination coverage obtained by voluntary pre-emptive vaccination and assess the costs of this policy when compared with mandatory vaccination. The group-optimal pre-emptive vaccine coverage is discussed in §4. We discuss parameter estimates and the procedure used to compare the various models fairly in §6. We compare the predictions of the various models, and emphasize important considerations for public health in §7. Notation and definitions are summarized in tables 1–3.

2. Vaccination scenarios

In this section, we give a brief description of the various post-outbreak vaccination scenarios considered in this paper.

Media coverage of a smallpox outbreak is likely to influence individual decisions concerning vaccination. Measures of severity of the outbreak that are likely to appear in the media include:

- death rates, as in ‘300 people died of smallpox today’,
- total number of people currently infected (prevalence), as in ‘there are now 30 000 people sick with smallpox’, and

Table 3. Summary of other notation.

quantity	interpretation
P	probability that an individual chooses to vaccinate pre-emptively (this defines the individual's strategy)
p	pre-outbreak vaccine coverage (proportion of the population vaccinated pre-emptively)
p_g	the group optimum, i.e. the proportion of the population vaccinated pre-emptively which minimizes mortality
p_i	the individual equilibrium, i.e. the level of pre-outbreak vaccine coverage which is the unique Nash equilibrium, as described in §3
$C(p)$	the mortality cost, i.e. the proportion of the population that is expected to die, given pre-emptive vaccine coverage p
t_{lag}^*	the critical lag, i.e. the response lag beyond which mortality is independent of vaccination effort (see §7.5.1)
\tilde{t}_{lag}^*	the effective critical lag, i.e. the response lag beyond which mortality is identical for all feasible values of vaccination effort (see §7.5.2)

Table 4. Summary of notable levels of the vaccination effort parameter, $\phi_{(model)}$, for the different models. The first column contains 'fair comparison' values for the vaccination effort parameters of the various models, as calculated in §6.2. In our simulations, we allowed $\phi_{(model)}$ to range between 0 and values generally above the 'fair comparison' values (except for ϕ_{inst} , for which we used the entire possible range of [0, 1]). The second column contains the minimal values of the vaccination effort parameter ($\phi_{(model)}$) for which the individual equilibrium is to delay (that is, $\phi_{(model)}$ at the end of the mortality plateau; see §7.2.1).

model	'fair comparison' value	value at end of mortality plateau
ϕ_{prev}	1582 per day	571 per day
ϕ_{susc}	0.1 per day	0.08 per day
ϕ_{inc}	5190 per day	1137 per day
ϕ_{inst}	—	0.82 per day
ϕ_{const}	0.1 per day	0.015 per day

— new cases (incidence), as in '200 new cases of smallpox were confirmed today'.

We consider separately how each of these types of information could affect smallpox vaccine uptake; in each case, we assume that the vaccination rate is proportional to the relevant quantity (e.g. prevalence). Note that, in standard epidemiological models [6], death rate is proportional to prevalence, so the first and second cases above are mathematically identical.

As a type of 'null model' for media-induced vaccination, we also consider the situation in which

— Vaccination rate is simply proportional to the size of the remaining susceptible population; this corresponds to a constant *per capita* vaccination rate for susceptible individuals (see the electronic supplementary material, appendix B.1). This can be regarded as a 'null model' to compare with models for the scenarios above in the following sense: individuals' proclivity to vaccinate is constant over time, and does not depend on the state of the epidemic (i.e. on prevalence or incidence, which are likely to be reported by the media), whereas the vaccination rate falls as the number of susceptibles decreases over time, meaning that fewer individuals per unit time are inclined to vaccinate.

We also consider two scenarios in which vaccine uptake is not influenced by the media, but is constrained by the capabilities of public health authorities:

- If an outbreak occurs, immediately vaccinate a proportion of the susceptible population. The proportion might describe the efficacy of a post-outbreak campaign in convincing those who have thus far avoided vaccination. Individuals who remain un-vaccinated after this post-outbreak campaign would be persons holding particularly radical anti-vaccine opinions.
- Susceptible individuals are vaccinated at a constant rate until there are no more susceptibles remaining.

Finally, for each of the above scenarios, we investigate the effect of a lag between the start of an outbreak and the initiation of the post-outbreak vaccination response (allowing for public health authorities to organize a response to the outbreak). Bauch *et al.* [4] assumed such a response lag in their model, which is otherwise identical to the final scenario described above.

The epidemic models associated with each of the above five scenarios are described in detail in §5.

3. Game-theoretical formulation

In this section, we adapt the game-theoretical framework of Bauch & Earn [5] to our current problem. We assume that all individuals have full knowledge and are rational (in the game-theoretical sense; see [11]).

We denote the proportion of the population vaccinated pre-emptively as p . Because a proportion r_v of those vaccinated will die, the pre-outbreak vaccine coverage (the proportion of the population that is immune prior to the outbreak) is $p_{eff} = p(1 - r_v)/(1 - pr_v)$ [4], which is slightly smaller than p . But, because none of the mathematical analysis and conclusions which follow are affected by this, and because the difference between p and p_{eff} is negligible, we refer to p as the pre-outbreak vaccine coverage level for simplicity (as in [4]).

Let $a \in [0, 1]$ be the probability of an outbreak (' a ' for 'bioterrorist attack probability' or 'accidental release probability') per lifetime (or whatever time period is under consideration). Consider two pure strategies: *vaccinate* and *delay*. The former vaccinates pre-emptively, before the beginning of an outbreak, and so receives (expected) pay-off $-r_v$; the latter delays vaccination until after an outbreak (at which point she/he may still be vaccinated during the public health post-outbreak vaccination campaign), and receives pay-off

$$-a[r_i\pi_p + \psi_p r_v], \quad (3.1)$$

where π_p and ψ_p are the probabilities of a delayer being infected, or vaccinated, respectively, after an outbreak (the delayer

infection and vaccination probabilities are discussed in more detail in §5). A *mixed strategy* is specified by the probability P that an individual will choose to vaccinate pre-emptively. We also assume $r_v < ar_i$ because if it were not so, even if all delayers were infected in an outbreak, the risk of dying in an outbreak would be smaller than the risk of dying from the side-effects of the vaccine, hence there would be no reason to vaccinate.

The pay-off to an individual playing a mixed strategy (vaccinating with probability P) in a population in which the coverage level is p is given by

$$E(P, p) = -Pr_v - (1 - P)a(\pi_p r_i + \psi_p r_v). \quad (3.2)$$

Equivalently, defining the *relative risk* of vaccination compared with infection as

$$r = \frac{r_v}{r_i}, \quad (3.3)$$

we have $E(P, p) = -r_i[rP + (1 - P)a(\pi_p + \psi_p r)]$. Because the parameter r_i simply scales the game pay-off by a constant, it does not change the dynamics. We therefore use the rescaled pay-off function

$$E(P, p) = -[rP + (1 - P)a(\pi_p + \psi_p r)]. \quad (3.4)$$

Suppose that a proportion ϵ of the population vaccinate with probability P and $1 - \epsilon$ vaccinate with probability Q . Following Bauch & Earn [5], we assume 100% vaccine efficacy, which implies coverage level $p = \epsilon P + (1 - \epsilon)Q$. (Note that in a homogeneous population where all individuals play the same strategy P , i.e. $\epsilon = 1$, the coverage is $p = P$.) The pay-offs to individuals playing P and Q in such a population are then

$$E_P(P, Q, \epsilon) = E(P, \epsilon P + (1 - \epsilon)Q) \quad (3.5a)$$

or

$$E_Q(P, Q, \epsilon) = E(Q, \epsilon P + (1 - \epsilon)Q), \quad (3.5b)$$

respectively, and the *pay-off gain* to an individual playing P rather than that to Q in this population is

$$\begin{aligned} \Delta E &= E_P(P, Q, \epsilon) - E_Q(P, Q, \epsilon) \\ &= -[rP + (1 - P)a(\pi_p + \psi_p r)] + [rQ + (1 - Q)a(\pi_p + \psi_p r)] \\ &= (\pi_p + r\psi_p - \frac{r}{a})a(P - Q), \quad \text{where } p = \epsilon P + (1 - \epsilon)Q. \end{aligned} \quad (3.6)$$

A strategy P^* is a *Nash equilibrium* (NE) if and only if (iff), in a population in which all individuals are playing P^* , no player employing a different strategy can achieve a higher pay-off. Mathematically, this means that for any other strategy $Q \in [0, 1]$ if the proportion playing Q is small enough (i.e. $1 - \epsilon$ is sufficiently small), then the pay-off gain to strategy P^* is non-negative, i.e. $\Delta E(P^*, Q, \epsilon) \geq 0$. When such an NE exists, we refer to this strategy as the *individual equilibrium* and denote it by p_i . This equilibrium is ‘individual’ in the sense that it is determined by individuals attempting to maximize their pay-offs (unlike the group optimum discussed in §4). Note, however, that this is a *population game* [5,12], so the pay-off to individuals depends on the frequencies of strategies in the entire population.

Additionally, consider a scenario whereby strategy P invades a population playing strategy Q . If, in this scenario, strategies P that are closer to the NE P^* than the prevalent strategy Q obtain a higher pay-off than the prevalent strategy, then P^* is called a *convergently stable Nash equilibrium* (CSNE). Mathematically, this is equivalent to demanding that

if $\epsilon \ll 1$, then

$$P^* < P < Q \leq 1 \Rightarrow \Delta E(P, Q, \epsilon) \geq 0$$

and

$$0 \leq Q < P < P^* \Rightarrow \Delta E(P, Q, \epsilon) \geq 0.$$

In order to proceed with the analysis, it is necessary to derive the probabilities π_p and ψ_p from an epidemiological model, either numerically or analytically (see the electronic supplementary material, appendix E). Proofs of existence and uniqueness of a CSNE are given for several cases in the electronic supplementary material, appendix G. These proofs depend on π_p being a decreasing function of p . We have shown this to be true when post-outbreak vaccination is instantaneous or proportional to incidence, and also when vaccination is proportional to prevalence and $\alpha\phi_{\text{prev}} > \gamma(1 - \alpha)$. Based on biological intuition corroborated with simulations, we have assumed that π_p decreases with p for all the models considered here. This has also recently been proved for other post-outbreak vaccination models not considered here (F Bai, F Brauer 2015, personal communication).

4. Group optimum

From the perspective of a public health official (i.e. group interest), it is desirable to attain the vaccine coverage that minimizes mortality. From this group perspective, a strategy is specified by the proportion p of the population that is pre-emptively vaccinated. The currency with which we compare strategies is the *mortality cost* $C(p)$, i.e. the proportion of the population that is expected to die (either from smallpox infection or from vaccination),

$$C(p) = rp + (1 - p)a(\pi_p + \psi_p r), \quad p \in [0, 1], \quad (4.1)$$

where we have ignored a factor of r_i as in equation (3.4). The minimum mortality cost yields the *group optimum* coverage level, which we denote p_g . The minimum of $C(p)$ on $[0, 1]$ may be attained either at a local minimum in $(0, 1)$ or at one of the endpoints

$$C(0) = a(\pi_0 + \psi_0 r) \quad (4.2a)$$

and

$$C(1) = r. \quad (4.2b)$$

To completely specify the cost $C(p)$, we need the probabilities π_p and ψ_p , derived from the epidemiological model (see §5 and the electronic supplementary material, appendix E), just as for the individual equilibrium. We have found an exact analytical expression for p_g in one subcase (see the electronic supplementary material, appendix H) and calculated it numerically in the other cases.

5. Epidemiological models

In order to find the group optimum (p_g) and individual equilibrium (p_i), two key quantities are calculated from the epidemic models: the *delayer infection probability* π_p (the probability of a delayer being infected after an outbreak) and the *delayer vaccination probability* ψ_p (the probability that a delayer is eventually vaccinated, given an outbreak).

Both π_p and ψ_p depend on the disease dynamics and the post-outbreak vaccination scenario. In the following, we

assume that, in the absence of post-outbreak vaccination, the SIR model is adequate to represent the disease dynamics [13, §4]. The models do not include vital dynamics (births and deaths from all natural causes other than the disease), because the mean serial interval (also called the disease generation time, $t_{\text{ser}} = 22$ days [9, p. 141]) is much smaller than the mean lifetime (approx. 80 years in the USA [14]). Note that, for diseases for which the outbreak time scale is similar to the mean lifetime, vital dynamics can easily be included in the analysis (e.g. as in [5], where much longer term dynamics were considered).

Let $S(t)$, $I(t)$, $R(t)$ and $V(t)$ be the proportions of susceptible, infected, removed (recovered or dead from smallpox infection) and vaccinated individuals (immune or dead from vaccination), respectively, at time t . Our basic framework is the SIRV model, described by the differential equations

$$\dot{S} = -\beta SI - \dot{V}, \quad (5.1a)$$

$$\dot{I} = \beta SI - \gamma I, \quad (5.1b)$$

$$\dot{R} = \gamma I \quad (5.1c)$$

and
$$\dot{V} \geq 0, \quad (5.1d)$$

where \dot{V} must be non-negative as indicated and is defined differently for each of the distinct scenarios of post-outbreak vaccination described in §2.

We assume that no one has natural immunity or retains immunity from vaccination decades earlier. This is an approximation, because many living individuals were vaccinated before smallpox was declared eradicated in 1979 [1], and many of those vaccinated individuals are probably still immune (vaccine-derived immunity seems to wane quite slowly and lifelong immunity is common [15]). However, smallpox is considered to have been eliminated in the USA as early as 1950, and while routine vaccination continued in some states well after that [1], the proportion of US residents younger than 60 who have been vaccinated is likely to be very small.

Thus, we assume that the coverage level prior to an outbreak is p , the proportion pre-emptively vaccinated. Consequently, prior to the outbreak, a proportion $1 - p$ of the population is susceptible. When a bioterrorist attack or accidental release takes place (at time $t = 0$), an *initial attack proportion* α of the susceptible population is infected. Thus,

$$S(0) = (1 - p)(1 - \alpha), \quad (5.2a)$$

$$I(0) = (1 - p)\alpha, \quad (5.2b)$$

$$R(0) = 0 \quad (5.2c)$$

and
$$V(0) = p. \quad (5.2d)$$

After an outbreak, the epidemic is over when no one remains infective ($I = 0$). In the electronic supplementary material, appendix C, we show rigorously that this is guaranteed to occur, either in finite time or in the limit as $t \rightarrow \infty$. In either case, we use the subscript ∞ to refer to the time at which the epidemic ends. Thus, S_∞ , I_∞ , R_∞ and V_∞ refer to the proportions of the population in the susceptible, infective, removed and vaccinated compartments at the end of the epidemic. With this notation, the probabilities of infection and vaccination for delayers are, respectively,

$$\pi_p = \frac{R_\infty - R(0)}{S_0 + I_0} = \frac{R_\infty}{1 - p} \quad (5.3a)$$

$$\begin{aligned} \text{and } \psi_p &= \frac{V_\infty - V_0}{S_0 + I_0} = \frac{V_\infty - p}{1 - p} = 1 - \frac{1 - V_\infty}{1 - p} \\ &= 1 - \frac{S_\infty + I_\infty + R_\infty}{1 - p} = 1 - \pi_p - \frac{S_\infty}{1 - p}. \end{aligned} \quad (5.3b)$$

We emphasize that R is the proportion of the population that has been infected (and consequently is either immune or has died); hence, $R(0) = 0$, because anyone who is immune at time $t = 0$ is immune from vaccination. Intuitively, there is no endemic equilibrium in these models, because the combination of vaccination and natural spread of disease must eventually cause susceptibles to be so rare that the disease cannot spread (recall that these models neglect vital dynamics).

Lastly, note that π_p is undefined at $p = 1$ (i.e. if everyone pre-emptively vaccinates), as there are no delayers for whom to calculate the probability of being infected. We define π_1 as the limit of the delayer infection probability,

$$\pi_1 = \lim_{p \rightarrow 1^-} \pi_p, \quad (5.4)$$

i.e. π_1 is the limit of π_p as pre-emptive vaccination approaches full coverage. In the electronic supplementary material, appendix D, we show that this limit is equal to the proportion of susceptibles initially infected in an outbreak, i.e. $\pi_1 = \alpha$ for all models considered.

In the following, we describe (and interpret mechanistically) various models that we compare, and present some analytical results. In all models, the vaccination rate depends on a vaccination effort parameter, $\phi_{(\text{model})}$, the exact interpretation of which is model dependent.

5.1. Vaccination rate \propto disease prevalence

In this model, vaccination occurs at a rate proportional to disease prevalence (I). A plausible scenario to which such a model would apply is if people respond to media reports on disease prevalence. As a result of increasing disease prevalence, the public might perceive the risk of being infected as higher, and be moved to vaccinate. Consequently,

$$\dot{V} = \phi_{\text{prev}} \text{sgn}(S)I, \quad (5.5)$$

where

$$\text{sgn}(x) = \begin{cases} -1 & \text{if } x < 0, \\ 0 & \text{if } x = 0, \\ 1 & \text{if } x > 0. \end{cases}$$

This model could also represent the case where vaccination rate is proportional to death rate, i.e. people vaccinate in response to media reports on new disease-induced deaths. Because the death rate is proportional to the rate at which the removed compartment, R , grows, which is proportional to I , the vaccination rate would also be proportional to I .

In the electronic supplementary material, appendix E.1.1, we find the final size relations [16–18] for the model defined by equation (5.5). These are given by

$$S_\infty = \begin{cases} 0 & \text{if } p < p_0 \text{ or } 1 \leq p_m, \\ S_\infty^1 & \text{if } p_0 \leq p \leq 1, \end{cases} \quad (5.6)$$

$$R_\infty = \begin{cases} 1 - p - \frac{\phi_{\text{prev}}}{\beta} \ln \left(\frac{\beta}{\phi_{\text{prev}}} S(0) + 1 \right) & \text{if } p < p_0 \text{ or } 1 \leq p_m, \\ \frac{\gamma}{\gamma + \phi_{\text{prev}}} (1 - p - S_\infty^1) & \text{if } p_0 \leq p \leq 1 \end{cases} \quad (5.7)$$

and

$$V_\infty = \begin{cases} p + \frac{\phi_{\text{prev}}}{\beta} \ln\left(\frac{\beta}{\phi_{\text{prev}}} S(0) + 1\right) & \text{if } p < p_0 \text{ or } 1 \leq p_m, \\ \frac{1}{\gamma + \phi_{\text{prev}}} (\phi_{\text{prev}}(1 - S_\infty^1) + \gamma p) & \text{if } p_0 \leq p \leq 1, \end{cases} \quad (5.8)$$

with

$$S_\infty^1 = -\frac{1}{\beta} \left(\phi_{\text{prev}} + (\gamma + \phi_{\text{prev}}) \times W_0 \left(-\frac{\beta S(0) + \phi_{\text{prev}}}{\gamma + \phi_{\text{prev}}} e^{-\frac{\beta(1-p) + \phi_{\text{prev}}}{\gamma + \phi_{\text{prev}}}} \right) \right), \quad (5.9)$$

$$p_m = 1 + \frac{\alpha \phi_{\text{prev}} - \gamma(1 - \alpha)}{\beta(1 - \alpha)} \quad (5.10)$$

and

$$p_0 = 1 + \frac{\phi_{\text{prev}}}{\beta(1 - \alpha)} + \frac{\gamma + \phi_{\text{prev}}}{\beta} \times W_k \left(-\frac{\phi_{\text{prev}}}{(1 - \alpha)(\gamma + \phi_{\text{prev}})} e^{-\frac{\phi_{\text{prev}}}{(1 - \alpha)(\gamma + \phi_{\text{prev}})}} \right), \quad (5.11)$$

where $k = 0$ if $p_m < 1$ and $k = -1$ if $p_m \geq 1$. W_0 is the principal branch of Lambert's W function [19,20], and W_{-1} is its other real branch (see the electronic supplementary material, appendix A). p_m is the unique maximum of the function S_∞^1 . S_∞^1 has two roots, one at $p = 1$, and the other at p_0 , which need not lie in the interval $[0, 1]$ (p_0 is a formal root and need not correspond to a meaningful probability). Note that if $p_m > 1$ then $p_0 > p_m$, and if $p_m < 1$ then $p_0 < p_m$.

If $p_m > 1$, then no delayers will remain susceptible at the end of the epidemic (i.e. all delayers will be either vaccinated or infected), regardless of the initial vaccine coverage level p . Moreover, if $p_m < 1$, but $p_0 < 0$, there are always some delayers who remain susceptible at the end of the epidemic, regardless of the initial coverage level p . If $0 < p_0 < p_m < 1$, then for $p \in [0, p_0]$ there will be no susceptibles left at the end of the epidemic, and for $p \in (p_0, 1)$ there will be some remaining susceptibles. Thus, there is a wide range of parameter values for which some susceptibles remain at the end of the epidemic; in such cases, $\pi_p + \psi_p < 1$. Numerical evidence and biological intuition suggest that π_p is a decreasing function of p , and we assume that this is the case from here on (this is proven for $p_m \geq 1$ in the electronic supplementary material, appendix E.1.3).

Finally, because the mean infectious period (7 days; see table 2 and §6.1) is longer than the time required to complete the vaccination programme (possibly as short as 3 days [21]), it is interesting to take the limit $\gamma \rightarrow 0$ (corresponding to an infinite infectious period) while keeping $\mathcal{R}_0 = \beta/\gamma$ fixed. In this limit, $p_m \rightarrow \infty$ so $S_\infty = 0$ (equation (5.6)), which is in accordance with the assumption—made in [4]—that individuals are ultimately either removed or vaccinated.

We show in the electronic supplementary material, appendix G.1, that there is always a unique CSNE, that is, a 'best strategy' from the individual perspective. Moreover, an analytical expression for this individual equilibrium can be found if

$$\text{either } \phi_{\text{prev}} \geq \gamma(1 - \alpha)/\alpha \quad (5.12a)$$

$$\text{or } 0 \leq p_0 < p_m \leq 1, \quad \pi_0 > \rho_1 > \pi_{p_0} \text{ and } \pi_{p_0} < \rho_2. \quad (5.12b)$$

In addition, we find an analytical formula for the group optimum when $\phi_{\text{prev}} \geq \gamma(1 - \alpha)/\alpha$ (see the electronic supplementary material, appendix H, for details).

5.2. Vaccination rate \propto incidence

A vaccination rate proportional to incidence again reflects media-induced vaccination. However, in this model, the public reacts to reports of new cases, rather than reports of the total number of sick individuals. Thus,

$$\dot{V} = \phi_{\text{inc}} SI. \quad (5.13)$$

In the electronic supplementary material, appendix E.2, we show that

$$S_\infty = -\frac{\gamma}{\beta} W_0 \left(-\frac{\beta(1-p)(1-\alpha)}{\gamma} e^{-\frac{\beta + \phi_{\text{inc}} \alpha}{\gamma}(1-p)} \right), \quad (5.14)$$

$$V_\infty = p + \frac{\phi_{\text{inc}}}{\beta + \phi_{\text{inc}}} ((1-p)(1-\alpha) - S_\infty) \quad (5.15)$$

and $R_\infty = 1 - p - \frac{\phi_{\text{inc}}(1-p)(1-\alpha) + \beta S_\infty}{\beta + \phi_{\text{inc}}}$. (5.16)

Again, because there are susceptible individuals left at the end of the epidemic, $\pi_p \neq 1 - \psi_p$. We show that $\partial_p \pi_p < 0$ (in the electronic supplementary material, appendix E.2) and find that there is a unique CSNE, p_i , for which an exact formula is derived in the electronic supplementary material, appendix G.2.

5.3. Vaccination rate \propto proportion still susceptible

In this scenario, susceptible individuals vaccinate at a rate

$$\dot{V} = \phi_{\text{susc}} S. \quad (5.17)$$

This is a null model, in the sense that susceptible individuals have a constant probability per unit time of being vaccinated, ϕ_{susc} , independent of the state of the outbreak, as shown in the electronic supplementary material, appendix B.1.

We were able to obtain analytical final size relations for this model (see the electronic supplementary material, appendix E.3), but we found the formulae too cumbersome to be useful. Thus, the remainder of our analysis of this model was performed by integrating the differential equations numerically. In our numerical simulations, we always find that π_p decreases with p (in the electronic supplementary material, appendix G.3, our proof of the existence of a CSNE depends on this being true).

5.4. Instantaneous vaccination of a proportion ϕ_{inst} of the population

Some experts believe that the entire USA could be vaccinated in 3 days [21], which is less than the latent period of smallpox. Consequently, instantaneous vaccination of a proportion ϕ_{inst} of the population remaining susceptible after the outbreak is also a realistic scenario to model. In this case, once vaccination has occurred, the disease simply spreads according to the standard SIR model,

$$\dot{S} = -\beta SI, \quad (5.18a)$$

$$\dot{I} = (\beta S - \gamma) I \quad (5.18b)$$

and $\dot{R} = \gamma I$, (5.18c)

with initial conditions given by

$$S(0) = (1-p)(1-\alpha)(1-\phi_{\text{inst}}),$$

$$I(0) = (1-p)\alpha,$$

$$R(0) = 0,$$

$$V(0) = p + \phi_{\text{inst}}(1-p)(1-\alpha).$$

Note that in this scenario we deviate from the convention we use for all the other models, in which $S(0)$ is the initial density of susceptibles prior to the beginning of the post-outbreak vaccination response. Here, $S(0)$ is the density of susceptibles *after* the post-outbreak vaccination response has taken place.

For this scenario, we find (in the electronic supplementary material, appendix E.4)

$$S_\infty = -\frac{\gamma}{\beta} W_0 \left(-\frac{\beta}{\gamma} S(0) e^{-\frac{\beta}{\gamma}(1-V(0))} \right) \quad (5.19)$$

and

$$R_\infty = \frac{\gamma}{\beta} \ln \frac{S(0)}{S_\infty}. \quad (5.20)$$

We also show that π_p is a decreasing function of p , ψ_p is constant and $\pi_p + \psi_p \neq 1$ (see the electronic supplementary material, appendix E.4). In addition, we have proved that, for this model, there is always a unique CSNE, for which we derive an exact formula in the electronic supplementary material, appendix G.4.

5.5. Constant rate vaccination

This is the model of Bauch *et al.* [4], in which vaccination occurs at a constant rate ϕ_{const} . Note that in [4] vaccination begins after a response lag t_{lag} , which is the public health services' response time. This lag is taken to be $t_{\text{lag}} = 0$ except in §7.5.

For consistency with [4], we included an exposed (but not infective) stage (E), in this model, making it an SEIRV model. This contrasts all the other scenarios, which we have modelled using a simpler SIRV formulation. Our choice of the SIRV framework for the new scenarios is motivated by mathematical tractability and by work subsequent to [4], indicating that SEIR dynamics are captured by an appropriately parametrized SIR model (see §6.1, but see §7.5.1 for an exception).

The model equations for the constant rate vaccination scenario are

$$\dot{S} = -\beta SI - \dot{V}, \quad (5.21a)$$

$$\dot{E} = \beta SI - \sigma E, \quad (5.21b)$$

$$\dot{I} = \sigma E - \gamma I, \quad (5.21c)$$

$$\dot{R} = \gamma I \quad (5.21d)$$

$$\text{and} \quad \dot{V} = \begin{cases} \phi_{\text{const}} & \text{if } t_{\text{lag}} < t \text{ and } S > 0, \\ 0 & \text{if } t \leq t_{\text{lag}} \text{ or } S \leq 0. \end{cases} \quad (5.21e)$$

We have not found a final size relation for this model.

Under the biologically plausible assumption that π_p decreases with p (verified by simulation), Bauch *et al.* [4] have shown the existence of a unique CSNE for this model.

6. Parameter estimates, fair comparisons of models and numerical procedures

Because one of the models we investigate includes an exposed class, and the vaccination effort parameter $\phi_{\text{(model)}}$ has a different meaning in each scenario we examine, fair comparisons of model results are not completely straightforward. In this section, we consider how the various models can be compared.

6.1. SIR versus SEIR

It is well known that similar dynamics are obtained with the standard SIR and SEIR models with identical basic reproductive number, \mathcal{R}_0 , if the mean infectious period in the SIR

model is set equal to the sum of the mean latent and infectious periods in the SEIR model [6, p. 668]. More generally, models can be fairly compared if they have the same mean serial interval [13, §4].

Estimates of the basic reproductive ratio \mathcal{R}_0 of smallpox vary in the range $3 \leq \mathcal{R}_0 \leq 10$ [6–8]. Following [4], we take $\mathcal{R}_0 = 5$. We take the mean serial interval to be $t_{\text{ser}} = 22$ days, as in [9, p. 141] (but note that [4] used $t_{\text{ser}} = 14$ days, and [22] estimated $t_{\text{ser}} = 17.7$ days).

In the constant rate vaccination model, we take the mean latent period to be $1/\sigma = 15$ days [9, p. 141] (based on summing the incubation and prodrom periods, which typically last 12 and 3 days, respectively; see [1, p. 188]). In an SEIR model, the mean serial interval is the sum of the mean latent and infectious periods [13,23]; hence, $1/\gamma = 22 - 15 = 7$ days and $\beta = \gamma \mathcal{R}_0 = 5/7$ per day. In the SIRV models, we take $1/\gamma = t_{\text{ser}}$, whereas β is modified so that $\mathcal{R}_0 = 5$ (i.e. $\beta = \gamma \mathcal{R}_0 = 5/22$ per day).

6.2. Vaccination effort parameter $\phi_{\text{(model)}}$

Public health policy changes will affect the vaccination effort parameter $\phi_{\text{(model)}}$, where (model) refers to any of 'prev', 'inc', 'susc', 'inst' or 'const'. In order to compare the outcomes of the various vaccination scenarios, for each vaccination model, we find the *fair comparison value* of $\phi_{\text{(model)}}$, that is, the value of $\phi_{\text{(model)}}$ that yields a maximal vaccination rate that is equal to the fixed rate in the constant rate vaccination model of Bauch *et al.* [4], $\dot{V} = 0.1$ per day (see description under ϕ_{const} below). This allows us to identify, for each scenario, ranges of $\phi_{\text{(model)}}$ that can feasibly be attained in reality (i.e. $\phi_{\text{(model)}}$ between 0 and the fair comparison value). Our aim is then to compare the different vaccination strategies in terms of vaccine doses used and total expected mortality (we will be interested in the values of these observables at both the individual equilibrium and the group optimum). The fair comparison values are summarized in table 4.

ϕ_{prev} . In the prevalence model, $\dot{V} = \phi_{\text{prev}} I$, the vaccination rate is proportional to the prevalence, I , and the vaccination effort parameter ϕ_{prev} is the rate of vaccination *per infected individual*. In the electronic supplementary material, appendix F.1, we calculate the maximal vaccination rate as a function of the model parameters, α , β , γ and ϕ_{prev} and p . We find that the maximal vaccination rate for a given initial coverage, p , decreases with the vaccination effort, ϕ_{prev} . We also find that, when α , β and γ are as in tables 1 and 2, a maximal vaccination rate of 0.1 per day is obtained when $\phi_{\text{prev}} \approx 1582$ per day.

ϕ_{inc} . In the incidence model, $\dot{V} = \phi_{\text{inc}} SI$, the vaccination effort parameter ϕ_{inc} is the vaccination rate *per infective per susceptible*. In the electronic supplementary material, appendix F.2, we calculate the maximal vaccination rate, as a function of the model parameters, α , β , γ and ϕ_{inc} . We show that the maximal vaccination rate, $\max\{\dot{V} : t \geq 0, p \in [0, 1]\}$, is an increasing function of ϕ_{inc} and that in order to obtain a maximal vaccination rate of 0.1 per day or lower, with α , β and γ as in tables 1 and 2, one needs $\phi_{\text{inc}} \approx 5190$ per day.

ϕ_{susc} . With $\dot{V} = \phi_{\text{susc}} S$, the vaccination effort parameter ϕ_{susc} is the vaccination rate *per susceptible individual* (alternatively, ϕ_{susc} can be interpreted as the probability per unit time of a delayer being vaccinated; see the electronic supplementary material, appendix B.1). In this model, the vaccination

rate \dot{V} is always decreasing, because S can only decrease, so $\max\{\dot{V}\} = \phi_{\text{susc}}S(0) = \phi_{\text{susc}}(1 - \alpha)(1 - p)$ (cf. equation (5.2*a*)). Because the maximal vaccination rate decreases with increasing initial coverage, p , the maximal vaccination rate is attained with no pre-emptive vaccination ($p = 0$). Because $S(0) = 1 - \alpha$, the maximal vaccination rate is $\max\{\dot{V}\} = (1 - \alpha)\phi_{\text{susc}}$, and a vaccination rate of 0.1 per day is attained for $\phi_{\text{susc}} = 0.1 / (1 - \alpha) \approx 0.1$ per day (because $\alpha \ll 1$).

ϕ_{inst} . For instantaneous vaccination, the vaccination effort parameter ϕ_{inst} is the proportion of susceptibles instantaneously vaccinated when an outbreak occurs. Thus, $\phi_{\text{inst}} \in [0, 1]$. The vaccination rate is either 0 (if $\phi_{\text{inst}} = 0$) or effectively infinite (if $0 < \phi_{\text{inst}} \leq 1$, because vaccination occurs all at once). We thus consider the entire range $0 \leq \phi_{\text{inst}} \leq 1$, because there is no value of ϕ_{inst} that results in a vaccination rate of 0.1 per day.

ϕ_{const} . With $\dot{V} = \phi_{\text{const}}$, the vaccination rate is constant, so ϕ_{const} is simply the proportion of the total population that can be vaccinated per unit time. Bauch *et al.* [4] estimated ϕ_{const} for New York City to be

$$\phi_{\text{const}} = (5000 \text{ vaccinators}) \times \left(\frac{200 \text{ people per day}}{\text{vaccinator}} \right) \times \frac{1}{10^7 \text{ people}} = \frac{0.1}{\text{day}}. \quad (6.1)$$

A rate of $\phi_{\text{const}} = 0.1$ per day means the entire population can be vaccinated in 10 days.

6.3. Numerical procedures

When generating figures 1–4, calculations of the following quantities were necessary: the mortality cost, $C(p)$ (equation (4.1)), the group optimum, p_g (§4), and the individual equilibrium, p_i (§3).

To find p_g , $C(p)$ was numerically minimized using the `optimize` function in R [24]. p_i was found by implementing the procedures described in the electronic supplementary material, appendix G, for the various models, using R's `uniroot` function.

The calculations of both p_g and p_i depend on π_p and ψ_p , the probabilities of a delayer being infected or vaccinated, respectively (equation (5.3)). For the models in which the vaccination rate is proportional to prevalence or incidence, we used the final size relations reported in §§5.1 and 5.2, respectively, to calculate π_p and ψ_p . For the remaining models, π_p and ψ_p were obtained by numerically integrating the differential equations using the `deSolve` package [25] in R [24].

When generating figure 5, for *all* the models π_p and ψ_p were calculated by numerical integration of the differential equations.

7. Results and discussion

7.1. Group optimum versus individual equilibrium

Figure 1 shows the group optimum p_g (red) and individual equilibrium p_i (black), as the vaccination effort parameter $\phi_{\text{(model)}}$ is varied, for the different models. As expected, the group-optimal coverage is never smaller than the individual equilibrium, and both decrease as $\phi_{\text{(model)}}$ is increased. The difference, $p_i - p_g$, tends to grow initially with $\phi_{\text{(model)}}$, but eventually decreases to 0, because the coverage at both the

group optimum and individual equilibrium always drops to 0 if the vaccination rate parameter $\phi_{\text{(model)}}$ is increased sufficiently. It is also evident that the difference between the group-optimal coverage and the individual equilibrium depends strongly on the vaccination model used. In general, this difference is much smaller for the instantaneous and constant rate vaccination models than it is for the other models in which vaccination is affected by the state of the outbreak.

7.2. Mortality cost versus vaccination cost

Figure 2 shows the mortality cost (proportion of the population that dies; figure 2*a*) and the vaccination cost (proportion of the population that is vaccinated by the end of the outbreak; figure 2*b*) as functions of the vaccination response effort parameter, $\phi_{\text{(model)}}$, for the various post-outbreak response scenarios.

7.2.1. Mortality plateau

The most striking feature of figure 2 is the plateau in mortality cost at the individual equilibrium for low values of $\phi_{\text{(model)}}$. This plateau can be explained using the Bishop–Cannings theorem [12,26], which implies that if the individual equilibrium is a mixed strategy then the pay-off for vaccinating and delaying must be the same. For low values of $\phi_{\text{(model)}}$, the individual equilibrium is mixed ($0 < p_i < 1$), so the mortality cost associated with vaccinating is the same as for delaying, which is therefore the same as the overall mortality cost. Because the mortality cost for vaccinating is equal to the risk from vaccination (r_v , or r in normalized units; cf. equation (3.3) and tables 1 and 2), the overall mortality cost is constant at r_v (or r in normalized units) as long as the individual equilibrium is mixed. As $\phi_{\text{(model)}}$ is increased, the individual equilibrium p_i is decreased (see §7.1). When p_i reaches 0, there is a pure strategy equilibrium (i.e. always delay), so the Bishop–Cannings theorem no longer applies; then, the overall mortality is the mortality of delayers, which is $-a[r_i\pi_p + \psi_p r_v]$ (see equation (3.1)) and this decreases as $\phi_{\text{(model)}}$ is increased (because the epidemic is extinguished faster).

7.2.2. Public health strategy implications of the mortality plateau

There is an important implication of the plateau in mortality that occurs for small $\phi_{\text{(model)}}$ if vaccination is voluntary: in order to achieve *any* reduction in overall mortality, the post-outbreak vaccination response must be so strong that no individual would choose to vaccinate pre-emptively ($p_i = 0$, i.e. the equilibrium is for everyone to delay). Only if the post-outbreak vaccination response is already sufficiently efficient ($\phi_{\text{(model)}}$ is already sufficiently large; figure 1) can outbreak size (and hence overall mortality) be reduced by further enhancing the post-outbreak vaccination response (i.e. by increasing $\phi_{\text{(model)}}$).

Note that, for every model examined here, the right-hand (high effort) edge of the mortality plateau in figure 1 occurs for a value of vaccination effort $\phi_{\text{(model)}}$ lower than the fair comparison value (table 4). Thus, at the fair comparison values of $\phi_{\text{(model)}}$, the individual equilibrium is always to delay vaccination, and mortality can be reduced by increasing vaccination effort, $\phi_{\text{(model)}}$.

However, in §7.5, we show that any lag between the start of an outbreak and the beginning of post-outbreak vaccination extends the mortality plateau to higher vaccination efforts, $\phi_{\text{(model)}}$, and a long enough lag makes reducing mortality by increasing vaccination effort impossible. We discuss the implications of this for public health strategies further in §7.5.




Figure 1. Variation of the group optimum p_g (red) and the individual equilibrium p_i (black) with $\phi_{(\text{model})}$ (each panel presents results for a different vaccination model). Note the different ranges of $\phi_{(\text{model})}$ (on the abscissa) for different models. (Online version in colour.)

7.2.3. Generality of the mortality plateau

It is important to note that the mortality plateau described earlier is a general phenomenon that applies not only to the post-outbreak vaccination scenarios that examined here, but also to any reasonable post-outbreak vaccination scenario. More precisely, suppose public health agencies have some measure of control over a vaccination effort parameter, ϕ . Suppose also that $\phi = 0$ corresponds to no possibility of obtaining vaccine post-outbreak, and that the probabilities of a delayer being infected or vaccinated after an outbreak (π_p and ψ_p , respectively) are continuous functions of p and ϕ (for $0 \leq p < 1$ and $\phi \geq 0$). As in §3, the costs for delaying and vaccinating individuals are then $a(r_i\pi_p + r_v\psi_p)$ and r_v , respectively. Now suppose the following additionally:

- (1) If there is no possibility of being vaccinated post-outbreak ($\phi = 0$), and no one is vaccinated pre-emptively ($p = 0$),

then individuals are at greater risk than if they had been vaccinated pre-emptively (i.e. $a(r_i\pi_p + r_v\psi_p)|_{p=0, \phi=0} > r_v$).

- (2) As the initial coverage approaches 100% ($p \rightarrow 1$), the disease does not spread any further than the initial infected cohort ($\pi_p \rightarrow \alpha$). Note that, as shown in the electronic supplementary material, appendix D, this assumption holds for all of the models considered in this paper, and the mathematical argument used to show this is quite general.
- (3) The risk that a delayer is infected in the initial infection event and then dies is smaller than the risk of mortality from the vaccine alone ($\alpha r_i < r_v$).

The vaccination game with this post-outbreak vaccination scenario is a population game, and thus must have at least one Nash equilibrium [27, theorem 2.1.1, p. 24].

So for, low enough vaccination effort ϕ , if coverage p is low, it is more costly to delay than to vaccinate (from the




Figure 2. Variation of the mortality cost (proportion of the population that dies) and vaccine dose cost (proportion of the population that is vaccinated by the end of the outbreak) as functions of the vaccination effort parameter $\phi_{(\text{model})}$, for different vaccination models. Each row depicts the mortality costs (panel *a*) and vaccine dose costs (panel *b*), at the group optimum (red) and individual equilibrium (black) for one model. Note the different ranges of $\phi_{(\text{model})}$ for different models. (Online version in colour.)

first assumption above). Conversely, if coverage p is high enough, the third assumption above implies that delaying is preferable to vaccinating pre-emptively. It follows that any individual equilibrium that results from the vaccination game is a mixed NE ($0 < p_i < 1$). The preceding argument presented in §7.2.1 (using the Bishop–Cannings theorem) now implies the existence of a plateau in mortality.

7.2.4. Vaccination cost plateau

Figure 2*b* also show a plateau for sufficiently *large* vaccination efforts (except for the constant rate vaccination model). Unlike the mortality plateau, this vaccination cost plateau is not rigorously a constant (it changes very slightly as a function of $\phi_{(\text{model})}$), but it is certainly a plateau for all intents and purposes. This plateau occurs because overall

vaccination rises with the vaccination effort, $\phi_{(\text{model})}$, and cannot exceed $V_\infty = 1$, so vaccination costs must eventually taper off.

7.2.5. Perceived versus real risks

The general public is likely to overestimate vaccine-induced mortality [28–30], which would tend to decrease the pre-outbreak vaccine coverage under voluntary vaccination. The game-theoretical framework we employ assumes individuals behave rationally and possess perfect information on which to base their decisions, but it is possible to relax the assumption of perfect information while maintaining that of rationality. Thus, to account for misinformation regarding the dangers of vaccination (possibly as a result of vaccine scares), we can interpret r_i and r_v as the *perceived* risks of infection and vaccination (rather than the actual risks) to predict the effective level of vaccine coverage prior to an




Figure 3. Variation of relative difference in mortality and vaccine dose costs with $\phi_{(model)}$, for different vaccination models. Each row depicts the relative mortality cost difference (panel *a*) and relative vaccine dose cost difference (panel *b*) for one model. Note the different ranges of $\phi_{(model)}$ for different models.

outbreak (note that perceived risks are to be used to predict the individual equilibrium, p_i , but not when predicting the group optimum p_g , nor when predicting the mortality and vaccination costs at *either* of these coverages). Consequently, public health agencies can potentially reduce mortality by attempting to influence the public's estimate of r (the risk of vaccination relative to infection). For example, risk perception might be influenced by a media campaign aiming to increase the accuracy of the public's perception of vaccine safety and promote pre-emptive vaccination.

7.3. Comparison of relative costs

In figure 3*a*, we look at the relative mortality cost difference, that is, in units of the cost of optimal mandatory vaccination. Explicitly, we examine how $(C(p_i) - C(p_g))/C(p_g)$ varies with $\phi_{(model)}$ for each model. Similarly, we plot the relative difference in vaccination ($V_\infty(p_i) - V_\infty(p_g)/V_\infty(p_g)$) (figure 3*b*), which is the relative vaccine dose cost difference between voluntary and mandatory vaccination.

7.3.1. Large variation in relative mortality cost

Observe that, in figure 3*a*, the relative mortality cost difference is always non-negative (as expected from the definition of the group optimum as the pre-outbreak coverage for which expected mortality cost is minimal). There is substantial variability among the models in the dependence of the relative mortality cost differences on the vaccination parameter $\phi_{(model)}$. In particular, if vaccination rate is proportional to incidence or prevalence, variation in relative mortality cost is an order of magnitude smaller than if vaccination is instantaneous or at a constant rate. The vaccination scenario that exhibits the largest variation in relative mortality costs is instantaneous vaccination. In this scenario, a voluntary vaccination policy could result in over 150% more deaths than if vaccination were mandatory.

7.3.2. Modest variation in relative vaccine dose cost

There is also substantial variability in the pattern of variation of relative vaccine dose cost as a function of $\phi_{(model)}$ among the

different models (figure 3*b*). However, for all the models, variation in relative vaccine dose cost as a function of $\phi_{(\text{model})}$ is much less than the corresponding variation in relative mortality cost. The maximum variation in relative vaccine dose cost reaches approximately 16% for the models in which vaccination is proportional to prevalence or incidence. This relatively large variation can be attributed to low pre-outbreak vaccination coverage (at the individual equilibrium) causing high disease prevalence and incidence; consequently, because vaccination rate is proportional to prevalence or incidence, there is correspondingly high post-outbreak vaccination, which overshoots that which would be required to minimize group mortality. In these two situations, the vaccine dose cost at the individual equilibrium can be greater than at the group optimum. In any case, the relatively small difference in overall vaccine dose costs, both as a function of vaccination effort ($\phi_{(\text{model})}$) and among vaccination scenarios (see figure 2*b*), suggests that vaccine dose cost should probably not be a factor in public health policy decisions.

7.4. Vaccine dose cost as a function of mortality cost

Figures 2 and 3 present mortality costs and vaccine dose costs as functions of vaccination effort for the various models. Because the meaning of the vaccination effort parameter $\phi_{(\text{model})}$ differs among models, it is not straightforward to make meaningful comparisons among the various models (which is why we calculated ‘fair comparison’ values in §6). In this section, we display results for the various models, factoring out the vaccination effort parameter. For each model, figure 4 shows the vaccine dose cost as a function of mortality cost. In health economics terms, this can be considered a *cost-effectiveness analysis* [31].

In figure 4, the squares indicate the point in the mortality-cost–vaccination-cost plane where the vaccination effort ($\phi_{(\text{model})}$) is the lowest that we considered. Increasing vaccination effort (while remaining at the individual equilibrium or the group optimum) corresponds to moving away from the square along the plotted curves.

The graphs in figure 4 allow us to answer practical questions such as ‘If we want to ensure that no more than one in every 10 million citizens dies, how many vaccine doses are required in each scenario?’ or ‘If we have a stockpile of vaccine doses sufficient for 30% of the population, what percentage of the population can be expected to die if there is an outbreak in each scenario?’ Of course, by construction the graphs do not indicate how much effort ($\phi_{(\text{model})}$) is required to achieve the desired results. We emphasize that—as shown in the previous section—vaccine dose cost at the individual equilibrium or group optimum hardly varies as a function of vaccination effort ($\phi_{(\text{model})}$), so the ‘practical’ questions are not necessarily well posed (e.g. if we have sufficient vaccine doses for only 30% of the population, then neither the individual equilibrium nor the group optimum can ever be achieved). This is true for all the models *with parameters appropriate for smallpox*; for another disease graphs like figure 4 could have genuine practical value for public health policy analysis (e.g. setting $\mathcal{R}_0 = 1.25$ and keeping all other model parameters as in tables 1 and 2 causes the vaccination cost to vary between 25% and 99.999%).

In figure 4, when the vaccination rate is proportional to either prevalence or incidence, note that, as the vaccination effort, $\phi_{(\text{model})}$, increases, two phases of behaviour are apparent for the costs at both p_i and p_g : first, vaccine dose cost

rises but no change in mortality cost is observed (this is caused by the plateau in mortality described in §7.2.1). Then, for all but the instantaneous vaccination model, vaccine dose cost remains virtually constant (note the differences in the scales of the vertical axes among the various panels), but mortality costs decrease.

It is interesting to note that the dependence of vaccine costs on mortality costs at the group optimum varies among the models. For example, when vaccination is proportional to remaining susceptibles, and for the constant rate vaccination model, we see in figure 4 that at the group optimum, as mortality cost is decreased, vaccine dose cost decreases at first, but then increases. Thus, in these situations, one can lower both the mortality and the vaccine dose cost at the same time by increasing vaccination effort (in health economics terms, the decision to use higher vaccination effort has negative marginal cost in vaccine doses per life saved). This contrasts the models in which vaccination is instantaneous, or proportional to incidence or prevalence, in which we observe that, as mortality cost is decreased, the vaccine dose cost at the group optimum remains constant and then increases sharply.

Finally, for the instantaneous vaccination model, there is a range of vaccination efforts for which one can reduce mortality without increasing vaccine dose costs at the group optimum. In this parameter range, the increase in vaccine dose cost necessary to decrease mortality at the individual equilibrium is small at first, but grows larger as mortality is decreased.

7.5. Effect of vaccination response lag t_{lag}

We have implicitly assumed that in any of the scenarios we have considered the vaccination response will begin as soon as an outbreak is seeded by a bioterrorist attack or accidental release. In contrast, Bauch *et al.* [4] assumed a lag of two weeks between the seeding of an outbreak and the initiation of a vaccination response. In this section, we investigate the effect of a *response lag* of t_{lag} days between an outbreak being seeded and the post-outbreak vaccination campaign beginning (so far, we have assumed $t_{\text{lag}} = 0$ days; in [4], $t_{\text{lag}} = 14$ days was assumed).

Intuitively, adding a lag between the beginning of an outbreak and the vaccination response allows the disease to spread unhindered for some time, which increases the probability of delayers being infected, thus decreasing the pay-off for delaying. As a result, the individual equilibrium p_i increases, which consequently extends the mortality plateau (§7.2.1) to higher values of $\phi_{(\text{model})}$.

7.5.1. The critical lag, t_{lag}^*

For a disease such as smallpox with $\mathcal{R}_0 \sim 5$, the expected final size of an uncontrolled epidemic is greater than 99.9% of the population. If no one is pre-emptively vaccinated, and the response lag after an outbreak is seeded is sufficiently long, almost everyone will have been infected before the response begins, i.e. if t_{lag} is sufficiently long then delayers will almost certainly be infected before they can be vaccinated. Consequently, unless the probability of an outbreak (a) is negligible, delaying will be riskier than vaccinating pre-emptively, so the individual equilibrium will not be for everyone to delay: we will certainly have $p_i > 0$. It follows that for response lags longer than some *critical lag*, t_{lag}^* , mortality cannot be reduced no matter how much effort is applied in the post-outbreak vaccination response




Figure 4. Vaccine dose cost as a function of mortality cost at the group optimum, p_g (red), and the individual equilibrium, p_i (black), for the different models. Squares represent values at lowest $\phi_{(\text{model})}$ simulated. (Online version in colour.)

(i.e. the mortality plateau described in §7.2.1 continues for arbitrarily large values of $\phi_{(\text{model})}$).

A more precise argument allows us to estimate t_{lag}^* . Suppose the initial coverage is $p = 0$ (no pre-emptive vaccination). If the risk of becoming infected and dying is larger than the risk from vaccinating, i.e. $ar_i\pi_0 > r_v$ (or, equivalently, $\pi_0 > r/a$), then delaying will not be the individual equilibrium. For any vaccination scenario, the delayers' probability of being infected by the end of the outbreak (equation (5.3a)) is greater than or equal to their probability of being infected before the vaccination response begins (at time t_{lag}),

$$\pi_0 \geq \left(I(t_{\text{lag}}) + R(t_{\text{lag}}) \right) \Big|_{p=0}. \quad (7.1)$$

Therefore, if

$$\left(I(t_{\text{lag}}) + R(t_{\text{lag}}) \right) \Big|_{p=0} > \frac{r}{a}, \quad (7.2)$$

then $\pi_0 > r/a$ and delaying is guaranteed not to be the individual equilibrium. But for *any* post-outbreak vaccination scenario that includes a response lag, when $t < t_{\text{lag}}$ the removed proportion of the population, $R(t)$, follows the standard SIR model solution (with no vaccination). For the SIR model with

no vaccination ($p = 0$; a , α and \mathcal{R}_0 as in table 1), numerical simulation shows that equation (7.2) is satisfied for $t_{\text{lag}} \gtrsim 15.1$ days. Hence, if the public health response lag is 16 days or longer, then it is guaranteed that (regardless of the vaccination scenario or corresponding vaccination effort, $\phi_{(\text{model})}$) delaying will not be the individual equilibrium.

We emphasize that our estimate of 16 days as an upper bound for t_{lag}^* depends on a number of factors, including:

- the probability of an outbreak (a);
- the proportion of susceptibles infected in the initial outbreak (α); and
- the epidemiological model: the estimate is obtained using the SIR model, but adding an exposed class (SEIR) with parameters as in table 1 increases the critical lag. Repeating the calculation for the SEIR model yielded the upper bound $t_{\text{lag}}^* \leq 26.3$ days. The reason for this difference in critical lags is that, when the outbreak is seeded, all individuals initially infected begin their latent period simultaneously, and take on average 15 days to become infectious.

Thus, $t_{\text{lag}}^* < 16$ days should be regarded as a rough estimate at best. Nonetheless, the existence of a critical lag,




Figure 5. Variation of the effective critical lag \tilde{t}_{lag}^* (§7.5.2) with vaccination effort $\phi_{(model)}$. (a) Effective critical lag versus percentage of the fair comparison vaccination effort when the fair comparison vaccination effort is defined. (b) Effective critical lag versus vaccination effort ϕ_{inst} when vaccination is instantaneous. (Online version in colour.)

beyond which it is impossible to reduce mortality by increasing vaccination effort, is an important consideration for public health agencies, in devising contingency plans for post-outbreak vaccination against diseases.

Note, however, that, in the case of a bioterrorist attack, an outbreak will probably not be discovered until individuals show symptoms, i.e. until someone's latent period has passed (12 days at a minimum). Taking this delayed detection into account, it follows that in order to avoid extending the mortality plateau to all feasible values of vaccination effort, ϕ_{const} , the response lag from discovery of the epidemic to the beginning of the post-outbreak vaccination response must, in practice, be substantially shorter than 26 days. This is in contrast to an accidental release, where public health authorities might know of the outbreak well before anyone has shown symptoms. In this latter case, because it is more likely that the critical lag has not been exceeded, it is especially important to begin the vaccination response as early as possible in order to reduce mortality.

Lastly, it is important to note that the effect of a response lag on the mortality plateau presupposes that both the vaccination effort and the response lag are known to the public in advance. This limits the applicability of this effect, because, in the case of a bioterrorist attack, the response lag probably

depends on when an infective first shows symptoms (which introduces a stochastic effect). Further analysis would be needed to determine the effects of a stochastic response lag on individual behaviour, and thus on mortality.

7.5.2. The effective critical lag, \tilde{t}_{lag}^*

We have seen that if the response lag is longer than the critical lag ($t_{lag} > t_{lag}^*$), then no matter how large the vaccination effort ($\phi_{(model)}$), it is impossible to reduce mortality. Of course, in practice, the vaccination effort cannot be arbitrarily large and will be constrained by public health resources. Given a maximum *feasible* vaccination effort, it would be helpful to know how long the response lag can be before the mortality plateau extends to all feasible levels of vaccination effort.

To address this issue, we define the *effective critical lag*, \tilde{t}_{lag}^* , to be the minimal response lag, such that the individual equilibrium is no longer to delay (i.e. $p_i > 0$) given a maximum *feasible vaccination effort* $\phi_{(model)}$. Thus, the critical lag t_{lag}^* is the limit of \tilde{t}_{lag}^* as the maximum feasible vaccination effort becomes arbitrarily large.

In figure 5, we plot the effective critical lag, \tilde{t}_{lag}^* , against the vaccination effort, $\phi_{(model)}$, for the various models

examined in this paper. For the models for which fair comparison values of $\phi_{(\text{model})}$ are well defined (see §6.2), we used these as estimates for feasible vaccination efforts. However, because the fair comparison level of vaccination effort is a crude estimate for the range of feasible vaccination efforts, in figure 5a, we plot the effective critical lag at values of $\phi_{(\text{model})}$ ranging from 50% to 150% of the fair comparison levels of vaccination efforts for the various models, in increments of 10% of the fair comparison level of $\phi_{(\text{model})}$.

The instantaneous vaccination model was the only model for which a fair comparison value of vaccination effort ϕ_{inst} could not be defined (see §6.2). For this model, we show the effective critical lag $\widetilde{t}_{\text{lag}}^*$ for ϕ_{inst} ranging from 0.8 to 1 (if $\phi_{\text{inst}} < 0.8$, then $\widetilde{t}_{\text{lag}}^* < 1$ day) in figure 5b.

We see in figure 5 that, for some vaccination scenarios, minimizing the response lag t_{lag} is essential: even a short lag extends the mortality plateau to all feasible vaccination effort levels, making it impossible to reduce mortality by increasing effort after the lag. We also note that, for some scenarios, a good estimate of the attainable vaccination effort is necessary, because the critical effective lag is very sensitive to the vaccination effort. These two facts further underline the importance of accurately modelling post-outbreak vaccination to inform public health decisions relating to post-outbreak contingency plans. When the response lag is longer than the effective critical lag ($t_{\text{lag}} \geq \widetilde{t}_{\text{lag}}^*$), the only plausible way for public health officials to decrease mortality (while allowing individuals to choose whether or not to vaccinate) is to reduce the relative mortality risk (by decreasing the probability of dying from vaccination, i.e. developing a safer vaccine).

7.5.3. The response lag should be minimized

Based on §§7.5.1 and 7.5.2, reducing the response lag lowers expected mortality and makes it easier to decrease mortality further:

- For vaccination efforts higher than the end of the mortality plateau, everyone will choose to delay vaccination (§7.2.1). From the discussion leading up to equation (7.1), it follows that, even in the best-case scenario where the epidemic is stopped immediately at $t = t_{\text{lag}}$, mortality will be no less than

$$r_i(I(t_{\text{lag}}) + R(t_{\text{lag}}))|_{p=0}, \quad (7.3)$$

which increases with the response time t_{lag} . Thus, increasing the response lag increases the lowest attainable mortality (even if vaccination effort can be increased without bound).

- Increasing the response lag increases the vaccination effort at the end of the mortality plateau (i.e. the minimal vaccination effort beyond which increasing vaccination effort decreases mortality). Thus, longer response lags make it harder to achieve a decrease in mortality.

However, note that if a response time lower than the effective critical lag ($t_{\text{lag}} < \widetilde{t}_{\text{lag}}^*$) cannot be achieved, neither increasing the vaccination effort, $\phi_{(\text{model})}$, nor decreasing the response time, t_{lag} , can decrease mortality.

8. Conclusion

We have analysed five distinct scenarios (§2) associated with a potential smallpox outbreak triggered by a bioterrorist attack

or accidental release. The scenarios differ in the factors that influence individuals' perception of risk and how a post-outbreak vaccination response plays out. We examined these scenarios both with and without an assumed lag between an outbreak starting and a public health response being initiated. Our work generalizes the analysis of Bauch *et al.* [4], who investigated a single scenario with a response lag of 14 days.

As in [4], we considered separately group interest (optimal strategies for minimizing overall mortality) and self-interest (stable strategies for individual choices with respect to pre-emptive vaccination). From each perspective, we obtained the (imposed or expected) pre-emptive vaccination coverage (p) for each scenario (the group optimum p_g in the case of group interest and the individual equilibrium p_i in the case of self-interest; figure 1).

Our principal conclusions are the following.

- (1) For a given level of post-outbreak vaccination effort, the group optimum pre-emptive coverage is always greater than the individual equilibrium ($p_g > p_i$; figure 1) and the expected total mortality is always less if public health authorities impose the group optimum rather than letting individuals make their own vaccination decisions (figure 2a). If no outbreak occurs, then some people will die unnecessarily from pre-emptive vaccination. Given the difficulty of estimating the probability of an attack or accidental release, it would be hard for governments to justify an imposed pre-emptive vaccination policy for a disease like smallpox for which the vaccine can cause death.
- (2) The number of vaccine doses required at the group optimum and individual equilibrium does not vary substantially as a function of vaccination effort (e.g. speed of vaccine distribution post-outbreak) for any of the scenarios (figure 2b). Consequently, the economic cost of vaccine production is not likely to play a significant role in policy decisions.
- (3) Total expected mortality as a function of vaccination effort depends strongly on which scenario is considered (figure 2a). Some vaccination scenarios are affected by the public reaction to media reports on the epidemic's progress, whereas some (the instantaneous and constant rate vaccination scenarios) are under the direct control of public health authorities. To assist public health authorities preparing for potential outbreaks, further research is needed to determine which factors have the greatest influence on individuals' perception of risk and which post-outbreak vaccination strategies are most feasible.
- (4) For any realistic vaccination scenario, there is a range of vaccination effort levels in which increasing vaccination effort does not reduce overall mortality. In this mortality plateau, increasing vaccination effort leads only to fewer people vaccinating pre-emptively, until the individual equilibrium becomes to delay vaccination (at which point it is possible to reduce mortality by increasing the vaccination effort). Thus, under voluntary vaccination, in order for public health authorities to expect to reduce mortality by increasing vaccination effort post-outbreak, their planned post-outbreak vaccination response must be so efficient that no one would choose to vaccinate pre-emptively ($p_i = 0$).
- (5) Any lag between the beginning of an outbreak and the post-outbreak vaccination response makes it harder for higher vaccination effort levels to make a difference to overall mortality, and a large enough lag will make it

impossible to reduce mortality, *regardless* of the level of vaccination effort. Given a maximum feasible vaccination effort level, there is an effective critical lag, beyond which it is impossible to reduce mortality by increasing vaccination effort. The dependence on the post-outbreak vaccination scenario, of both the effective critical lag at feasible levels of vaccination effort and the effect of changes in vaccination effort on the effective critical lag, further highlights the importance of researching realistic post-outbreak vaccination responses.

It is not possible to know with certainty how governments and health agencies will react, or how individuals

will behave, in the event of an outbreak. However, the above-mentioned conclusions are based on our analysis of five distinct post-outbreak scenarios (and some model features that are much more generic), so it seems likely that our conclusions would remain valid if further plausible scenarios were considered.

Acknowledgements. Some preliminary work on the problem addressed in this paper was carried out by Adelia Yu as part of her undergraduate summer research project in 2004. We are grateful to Sigal Balshine, Paul Higgs, Rufus Johnstone and two anonymous referees for valuable comments.

Funding statement. This study is supported by NSERC (D.J.D.E.) and the Trillium Foundation (C.M.).

References

- Fenner F. 1988 Smallpox and its eradication, Parts 1–14. In *History of international public health*, vol. 6. Geneva, Switzerland: World Health Organization.
- Koplow DA. 2004 *Smallpox: the fight to eradicate a global scourge*. Oakland, CA: University of California Press.
- Christensen J. 2014 *CDC: smallpox found in NIH storage room is alive*. <http://www.cnn.com/2014/07/11/health/smallpoxfound-nih-alive/> (accessed 12 August 2014).
- Bauch CT, Galvani AP, Earn DJD. 2003 Group interest versus self-interest in smallpox vaccination policy. *Proc. Natl Acad. Sci. USA* **100**, 10564–10567. (doi:10.1073/pnas.1731324100)
- Bauch CT, Earn DJD. 2004 Vaccination and the theory of games. *Proc. Natl Acad. Sci. USA* **101**, 13 391–13 394. (doi:10.1073/pnas.0403823101)
- Anderson RM, May RM. 1991 *Infectious diseases of humans: dynamics and control*. Oxford, UK: Oxford University Press.
- Eichner M, Dietz K. 2003 Transmission potential of smallpox: estimates based on detailed data from an outbreak. *Am. J. Epidemiol.* **158**, 110–117. (doi:10.1093/aje/kwg103)
- Gani R, Leach S. 2001 Transmission potential of smallpox in contemporary populations. *Nature* **414**, 748–751. (doi:10.1038/414748a)
- Krylova O. 2011 Predicting epidemiological transitions in infectious disease dynamics: smallpox in historic London (1664–1930). PhD thesis, McMaster University, Canada.
- Kaplan EH, Craft DL, Wein LM. 2002 Emergency response to a smallpox attack: the case for mass vaccination. *Proc. Natl Acad. Sci. USA* **99**, 10 935–10 940. (doi:10.1073/pnas.162282799)
- Fudenberg D, Tirole J. 1991 *Game theory*. Cambridge, MA: MIT Press.
- Maynard Smith J. 1982 *Evolution and the theory of games*. Cambridge, UK: Cambridge University Press.
- Krylova O, Earn DJD. 2013 Effects of the infectious period distribution on predicted transitions in childhood disease dynamics. *J. R. Soc. Interface* **10**, 20130098. (doi:10.1098/rsif.2013.0098)
- World Health Organization. 2014 *World health statistics 2014*. Geneva, Switzerland: WHO.
- Hammarlund E, Lewis MW, Hansen SG, Strelow LJ, Nelson JA, Sexton GJ, Hanifin JM, Slika MK. 2003 Duration of antiviral immunity after smallpox vaccination. *Nat. Med.* **9**, 1131–1137. (doi:10.1038/nm917)
- Kermack WO, McKendrick AG. 1927 A contribution to the mathematical theory of epidemics. *Proc. R. Soc. Lond. A* **115**, 700–721. (doi:10.1098/rspa.1927.0118)
- Ma J, Earn DJD. 2006 Generality of the final size formula for an epidemic of a newly invading infectious disease. *Bull. Math. Biol.* **68**, 679–702. (doi:10.1007/s11538-005-9047-7)
- Earn DJD, Andrews PW, Bolker BM. 2014 Population-level effects of suppressing fever. *Proc. R. Soc. B* **281**, 20132570. (doi:10.1098/rspb.2013.2570)
- Weisstein EW. Lambert W-function. From MathWorld: a Wolfram Web Resource. <http://mathworld.wolfram.com/LambertW-Function.html> (accessed 17 April 2015).
- Corless RM, Gonnet GH, Hare DEG, Jeffrey DJ, Knuth DE. 1996 On the Lambert W function. *Adv. Comput. Math.* **5**, 329–359. (doi:10.1007/BF02124750)
- McNeil Jr DG. 2013 Wary of attack with smallpox, U.S. buys up a costly drug. *The New York Times*. See <http://www.nytimes.com/2013/03/13/health/us-stockpiles-smallpox-drug-in-case-of-bioterror-attack.html> (accessed 12 August 2014).
- Vink MA, Bootsma MJC, Wallinga J. 2014 Serial intervals of respiratory infectious diseases: a systematic review and analysis. *Am. J. Epidemiol.* **180**, 865–875. (doi:10.1093/aje/kwu209)
- Svensson A. 2007 A note on generation times in epidemic models. *Math. Biosci.* **208**, 300–311. (doi:10.1016/j.mbs.2006.10.010)
- R Core Team. 2013 *R: a language and environment for statistical computing*. Vienna, Austria. See <http://www.R-project.org/>.
- Soetaert K, Petzoldt T, Setzer RW. 2010 Solving differential equations in R: package deSolve. *J. Stat. Softw.* **33**, 1–25.
- Sandholm WH. 2010 *Population games and evolutionary dynamics*. Economic Learning and Social Evolution. Cambridge, MA: MIT Press.
- Bishop DT, Cannings C. 1978 A generalized war of attrition. *J. Theor. Biol.* **70**, 85–124. (doi:10.1016/0022-5193(78)90304-1)
- Leach M, Fairhead J. 2007 *Vaccine anxieties: global science, child health and society*. Science in Society Series. London, UK: Earthscan.
- Brown KF, Kroll JS, Hudson MJ, Ramsay M, Green J, Long SJ, Vincent CA, Fraser G, Sevdails N. 2010 Factors underlying parental decisions about combination childhood vaccinations including MMR: a systematic review. *Vaccine* **28**, 4235–4248. (doi:10.1016/j.vaccine.2010.04.052)
- Gangarosa EJ, Galazka A, Wolfe C, Phillips L, Gangarosa R, Miller E, Chen RT, Gangarosa RE. 1998 Impact of anti-vaccine movements on pertussis control: the untold story. *Lancet* **351**, 356–361. (doi:10.1016/S0140-6736(97)04334-1)
- Phillips CJ. 2008 *Health economics: an introduction for health professionals*. London, UK: John Wiley & Sons.

Game theory of pre-emptive vaccination before
bioterrorism or accidental release of smallpox

Electronic Supplementary Material

Journal of the Royal Society Interface (2015)

Chai Molina and David J.D. Earn

Department of Mathematics and Statistics

McMaster University

Hamilton, Ontario, Canada, L8S 4K1

April 7, 2015 @ 17:51

APPENDICES

A	Lambert W function	3
B	Interpretation of vaccination effort parameters	4
B.1	ϕ_{susc}	4
C	Convergence to disease-free equilibrium	4
D	Calculation of π_1	5
E	Final size relations, π_p and ψ_p	7
E.1	Vaccination rate \propto disease prevalence	7
E.1.1	Final size relations	7
E.1.2	Qualitative behaviour of S_∞^1 for high vaccine coverage	12
E.1.3	π_p decreases and ψ_p increases with p when $p_m \geq 1$	13
E.1.4	Behaviour of $p_0(-1)$ as $\phi_{\text{prev}} \rightarrow 0^+$	14
E.2	Vaccination rate \propto incidence	14
E.3	Vaccination rate \propto proportion still susceptible	16
E.4	Instantaneous vaccination of a proportion ϕ_{inst} of the population	18
F	Maximal vaccination rate for fair comparison of models	20
F.1	Maximal Vaccination rate when $\dot{V} = \phi_{\text{prev}}I$	20
F.2	Maximal Vaccination rate when $\dot{V} = \phi_{\text{inc}}SI$	21
G	The individual equilibrium	24
G.1	Vaccination rate \propto disease prevalence	24
G.1.1	$p_m \geq 1$	24
G.1.2	$p_m \leq 1$ and $p_0 \leq 0$	26
G.1.3	$p_m \leq 1$ and $p_0 \in (0, 1)$	27
G.2	Vaccination rate \propto incidence	29
G.3	Vaccination rate \propto proportion still susceptible	30
G.4	Instantaneous vaccination of a proportion ϕ_{inst} of the population	30
G.5	constant rate vaccination	31
H	The group optimum	31
H.1	Vaccination rate \propto disease prevalence	31
	References	33

1 A Lambert W function

2 The standard final size relation, which can be derived from the SIR model [1] and many
3 other epidemic models [2], is

$$4 \quad Z = 1 - e^{-\mathcal{R}_0 Z}. \quad (\text{S1})$$

5 Here, Z is the final size ($Z = 1 - S_\infty$) and \mathcal{R}_0 is the basic reproduction number. Z can be
6 expressed explicitly as a function of \mathcal{R}_0 [2, 3],

$$7 \quad Z(\mathcal{R}_0) = 1 + \frac{1}{\mathcal{R}_0} W[-\mathcal{R}_0 e^{-\mathcal{R}_0}], \quad (\text{S2})$$

8 where the Lambert W function [4, 5] is the inverse function of

$$9 \quad f(W) = W e^W. \quad (\text{S3})$$

10 Use of the Lambert W function is critical for our derivations of final size formulae for models
11 we consider here. $W(x)$ is real for $x \geq -1/e \simeq -0.368$ and is two-valued for $-1/e < x < 0$.
12 The upper “principal” branch, for which $W(x) \geq -1$, is denoted W_0 and the lower branch
13 is denoted W_{-1} . See Figure S1.




Figure S1: The Lambert W function, showing the principal branch W_0 and the secondary branch W_{-1} .

14 B Interpretation of vaccination effort parameters

15 The vaccination effort parameters are explained § 5. “Fair comparison” values for these
 16 parameters are derived in § 6.2 and listed in Table 3.

17 B.1 ϕ_{susc}

18 In § 6.2, we commented that ϕ_{susc} can be considered to be the probability per unit time of a
 19 delayer being vaccinated. To see this, note that the probability (p_{vacc}) of a susceptible delayer
 20 being vaccinated in the time interval $[t, t + \Delta t]$ is the ratio of the number of susceptibles
 21 vaccinated in that time interval, $V(t + \Delta t) - V(t)$, to the number of susceptibles present at
 22 the beginning of that time interval, $S(t)$. Thus,

$$23 \quad p_{\text{vacc}} = \frac{V(t + \Delta t) - V(t)}{S(t)} = \frac{V(t + \Delta t) - V(t)}{\Delta t} \frac{1}{S(t)} \Delta t.$$

24 Since $\lim_{\Delta t \rightarrow 0} \frac{V(t + \Delta t) - V(t)}{\Delta t} = \dot{V} = \phi_{\text{susc}} S$, for small Δt , we have $p_{\text{vacc}} \approx \phi_{\text{susc}} \Delta t$. Thus, ϕ_{susc}
 25 is the (constant) probability per unit time of a delayer being vaccinated.

26 C Convergence to disease-free equilibrium

27 In this appendix, we show that for all models considered, the epidemic must eventually die
 28 out (*i.e.*, the system converges to a disease-free equilibrium).

29 Consider the $SIRV$ model given by the differential equations

$$30 \quad \dot{S} = -\beta SI - \dot{V}, \tag{S4a}$$

$$31 \quad \dot{I} = \beta SI - \gamma I, \tag{S4b}$$

$$32 \quad \dot{R} = \gamma I, \tag{S4c}$$

$$33 \quad \dot{V} = f(t, S, I, R, V), \tag{S4d}$$

34 where f is continuously differentiable and satisfies $f(t, S, I, R, V) \geq 0$, $f(t, 0, I, R, V) = 0$
 35 whenever $t \geq 0$, $S \geq 0$, $I \geq 0$, $R \geq 0$, $V \geq 0$. From the fundamental existence and uniqueness
 36 theorem [6], there is a unique solution to Equation (S4) for any non-negative initial conditions
 37 $S(0) \geq 0$, $I(0) \geq 0$, $R(0) \geq 0$ and $V(0) \geq 0$. Suppose also that $S(0) + I(0) + R(0) + V(0) = 1$.

38 First, note that $S(t) \geq 0$ for all $t \geq 0$. To see this, suppose in order to derive a
 39 contradiction that $S(T) < 0$ for some $T > 0$. Then, since $S(t)$ is continuous, there must be
 40 some time $0 \leq \tau < T$ such that $S(\tau) = 0$. But because $\dot{S}|_{S=0} = 0$ it follows that $S(t) = 0$ for
 41 all $t \geq \tau$, which contradicts $S(T) < 0$.

42 Similarly, it follows that $I(t) \geq 0$. Consequently, $R(t)$ is nondecreasing in t , and in
 43 particular, $R(t) \geq 0$. Lastly, $\dot{V} \geq 0$ so $V(t) \geq 0$ as well.

44 Now, $\frac{d}{dt} (S(t) + I(t) + R(t) + V(t)) = 0$ for all t , so $S(0) + I(0) + R(0) + V(0) = 1$ implies
 45 $S(t) + I(t) + R(t) + V(t) = 1$ for all t . Consequently, $S(t)$, $I(t)$, $R(t)$ and $V(t)$ each lie in
 46 the interval $[0, 1]$ for all time.

48 In addition to being bounded, $S(t)$, $R(t)$ and $V(t)$ are monotonic (their time derivatives
49 are non-positive) and therefore have a limit as $t \rightarrow \infty$. It follows that $I(t)$ also has such a limit
50 ($I = 1 - S - V - R$). To see that this limit is 0, suppose instead that $I_\infty = \lim_{t \rightarrow \infty} I(t) > 0$.
51 Then, $\lim_{t \rightarrow \infty} \dot{R} = \gamma I_\infty$. Thus, there exists a time t_* such that $\dot{R}(t) > \gamma I_\infty/2$ for all $t_* < t$.
52 This implies that the proportion in the recovered class increases at least linearly, and must
53 eventually hit $R = 1$ (no later than at time $t_* + 2/(\gamma I_\infty)$) and be greater than 1 thereafter.
54 However, this contradicts the fact that the proportion of the population in any class cannot
55 exceed 1. Thus $I_\infty = \lim_{t \rightarrow \infty} I(t) = 0$.

56 A similar argument can be applied to the constant rate $SEIRV$ model by noting that

- 57 1. If $S(0) \geq 0$, $E(0) \geq 0$, $I(0) \geq 0$, $R(0) \geq 0$ and $V(0) \geq 0$, then S , E , I , R and V
58 remain non-negative for all time.
- 59 2. If $S(0) + E(0) + I(0) + R(0) + V(0) = 1$, then $S + E + I + R + V = 1$ for all time.
- 60 3. $S(t) \rightarrow 0$ in finite time in this model. To see this, suppose in order to derive a
61 contradiction, that $S(t) > 0$ for all time $t \geq 0$. Then, for all $t > t_{\text{lag}}$, $\dot{S} = \phi_{\text{const}}$,
62 and thus $S(t) \geq \phi_{\text{const}}(t - t_{\text{lag}})$. Consequently, $S(t) > 1$ for all $t > t_{\text{lag}} + 1/\phi_{\text{const}}$,
63 contradicting the fact that $S(t) \leq 1$ for all t .
- 64 4. Since $S(t) \rightarrow 0$ in finite time in this model, it follows that after a finite time $\dot{E} = -\sigma E$,
65 implying that $\lim_{t \rightarrow \infty} E = 0$. Since R and V are monotonic and thus have a limit as
66 $t \rightarrow \infty$, it follows that $\lim_{t \rightarrow \infty} I = \lim_{t \rightarrow \infty} 1 - S - E - R - V$ exists as well, and one
67 can continue as before.

68 Lastly, a corollary of this convergence to the disease-free equilibrium is that $S_\infty = \lim_{t \rightarrow \infty} S(t)$
69 is well defined and

$$70 \quad S_\infty < \frac{\gamma}{\beta} = \frac{1}{\mathcal{R}_0}. \quad (\text{S5})$$

71 To see this, observe that since S is monotonic and bounded, it must converge to some
72 finite limit within $[0, 1]$, so S_∞ is well defined. Next, note that $I(0) = \alpha(1 - p) > 0$ and
73 $\lim_{t \rightarrow \infty} I(t) = I_\infty = 0$, so there is some time t_* at which $\dot{I}(t_*) < 0$ and so $S(t_*) < \gamma/\beta$. But
74 $\dot{S} \leq 0$ and so for any $t_* < t$, we have $S(t) < \gamma/\beta$. Thus, Equation (S5) follows because of
75 the monotonicity of S .

76 D Calculation of π_1

77 In § 5, we stated that as pre-emptive vaccination approaches full coverage ($p \rightarrow 1$), the
78 probability of a delayer being infected (π_p) approaches the proportion of susceptibles initially
79 infected in an outbreak (α). Recalling the definition of π_1 in Equation (12), the claim is that
80 for all the models considered

$$81 \quad \pi_1 = \lim_{p \rightarrow 1^-} \pi_p = \alpha. \quad (\text{S6})$$

82 To verify Equation (S6), first consider the SIRV models defined in Equation (9). The
83 proportion of individuals who are eventually removed (R_∞) must be greater than the number

84 initially infected (Equation (10b)), so

$$85 \quad R_\infty \geq I(0) = \alpha(1-p). \quad (\text{S7})$$

86 Thus from Equation (11a), we have $\pi_p \geq \alpha$ for any $p \in [0, 1)$. It follows that $\pi_1 \geq \alpha$ if the
87 limit exists.

88 We now show that $\pi_1 \leq \alpha$. The basic reproduction number is $\mathcal{R}_0 = \beta/\gamma$ and the effective
89 reproduction number when the outbreak begins is (Equation (10a))

$$90 \quad \mathcal{R}_{\text{eff}}(0) = \mathcal{R}_0 S(0) = \mathcal{R}_0(1-p)(1-\alpha). \quad (\text{S8})$$

91 Thus, if $p > 1 - 1/\mathcal{R}_0$ then $\mathcal{R}_{\text{eff}}(0) < 1$, and

$$92 \quad \lim_{p \rightarrow 1^-} \mathcal{R}_{\text{eff}}(0) = 0. \quad (\text{S9})$$

93 To prove that $\pi_1 \leq \alpha$, we will show that

$$94 \quad \pi_p \leq \frac{\alpha}{1 - \mathcal{R}_{\text{eff}}(0)}, \quad \text{for all } p > 1 - \frac{1}{\mathcal{R}_0}. \quad (\text{S10})$$

95 Equations (9b) and (S4c) imply that $\dot{R} = \gamma I = \beta SI - \dot{I}$. Hence,

$$96 \quad R_\infty = \int_0^\infty \dot{R} dt = \int_0^\infty (\beta SI - \dot{I}) dt.$$

97 But $S(t)$ decreases monotonically with t , and $I_\infty = 0$ (Appendix C), so

$$98 \quad R_\infty \leq \beta S(0) \int_0^\infty I dt + I(0) = \mathcal{R}_{\text{eff}}(0) \int_0^\infty \gamma I dt + I(0) = \mathcal{R}_{\text{eff}}(0) R_\infty + \alpha(1-p). \quad (\text{S11})$$

99 Thus,

$$100 \quad (1 - \mathcal{R}_{\text{eff}}(0)) R_\infty \leq \alpha(1-p), \quad (\text{S12})$$

101 and consequently, for any $p > 1 - 1/\mathcal{R}_0$,

$$102 \quad \pi_p = \frac{R_\infty}{1-p} \leq \frac{\alpha}{1 - \mathcal{R}_{\text{eff}}(0)}. \quad (\text{S13})$$

103 In the limit $p \rightarrow 1^-$, Equation (S9) implies $\pi_1 \leq \alpha$, as required.

104 To see that $\pi_1 = \alpha$ for the constant rate vaccination (SEIRV) model (Equation (29)), we
105 need only note that in this case,

$$106 \quad R_\infty = \int_0^\infty \gamma I dt = \int_0^\infty (\beta SI - \dot{I} - \dot{E}) dt \leq \mathcal{R}_0 S(0) \int_0^\infty \gamma I dt + E(0), \quad (\text{S14})$$

107 where $E(0) = \alpha(1-p)$. Thus, Inequality (S11) holds for the SEIRV model as well, and the
108 remainder of the proof that $\pi_1 = \alpha$ is identical to the argument for SIRV models.

109 **E Final size relations, π_p and ψ_p**

110 **E.1 Vaccination rate \propto disease prevalence**

111 **E.1.1 Final size relations**

112 A naïve model in which vaccination is proportional to prevalence is

$$113 \quad \dot{S} = -\beta SI - \phi_{\text{prev}} I \quad (\text{S15a})$$

$$114 \quad \dot{I} = \beta SI - \gamma I \quad (\text{S15b})$$

$$115 \quad \dot{R} = \gamma I \quad (\text{S15c})$$

$$116 \quad \dot{V} = \phi_{\text{prev}} I. \quad (\text{S15d})$$

118 However, Equation (S15a) is not biologically sensible, since if $S = 0$ and $I > 0$ it follows that
 119 $\dot{S} < 0$ and so S attains negative values. Thus, a more realistic model is obtained by replacing
 120 the vaccination rate $\phi_{\text{prev}} I$ with $\phi_{\text{prev}} f(S) I$, where f is a nondecreasing and smooth “cutoff
 121 function” such that $f(S) = 1$ except for $0 \leq S < \delta$, and $f(0) = 0$. Thus, Equation (S15a) is
 122 replaced by

$$123 \quad \dot{S} = -\beta SI - \phi_{\text{prev}} f(S) I. \quad (\text{S15a}')$$

124 For convenience, we also choose f to be an odd function, *i.e.*, $f(-S) = -f(S)$ (however,
 125 since negative values of S are not biologically feasible and are unattainable by this model
 126 if $S(0) \geq 0$, this has no effect on the dynamics of the model for biologically sensible initial
 127 conditions).

128 As $\delta \rightarrow 0$, \dot{V} approaches $\phi_{\text{prev}} \text{sgn}(S) I$ and Equation (S15a') approaches

$$129 \quad \dot{S} = -\beta SI - \phi_{\text{prev}} \text{sgn}(S) I,$$

130 where

$$131 \quad \text{sgn}(x) = \begin{cases} -1 & \text{if } x < 0, \\ 0 & \text{if } x = 0, \\ 1 & \text{if } x > 0. \end{cases}$$

132 Thus, a more biologically sensible model where vaccination is proportional to prevalence is:

$$133 \quad \dot{S} = -\beta SI - \phi_{\text{prev}} \text{sgn}(S) I \quad (\text{S16a})$$

$$134 \quad \dot{I} = \beta SI - \gamma I \quad (\text{S16b})$$

$$135 \quad \dot{R} = \gamma I \quad (\text{S16c})$$

$$136 \quad \dot{V} = \phi_{\text{prev}} \text{sgn}(S) I. \quad (\text{S16d})$$

138 In the interior of the biologically meaningful domain,

$$139 \quad \Delta = \{(S, I, R, V) | S \geq 0, I \geq 0, R \geq 0, V = 1 - S + I + R\}, \quad (\text{S17})$$

140 the phase portrait for this model is similar to that of the original model and the dynamics
 141 change only as the hyper-plane $S = 0$ is reached. For this reason, we analyze the original

142 model (Equations (S15)) and make the necessary corrections to account for Equation (S16d)
 143 afterwards. We denote state-variable solutions to the original model (Equations (S15)) with
 144 a superscript 1, as in S^1 , *etc.*

145 From Appendix C, we know that solutions of Equations (S15) converge to a disease-free
 146 equilibrium. Thus, we wish to obtain final size relations for this model. We proceed as
 147 follows:

148 From

$$149 \quad \frac{dR^1}{dS^1} = -\frac{\gamma}{\beta S^1 + \phi_{\text{prev}}}, \quad (\text{S18})$$

150 we have

$$151 \quad R_\infty^1 = \frac{\gamma}{\beta} \ln \left(\frac{\beta S(0) + \phi_{\text{prev}}}{\beta S_\infty^1 + \phi_{\text{prev}}} \right), \quad (\text{S19})$$

152 where a subscript ∞ indicates the value of that variable at the end of the epidemic (recall
 153 that $S(0)$ also depends on p). S_∞^1 is obtained by a similar trick.

$$154 \quad \frac{dI^1}{dS^1} = -\frac{\beta S^1 - \gamma}{\beta S^1 + \phi_{\text{prev}}} = -1 + \frac{\phi_{\text{prev}} + \gamma}{\beta S^1 + \phi_{\text{prev}}} \quad (\text{S20})$$

$$155 \quad I(t) - I(0) = S(0) - S^1(t) + \frac{\phi_{\text{prev}} + \gamma}{\beta} \ln \left(\frac{\beta S^1(t) + \phi_{\text{prev}}}{\beta S(0) + \phi_{\text{prev}}} \right). \quad (\text{S21})$$

157 As $t \rightarrow \infty$, we have

$$158 \quad I_\infty - I(0) = S(0) - S_\infty^1 + \frac{\phi_{\text{prev}} + \gamma}{\beta} \ln \left(\frac{\beta S_\infty^1 + \phi_{\text{prev}}}{\beta S(0) + \phi_{\text{prev}}} \right). \quad (\text{S22})$$

159 Since $I_\infty^1 = 0$ and $S(0) + I(0) = 1 - V(0) = 1 - p$,

$$160 \quad S_\infty^1 = (1 - p) + \frac{\phi_{\text{prev}} + \gamma}{\beta} \ln \left(\frac{\beta S_\infty^1 + \phi_{\text{prev}}}{\beta S(0) + \phi_{\text{prev}}} \right) = (1 - p) - \frac{\phi_{\text{prev}} + \gamma}{\gamma} R_\infty^1. \quad (\text{S23})$$

161 Let

$$162 \quad w(x) = (1 - p) + \frac{\phi_{\text{prev}} + \gamma}{\beta} \ln \left(\frac{\beta x + \phi_{\text{prev}}}{\beta(1 - p)(1 - \alpha) + \phi_{\text{prev}}} \right). \quad (\text{S24})$$

163 We seek solutions to $S_\infty^1 = w(S_\infty^1)$ in the range $S_\infty^1 \in [0, 1]$. We note that no solutions
 164 of Equation (S15) cross the S -nullcline, $S = -\phi_{\text{prev}}/\beta$, and so solutions to Equation (S23)
 165 are in the range $[-\phi_{\text{prev}}/\beta, 1]$.

166 It is possible to use Equation (S23) to find which initial coverage causes solutions of Equa-
 167 tion (S15) to hit the $S = 0$ hyper-plane, as they will be those for which $S_\infty^1 \leq 0$. As long as
 168 this does not occur, Equations (S19), (S22) and (S23) are valid also for the modified system
 169 (Equations (S16)).

170 From Equation (S23), we have

$$171 \quad S_\infty^1 = -\frac{1}{\beta} \left(\phi_{\text{prev}} + (\gamma + \phi_{\text{prev}}) W_i \left(-\frac{\beta S(0) + \phi_{\text{prev}}}{\gamma + \phi_{\text{prev}}} e^{-\frac{\beta(1-p) + \phi_{\text{prev}}}{\gamma + \phi_{\text{prev}}}} \right) \right),$$

172 where $i = 0$ or -1 specifies the branch of the Lambert W function. To determine which
173 branch of W gives the correct final size, observe that $S_\infty^1 < \gamma/\beta$ for any initial condition (in
174 the biologically meaningful domain). This follows from the following argument: If $S(0) \leq$
175 γ/β , since S is non-increasing and $\dot{S} < 0$ at time $t = 0$ (since also $I(0) > 0$), we are done.
176 If $S(0) > \gamma/\beta$, we note that I is increasing for any S such that $S > \gamma/\beta$. But we have
177 seen that $I_\infty = 0$. Thus, at some point in time, $S < \gamma/\beta$, and since S is non-increasing,
178 we have $S_\infty^1 < \gamma/\beta$. From this we see that it is necessary to use the principal branch, W_0
179 (rather than W_{-1} , which satisfies $W_{-1}(x) \leq -1$ for all x in its domain of definition, that is
180 $(-1/e, 0)$). Thus,

$$181 \quad S_\infty^1 = -\frac{1}{\beta} \left(\phi_{\text{prev}} + (\gamma + \phi_{\text{prev}}) W_0 \left(-\frac{\beta S(0) + \phi_{\text{prev}}}{\gamma + \phi_{\text{prev}}} e^{-\frac{\beta(1-p) + \phi_{\text{prev}}}{\gamma + \phi_{\text{prev}}}} \right) \right). \quad (\text{S25})$$

182 For convenience, we rewrite Equations (S19), (S22) and (S23) to give:

$$183 \quad \ln \left(\frac{\beta S(0) + \phi_{\text{prev}}}{\beta S_\infty^1 + \phi_{\text{prev}}} \right) = \frac{\beta}{\phi_{\text{prev}} + \gamma} (1 - p - S_\infty^1), \quad (\text{S26})$$

184 and

$$185 \quad R_\infty^1 = \frac{\gamma}{\beta} \ln \left(\frac{\beta S(0) + \phi_{\text{prev}}}{\beta S_\infty^1 + \phi_{\text{prev}}} \right) = \frac{\gamma}{\phi_{\text{prev}} + \gamma} (1 - p - S_\infty^1). \quad (\text{S27})$$

186 Thus,

$$187 \quad \begin{aligned} V_\infty^1 &= 1 - R_\infty^1 - S_\infty^1 \\ &= 1 - \frac{\gamma}{\phi_{\text{prev}} + \gamma} (1 - p - S_\infty^1) - S_\infty^1 \\ &= \frac{\phi_{\text{prev}}}{\gamma + \phi_{\text{prev}}} (1 - S_\infty^1) + \frac{\gamma}{\gamma + \phi_{\text{prev}}} p. \end{aligned} \quad (\text{S28})$$

191 To see when we can use $S_\infty = S_\infty^1$, $R_\infty = R_\infty^1$ and $V_\infty = V_\infty^1$, it is necessary to find
192 when Equation (S25) yields a negative S_∞^1 . First, we evaluate how S_∞ changes with p . To
193 find $\partial_p S_\infty^1$, apply $\partial_p := \frac{\partial}{\partial p}$ to Equation (S23), to get (recall that $S(0) = (1 - \alpha)(1 - p)$, so
194 $\partial_p S(0) = -(1 - \alpha)$)

$$195 \quad \partial_p S_\infty^1 = \frac{\gamma + \phi_{\text{prev}}}{\beta} \left[\frac{\beta(1 - \alpha)}{\beta S(0) + \phi_{\text{prev}}} + \frac{\beta \partial_p S_\infty^1}{\beta S_\infty^1 + \phi_{\text{prev}}} \right] - 1 \quad (\text{S29})$$

$$196 \quad \left(1 - \frac{\gamma + \phi_{\text{prev}}}{\beta S_\infty^1 + \phi_{\text{prev}}} \right) \partial_p S_\infty^1 = \frac{(\gamma + \phi_{\text{prev}})(1 - \alpha)}{\beta S(0) + \phi_{\text{prev}}} - 1 \quad (\text{S30})$$

$$197 \quad \partial_p S_\infty^1 = \left(\frac{(\gamma + \phi_{\text{prev}})(1 - \alpha)}{\beta S(0) + \phi_{\text{prev}}} - 1 \right) / \left(1 - \frac{\gamma + \phi_{\text{prev}}}{\beta S_\infty^1 + \phi_{\text{prev}}} \right) \quad (\text{S31})$$

199 Since $S_\infty^1 < \gamma/\beta$, it follows that $1 - (\gamma + \phi_{\text{prev}})/(\beta S_\infty^1 + \phi_{\text{prev}}) < 0$. Thus,

$$200 \quad \text{sgn}(\partial_p S_\infty^1) = \text{sgn}[\beta S(0) + \phi_{\text{prev}} - (\gamma + \phi_{\text{prev}})(1 - \alpha)], \quad (\text{S32})$$

201 and since $S(0) = (1 - p)(1 - \alpha)$, we have:

$$202 \quad \text{sgn } \partial_p S_\infty^1 = \begin{cases} 1 & \text{if } p_m > p \\ 0 & \text{if } p_m = p, \\ -1 & \text{if } p_m < p, \end{cases} \quad (\text{S33})$$

203 where the local maximum is attained at

$$204 \quad p = p_m := 1 + \frac{\phi_{\text{prev}} - (\gamma + \phi_{\text{prev}})(1 - \alpha)}{\beta(1 - \alpha)} = 1 + \frac{\alpha\phi_{\text{prev}} - \gamma(1 - \alpha)}{\beta(1 - \alpha)}. \quad (\text{S34})$$

205 Observe that $p_m \in [0, 1] \iff \alpha\phi_{\text{prev}} \leq \gamma(1 - \alpha)$ and $\alpha\phi_{\text{prev}} \geq (\gamma - \beta)(1 - \alpha)$. However,
 206 the second condition, which is necessary to ensure $p_m \geq 0$, is trivially satisfied whenever
 207 $\beta/\gamma = \mathcal{R}_0 \geq 1$. Thus, if $\mathcal{R}_0 \geq 1$, then $p_m \in [0, 1]$ iff

$$208 \quad \alpha\phi_{\text{prev}} \leq \gamma(1 - \alpha). \quad (\text{S35})$$

209 The maximum value of S_∞^1 is thus:

$$210 \quad \max_{p \in [0, 1]} S_\infty^1 = -\frac{1}{\beta} \left(\phi_{\text{prev}} + (\gamma + \phi_{\text{prev}}) W_0 \left(-\frac{\beta S(0) + \phi_{\text{prev}}}{\gamma + \phi_{\text{prev}}} e^{-\frac{\phi_{\text{prev}} - \gamma(1 - \alpha)}{(\gamma + \phi_{\text{prev}})(1 - \alpha)}} \right) \right). \quad (\text{S36})$$

211 Now solving for $S_\infty^1 = 0$ using Equation (S25), we have

$$212 \quad p_0(i) = 1 + \frac{\phi_{\text{prev}}}{\beta(1 - \alpha)} + \frac{\gamma + \phi_{\text{prev}}}{\beta} W_i \left(-\frac{\phi_{\text{prev}}}{(1 - \alpha)(\gamma + \phi_{\text{prev}})} e^{-\frac{\phi_{\text{prev}}}{(1 - \alpha)(\gamma + \phi_{\text{prev}})}} \right) \quad (\text{S37})$$

213 for $i = 0$ or -1 . Note that for $\phi_{\text{prev}} > 0$, Equation (S37) gives two values for p_0 ; we cannot
 214 simply cancel out the operation of W with xe^x , since in this case $x = -\frac{\phi_{\text{prev}}}{(1 - \alpha)(\phi_{\text{prev}} + \gamma)} < 0$
 215 and W is not univalued for negative arguments. Instead, we have two possibilities for
 216 $W \left(\frac{\phi_{\text{prev}}}{(\alpha - 1)(\gamma + \phi_{\text{prev}})} e^{\frac{\phi_{\text{prev}}}{(\alpha - 1)(\gamma + \phi_{\text{prev}})}} \right)$ corresponding to the two branches, W_0 and W_{-1} .

217 If $\alpha\phi_{\text{prev}} < (1 - \alpha)\gamma$, then $W_0 \left(-\frac{\phi_{\text{prev}}}{(1 - \alpha)(\gamma + \phi_{\text{prev}})} e^{-\frac{\phi_{\text{prev}}}{(1 - \alpha)(\gamma + \phi_{\text{prev}})}} \right) = -\frac{\phi_{\text{prev}}}{(1 - \alpha)(\gamma + \phi_{\text{prev}})}$, which
 218 gives $p_0(0) = 1$. If $\alpha\phi_{\text{prev}} \geq (1 - \alpha)\gamma$ then similarly $p_0(-1) = 1$. This is in agreement with
 219 the fact that if $p = 1$, $S(0) = 0$, and so $S_\infty^1 = 0$.

220 There are now three cases, which we express as two main cases, the second of which has
 221 two subcases:

- 222 • if $p_m \geq 1$ (which happens iff $\alpha\phi_{\text{prev}} \geq (1 - \alpha)\gamma$), then $S_\infty^1 \leq 0 \quad \forall p \in [0, 1]$. This
 223 follows since if $p_m \geq 1$ then $p_0(-1) = 1 \leq p_m \leq p_0(0)$. Thus because S_∞^1 is increasing
 224 for $p < p_m$, so for $p \in [0, 1]$, $S_\infty^1 \leq S_\infty^1|_{p=1} = 0$. In this case, S_∞ is not given by S_∞^1 , but
 225 is simply $S_\infty = 0$.
- 226 • If $p_m \in (0, 1)$ then $S_\infty^1|_{p_m} > S_\infty^1|_{p=1} = 0$ and $p_0(-1) < p_m$.

227 – If $0 \leq p_0(-1)$ (and $p_0(-1) < p_m < 1$) then

$$228 \quad S_\infty = \begin{cases} 0 & \text{if } p < p_0(-1), \\ S_\infty^1 & \text{if } p_0(-1) \leq p. \end{cases} \quad (\text{S38})$$

229 – If $p_0(-1) \leq 0$ then $S_\infty^1 \geq 0 \quad \forall p \in [0, 1]$ and so $S_\infty = S_\infty^1$ for any $p \in [0, 1]$.

230 In all but the very last sub-case, it is also necessary to adjust our formulae for the final
 231 sizes of the removed and vaccinated compartments, R_∞ and V_∞ , for the values of p for which
 232 $S_\infty = 0$. Qualitatively, this adjustment is necessary because, when $\delta \rightarrow 0^+$, if a solution
 233 reaches $S = 0$ in finite time, S remains 0, while I decays exponentially to 0. However, the
 234 solutions of Equations (S15) are only identical to those of Equations (S16) so long as $S > 0$.
 235 Moreover, once $S = 0$, V remains constant and all the infectives move into the recovered
 236 compartment, which is not the case for solutions of Equations (S15).

To find formulas for R_∞ and V_∞ , fix $p \in [0, 1)$ and let t_0 be the first time at which no
 susceptibles remain ($S(t_0) = S^1(t_0) = 0$). Then Equations (S15) are valid for any $t < t_0$. We
 now have

$$R(t) = \frac{\gamma}{\beta} \ln \left(\frac{\beta S(0) + \phi_{\text{prev}}}{\beta S(t) + \phi_{\text{prev}}} \right)$$

$$I(t) = I(0) + S(0) - S(t) + \frac{\phi_{\text{prev}} + \gamma}{\beta} \ln \left(\frac{\beta S(t) + \phi_{\text{prev}}}{\beta S(0) + \phi_{\text{prev}}} \right),$$

in a manner analogous to Equation (S19) and Equation (S22). These equations depend on
 $S = S(t)$ in a way that is continuous at $S = 0$, and so taking $t \rightarrow t_0$ is equivalent to taking
 $S \rightarrow 0$:

$$R(t_0) = \frac{\gamma}{\beta} \ln \left(\frac{\beta S(0) + \phi_{\text{prev}}}{\phi_{\text{prev}}} \right) \quad (\text{S39})$$

$$I(t_0) = I(0) + S(0) + \frac{\phi_{\text{prev}} + \gamma}{\beta} \ln \left(\frac{\phi_{\text{prev}}}{\beta S(0) + \phi_{\text{prev}}} \right). \quad (\text{S40})$$

237 When $S = 0$, I decays exponentially to 0, until all the infectives present at t_0 transition into
 238 the removed class, R . Thus, we have

$$239 \quad R_\infty = R(t_0) + I(t_0) = 1 - p - \frac{\phi_{\text{prev}}}{\beta} \ln \left(\frac{\beta}{\phi_{\text{prev}}} S(0) + 1 \right). \quad (\text{S41})$$

240 Next, we know that once $S = 0$, V does not change either, since there are no more
 241 susceptibles to be vaccinated. Thus,

$$242 \quad V_\infty = V(t_0) = 1 - R(t_0) - I(t_0) = 1 - R_\infty = p + \frac{\phi_{\text{prev}}}{\beta} \ln \left(\frac{\beta}{\phi_{\text{prev}}} S(0) + 1 \right) \quad (\text{S42})$$




Figure S2: The proportion of individuals remaining susceptible at the end of the epidemic, S_∞ , as a function of the proportion of the population vaccinated preemptively, p , for the model in which vaccination rate is proportional to prevalence. The line $y(p) = (1 - \alpha)(1 - p)$ is overlaid in red. We take the proportion of susceptibles initially infected, α , to be the estimated value in the left panel ($\alpha = 1.72 \times 10^{-5}$) and a much larger value for comparison in the right panel ($\alpha = 0.1$); the remaining model parameters are as in Table 1. See § E.1.2.

243 To summarize our results so far,

$$244 \quad S_\infty = \begin{cases} 0 & \text{if } p < p_0 \text{ or } 1 \leq p_m, \\ S_\infty^1 & \text{if } p_0 \leq p \leq 1, \end{cases} \quad (\text{S43a})$$

$$245 \quad R_\infty = \begin{cases} 1 - p - \frac{\phi_{\text{prev}}}{\beta} \ln \left(\frac{\beta}{\phi_{\text{prev}}} S(0) + 1 \right) & \text{if } p < p_0 \text{ or } 1 \leq p_m, \\ \frac{\gamma}{\gamma + \phi_{\text{prev}}} (1 - p - S_\infty^1) & \text{if } p_0 \leq p \leq 1, \end{cases} \quad (\text{S43b})$$

$$246 \quad V_\infty = \begin{cases} p + \frac{\phi_{\text{prev}}}{\beta} \ln \left(\frac{\beta}{\phi_{\text{prev}}} S(0) + 1 \right) & \text{if } p < p_0 \text{ or } 1 \leq p_m, \\ \frac{1}{\gamma + \phi_{\text{prev}}} (\phi_{\text{prev}}(1 - S_\infty^1) + \gamma p) & \text{if } p_0 \leq p \leq 1, \end{cases} \quad (\text{S43c})$$

247 where S_∞^1 is given by Equation (S25).

249 E.1.2 Qualitative behaviour of S_∞^1 for high vaccine coverage

250 Qualitatively, observe that for high values of p , $S_\infty^1 \approx (1 - \alpha)(1 - p)$ (see Figure S2). This
 251 is because when $p > p_d := 1 - \frac{\gamma}{\beta(1-\alpha)}$ then $S(0) < \gamma/\beta$, in which case I decays to 0
 252 monotonically (the subscript “ d ” denotes decay of I). Now in this case, S decreases at least

253 as fast as it does in the vaccination-less *SIR* model, and so I decreases at least as fast as in
 254 the vaccination-less case too. Because S is monotonically decreasing, I decays faster than
 255 $I(0)e^{(\beta S(0) - \gamma)t}$. Because I decays at least exponentially, S hardly changes over the course of
 256 the epidemic, and so we see an approximately linear decay in $S_\infty^1 \approx S(0) = (1 - \alpha)(1 - p)$
 257 to 0 as we increase p . This last phenomenon is related to the herd immunity effect in the
 258 standard *SIR* model: when the entire population is susceptible, in the absence of post-
 259 outbreak vaccination ($\phi_{\text{prev}} = 0$), the critical vaccine coverage which stops the epidemic
 260 from taking off is $p = 1 - \frac{1}{\mathcal{R}_{\text{eff}}} = 1 - \frac{\gamma}{\beta(1 - \alpha)}$. But note, moreover, that the maximal value of
 261 S_∞^1 is not attained at p_d . Rather, we see that

$$262 \quad p_m - p_d = \frac{\alpha(\phi_{\text{prev}} + \gamma)}{\beta(1 - \alpha)} > 0. \quad (\text{S44})$$

263 This is because even when $p > p_d$ and I immediately decays to 0 (starting at $t = 0$), there
 264 are still some susceptibles converted into vaccinated individuals, due to $I > 0$. This number
 265 of susceptibles lost to vaccination decreases as p is increased, since this decreases $I(0)$ as
 266 well. Thus, for $p > p_d$, initially S_∞ increases with p . Only when $p > p_m$ does the decrease
 267 in S_∞ due to more susceptibles being vaccinated pre-emptively take over.

268 **E.1.3 π_p decreases and ψ_p increases with p when $p_m \geq 1$**

269 Consider the first case above: if $p_m \geq 1$ (see Equation (S43)), then $S_\infty = 0$ and we have

$$270 \quad V_\infty = p + \frac{\phi_{\text{prev}}}{\beta} \ln \left(\frac{\beta}{\phi_{\text{prev}}} S(0) + 1 \right)$$

$$271 \quad \psi_p = \frac{\phi_{\text{prev}}}{\beta(1 - p)} \ln \left(\frac{\beta}{\phi_{\text{prev}}} S(0) + 1 \right).$$

273 Note that since $S_\infty = 0$, $\pi_p = 1 - \psi_p$. Letting $x = \frac{\beta}{\phi_{\text{prev}}} S(0) = \frac{\beta(1 - \alpha)}{\phi_{\text{prev}}}(1 - p)$,

$$274 \quad \psi(x) = \psi_{p(x)} = (1 - \alpha) \frac{\ln(x + 1)}{x},$$

275 SO

$$276 \quad \partial_p \psi_p = - \frac{\beta(1 - \alpha)}{\phi_{\text{prev}}} \frac{\partial \psi(x)}{\partial x} = - \frac{\beta(1 - \alpha)}{\phi_{\text{prev}}} \frac{x - (1 + x) \ln(1 + x)}{x^2(x + 1)}$$

$$277 \quad = - \frac{\beta(1 - \alpha)}{\phi_{\text{prev}}} \frac{\frac{\beta}{\phi_{\text{prev}}} S(0) - \left(\frac{\beta}{\phi_{\text{prev}}} S(0) + 1\right) \ln\left(\frac{\beta}{\phi_{\text{prev}}} S(0) + 1\right)}{\left(\frac{\beta}{\phi_{\text{prev}}} S(0)\right)^2 \left(\frac{\beta}{\phi_{\text{prev}}} S(0) + 1\right)}$$

$$278 \quad = - \frac{\phi_{\text{prev}}(1 - \alpha)}{\beta} \frac{\beta S(0) - (\beta S(0) + \phi_{\text{prev}}) \ln\left(\frac{\beta}{\phi_{\text{prev}}} S(0) + 1\right)}{(S(0))^2 (\beta S(0) + \phi_{\text{prev}})}. \quad (\text{S45})$$

280 Since $x < (x + 1) \ln(1 + x) \quad \forall x > -1$, it follows that for $p \in [0, 1)$, $\partial_p \psi_p > 0$ and consequently,
 281 $\partial_p \pi_p < 0$.

282 E.1.4 Behaviour of $p_0(-1)$ as $\phi_{\text{prev}} \rightarrow 0^+$

283 Using Equation (S37), one can show that $\lim_{\phi_{\text{prev}} \rightarrow 0^+} p_0(-1) = -\infty$. This follows because
 284 $\lim_{\phi_{\text{prev}} \rightarrow 0^-} W_{-1}(x) = -\infty$. But because $p_0(-1) \rightarrow 1$ as $\phi_{\text{prev}} \rightarrow \frac{\gamma(1-\alpha)}{\alpha}$, it follows that there
 285 is some value of ϕ_{prev} for which $p_0(-1) = 0$. This ϕ_{prev} can be found from Equation (S37),
 286 but the formula is not needed here.

287 However, we also note that $\phi_{\text{prev}} < \frac{\gamma(1-\alpha)}{\alpha}$ (along with $\mathcal{R}_0 > 1$) implies $p_m \in (0, 1)$, so for
 288 small ϕ_{prev} , we have $p_m \in (0, 1)$ and $p_0(-1) < 0$ (in fact, as $\phi_{\text{prev}} \rightarrow 0^+$, $p_m \rightarrow 1 - 1/\mathcal{R}_0$).
 289 Thus, for $\phi_{\text{prev}} \rightarrow 0^+$ (small enough such that $p_0(-1) < 0$), $S_\infty^1 > 0$ for $p \in [0, 1)$.

290 E.2 Vaccination rate \propto incidence

291 The model equations are

$$292 \quad \dot{S} = -\beta SI - \phi_{\text{inc}} SI \quad (\text{S46a})$$

$$293 \quad \dot{I} = \beta SI - \gamma I \quad (\text{S46b})$$

$$294 \quad \dot{R} = \gamma I \quad (\text{S46c})$$

$$295 \quad \dot{V} = \phi_{\text{inc}} SI. \quad (\text{S46d})$$

297 Finding the final sizes for this model is somewhat similar to when vaccination is propor-
 298 tional to prevalence:

$$299 \quad \frac{dI}{dS} = -\frac{\beta SI - \gamma I}{(\beta + \phi_{\text{inc}})SI} = \frac{\gamma}{\beta + \phi_{\text{inc}}} \frac{1}{S} - \frac{\beta}{\beta + \phi_{\text{inc}}} \quad (\text{S47})$$

$$300 \quad I(t) - I(0) = \frac{\gamma}{\beta + \phi_{\text{inc}}} \ln \left(\frac{S(t)}{S(0)} \right) - \frac{\beta}{\beta + \phi_{\text{inc}}} (S(t) - S(0)) \quad (\text{S48})$$

$$301 \quad -(1-p)\alpha = \frac{\gamma}{\beta + \phi_{\text{inc}}} \ln \left(\frac{S_\infty}{(1-p)(1-\alpha)} \right) - \frac{\beta}{\beta + \phi_{\text{inc}}} (S_\infty - (1-p)(1-\alpha)) \quad (\text{S49})$$

$$302 \quad S_\infty = \frac{\gamma}{\beta} \ln \left(\frac{S_\infty}{(1-p)(1-\alpha)} \right) + \left(1 + \frac{\phi_{\text{inc}}}{\beta} \alpha \right) (1-p), \quad (\text{S50})$$

304 where Equation (S49) is obtained by taking $t \rightarrow \infty$ in Equation (S48). The solution of Equa-
 305 tion (S50) is given explicitly by

$$306 \quad S_\infty = -\frac{\gamma}{\beta} W_0 \left(-\frac{\beta(1-p)(1-\alpha)}{\gamma} e^{-\frac{\beta + \phi_{\text{inc}} \alpha}{\gamma} (1-p)} \right), \quad (\text{S51})$$

307 where we take the principle branch of the Lambert function, W_0 , because solutions are in the
 308 range $S_\infty \in [0, \gamma/\beta]$ (see Equation (S5) in Appendix C). Note also that $S_\infty > 0$ *iff* $S(0) > 0$
 309 (that is, $p < 1$ and $\alpha < 1$). Thus, $S = 0$ is not attainable in scenarios of interest here.

310 To find V_∞ , we proceed similarly:

$$311 \quad \dot{V} = -\frac{\phi_{\text{inc}}}{\beta + \phi_{\text{inc}}} \dot{S} \quad (\text{S52})$$

$$312 \quad V_\infty - V(0) = -\frac{\phi_{\text{inc}}}{\beta + \phi_{\text{inc}}}(S_\infty - S(0)) \quad (\text{S53})$$

$$313 \quad V_\infty - p = \frac{\phi_{\text{inc}}}{\beta + \phi_{\text{inc}}}((1-p)(1-\alpha) - S_\infty). \quad (\text{S54})$$

314

315 At the end of the epidemic, $1 = R_\infty + S_\infty + V_\infty$, hence,

$$316 \quad R_\infty = 1 - p + \frac{\phi_{\text{inc}}}{\beta + \phi_{\text{inc}}}(S_\infty - (1-p)(1-\alpha)) - S_\infty$$

$$317 \quad = 1 - p - \frac{\phi_{\text{inc}}(1-p)(1-\alpha) + \beta S_\infty}{\beta + \phi_{\text{inc}}}. \quad (\text{S55})$$

318

319 Thus,

$$320 \quad \pi_p = \frac{R_\infty}{1-p} = 1 - \frac{\phi_{\text{inc}}}{\beta + \phi_{\text{inc}}}(1-\alpha) - \frac{\beta S_\infty}{(\beta + \phi_{\text{inc}})(1-p)} \quad (\text{S56a})$$

$$321 \quad \psi_p = \frac{V_\infty - p}{1-p} = \frac{\phi_{\text{inc}}}{\beta + \phi_{\text{inc}}} \left((1-\alpha) - \frac{S_\infty}{1-p} \right). \quad (\text{S56b})$$

322

323 Note that whenever $S(0) > 0$, there are susceptible individuals left at the end of the epidemic,
324 and so $\pi_p \neq 1 - \psi_p$.

325 Using $\partial_p \left(\frac{S_\infty}{1-p} \right) = \frac{(1-p)\partial_p S_\infty + S_\infty}{(1-p)^2}$ we have

$$326 \quad \partial_p S_\infty = \frac{\gamma}{\beta} \left(\frac{\partial_p S_\infty}{S_\infty} + \frac{1}{1-p} \right) - \left(1 + \frac{\phi_{\text{inc}}}{\beta} \alpha \right)$$

$$327 \quad \frac{\beta S_\infty - \gamma}{\beta S_\infty} \partial_p S_\infty = \frac{\gamma}{\beta(1-p)} - 1 - \frac{\phi_{\text{inc}}}{\beta} \alpha$$

$$328 \quad \partial_p S_\infty = \frac{\gamma}{1-p} \frac{S_\infty}{\beta S_\infty - \gamma} - (\phi_{\text{inc}} \alpha + \beta) \frac{S_\infty}{\beta S_\infty - \gamma}$$

$$329 \quad = \frac{\gamma - (\phi_{\text{inc}} \alpha + \beta)(1-p)}{(1-p)} \frac{S_\infty}{\beta S_\infty - \gamma}$$

$$330 \quad (1-p)\partial_p S_\infty + S_\infty = \frac{(\beta S_\infty - (1-p)(\beta + \phi_{\text{inc}} \alpha))S_\infty}{\beta S_\infty - \gamma}$$

$$331 \quad \partial_p \left(\frac{S_\infty}{1-p} \right) = \frac{(\beta S_\infty - (1-p)(\beta + \phi_{\text{inc}} \alpha))S_\infty}{\beta S_\infty - \gamma} \frac{1}{(1-p)^2}. \quad (\text{S57})$$

332

333 From Equation (S50)

$$334 \quad \beta S_\infty - (1-p)(\beta + \phi_{\text{inc}} \alpha) = \gamma \ln \left(\frac{S_\infty}{S(0)} \right) < 0 \quad (\text{S58})$$

335 because $S_\infty < S(0)$ (unless $S(0)$ or $I(0)$ is 0, in which case an outbreak cannot take place).

336 Thus $\partial_p \left(\frac{S_\infty}{1-p} \right) > 0$ and so

$$337 \quad \partial_p \pi_p < 0 \quad (\text{S59})$$

$$338 \quad \partial_p \psi_p < 0. \quad (\text{S60})$$

340 Note also that S_∞ attains a local maximum (in p) at

$$341 \quad p_m = 1 - \gamma / (\phi_{\text{inc}} \alpha + \beta). \quad (\text{S61})$$

342 $p_m < 0$ for $\gamma > \phi_{\text{inc}} \alpha + \beta$, in which case S_∞ decreases with p on the interval $[0, 1]$. This can
 343 only happen when $\gamma > \beta$, that is when $\mathcal{R}_0 < 1$, implying that for any disease which can
 344 spread in the population (with no vaccination), pre-emptive vaccination initially raises, then
 345 lowers the proportion of susceptibles remaining at the end of the epidemic. The maximum
 346 level of remaining susceptibles is

$$347 \quad S_\infty|_{p=p_m} = -\frac{\gamma}{\beta} W_0 \left(-\frac{\beta(1-\alpha)}{(\phi_{\text{inc}} \alpha + \beta)e} \right). \quad (\text{S62})$$

348 However, R_∞ is more informative, since susceptibles can be depleted by either infection or
 349 vaccination, and so fewer remaining susceptibles does not necessarily imply a larger epidemic,
 350 nor does it imply that more individuals were vaccinated. However,

$$351 \quad \partial_p R_\infty = \partial_p ((1-p)\pi_p) = -\pi_p + (1-p)\partial_p \pi_p < 0, \quad (\text{S63})$$

353 which shows that increasing pre-emptive vaccine coverage decreases the size of the epidemic,
 354 as expected.

355 **E.3 Vaccination rate \propto proportion still susceptible**

356 The model equations are

$$357 \quad \dot{S} = -\beta SI - \phi_{\text{susc}} S \quad (\text{S64a})$$

$$358 \quad \dot{I} = \beta SI - \gamma I \quad (\text{S64b})$$

$$359 \quad \dot{R} = \gamma I \quad (\text{S64c})$$

$$360 \quad \dot{V} = \phi_{\text{susc}} S. \quad (\text{S64d})$$

362 In this case, a similar strategy to the one we employed for the case where vaccination is
 363 proportional to prevalence doesn't quite work. Calculating $S_\infty(I(0))$ is not enough, since we
 364 do not know how the remainder of the population is partitioned between the removed and
 365 vaccinated classes at the end of the epidemic. The following calculations are also helpful but
 366 insufficient:

$$367 \quad \frac{\dot{I}}{I} = \beta S - \gamma = \frac{\beta}{\phi_{\text{susc}}} \dot{V} - \gamma \quad (\text{S65})$$

$$368 \quad \frac{\dot{S}}{S} = -\beta I - \phi_{\text{susc}} = -\frac{\beta}{\gamma} \dot{R} - \phi_{\text{susc}}, \quad (\text{S66})$$

369

370 SO

$$371 \quad \ln \left(\frac{I(t)}{I(0)} \right) = \frac{\beta}{\phi_{\text{susc}}} (V(t) - V(0)) - \gamma t \quad (\text{S67})$$

$$372 \quad \ln \left(\frac{S(t)}{S(0)} \right) = -\frac{\beta}{\gamma} (R(t) - R(0)) - \phi_{\text{susc}} t. \quad (\text{S68})$$

373

374 However, it is not possible to extract V_∞ and R_∞ from here because phase-portrait arguments
 375 show that I and S tend to 0 as $t \rightarrow \infty$ (this is also implied by Equations (S67) and (S68)),
 376 thus both sides of these equations diverge as $t \rightarrow \infty$.

377 Nonetheless, a similar method to the one employed in Appendix E.2 yields a relation
 378 between $S(t)$ and $I(t)$ from which, using the previous relations, a relation between $R(t)$ and
 379 $V(t)$ can be obtained. These will not diverge as $t \rightarrow \infty$ (they are bounded), so any divergent
 380 components must cancel out.

$$381 \quad \frac{dS}{dI} = -\frac{(\beta I + \phi_{\text{susc}})S}{(\beta S - \gamma)I} \quad (\text{S69})$$

$$382 \quad \frac{(\beta S - \gamma)}{S} dS = -\frac{(\beta I + \phi_{\text{susc}})}{I} dI \quad (\text{S70})$$

$$383 \quad \left(\beta - \frac{\gamma}{S}\right) dS = -\left(\beta + \frac{\phi_{\text{susc}}}{I}\right) dI \quad (\text{S71})$$

$$384 \quad \beta(S(t) - S(0)) - \gamma \ln \left(\frac{S(t)}{S(0)} \right) = -\beta(I(t) - I(0)) - \phi_{\text{susc}} \ln \left(\frac{I(t)}{I(0)} \right) \quad (\text{S72})$$

385

386 from which we get:

$$387 \quad S(t) = -\frac{\gamma}{\beta} W \left(-\frac{\beta}{\gamma} e^{\frac{\beta}{\gamma}(I(t)-I(0)-S(0))} \left(\frac{I(t)}{I(0)} \right)^{\frac{\phi_{\text{susc}}}{\gamma}} S(0) \right). \quad (\text{S73})$$

388 By taking the limit $t \rightarrow \infty$ in Equation (S73), we see that $S_\infty = 0$, and consequently,
 389 $\psi_p = 1 - \pi_p$.

390 We now determine under which conditions each of the two branches of the Lambert W
 391 function is used in Equation (S73). First, note that

$$392 \quad S(I) = -\frac{\gamma}{\beta} W(z(p))$$

$$393 \quad z(I) = -\frac{\beta}{\gamma} e^{\frac{\beta}{\gamma}(I(t)-I(0)-S(0))} \left(\frac{I(t)}{I(0)} \right)^{\frac{\phi_{\text{susc}}}{\gamma}} S(0)$$

$$394 \quad z(I(0)) = -\frac{\beta}{\gamma} S(0) e^{-\frac{\beta}{\gamma} S(0)}.$$

395

396 For $I = I(0)$, we expect to get $W(-\frac{\beta}{\gamma} S(0)) = -\frac{\beta}{\gamma} S(0)$ so that $S(I = I(0)) = S(t = 0)$.
 397 We know that $S(0) > \frac{\gamma}{\beta}$ which implies that for $t = 0$, we must use W_1 . Because $S(t)$
 398 monotonically decreases to 0 as $t \rightarrow 0$, we know that the branch W_1 is used until the peak

399 prevalence is attained (at which time $S = \gamma/\beta$), and then the principal branch W_0 is used.
 400 Now,

$$\begin{aligned}
 401 \quad \frac{dR}{dI} &= \frac{\gamma I}{\beta S I - \gamma I} \\
 402 \quad &= \frac{\gamma}{\beta S - \gamma} \\
 403 \quad &= \frac{-1}{W \left(-\frac{\beta}{\gamma} e^{\frac{\beta}{\gamma}(I-I(0)-S(0))} \left(\frac{I}{I(0)}\right)^{\frac{\phi_{\text{susc}}}{\gamma}} S(0) \right) + 1}. \tag{S74}
 \end{aligned}$$

405 This can be integrated, to give

$$406 \quad R_\infty = R(0) + \int_{I(0)}^{I_\infty} \frac{\gamma}{\beta S - \gamma} dI \tag{S75}$$

$$\begin{aligned}
 407 \quad &= R(0) + \int_{I(0)}^{I_\infty} \frac{-1}{W_i \left(-\frac{\beta}{\gamma} e^{\frac{\beta}{\gamma}(I-I(0)-S(0))} \left(\frac{I}{I(0)}\right)^{\frac{\phi_{\text{susc}}}{\gamma}} S(0) \right) + 1} dI \\
 408 \quad &= \int_0^{I(0)} \frac{1}{W_i \left(-\frac{\beta}{\gamma} e^{\frac{\beta}{\gamma}(I-I(0)-S(0))} \left(\frac{I}{I(0)}\right)^{\frac{\phi_{\text{susc}}}{\gamma}} S(0) \right) + 1} dI, \tag{S76} \\
 409
 \end{aligned}$$

410 where the appropriate branch of W_i is determined as above, as the integration variable I is
 411 varied. Note, however, that the integral in Equation (S75) is improper, because the integral
 412 diverges at the peak prevalence (when $S = \gamma/\beta$).

413 **E.4 Instantaneous vaccination of a proportion ϕ_{inst} of the popula-** 414 **tion**

415 In this case, the disease progresses according to the standard *SIR* model,

$$416 \quad \dot{S} = -\beta S I, \tag{S77a}$$

$$417 \quad \dot{I} = (\beta S - \gamma) I, \tag{S77b}$$

$$418 \quad \dot{R} = \gamma I, \tag{S77c} \\
 419$$

420 with initial conditions given by

$$421 \quad S(0) = (1 - p)(1 - \alpha)(1 - \phi_{\text{inst}})$$

$$422 \quad I(0) = (1 - p)\alpha$$

$$423 \quad R(0) = 0$$

$$424 \quad V(0) = p + \phi_{\text{inst}}(1 - p)(1 - \alpha). \\
 425$$

426 Note that for this model, $S(0)$ is the density of susceptibles after the post-outbreak vaccina-
 427 tion response has taken place.

428 Equation (S77a) implies that

$$429 \quad -\frac{\gamma}{\beta} \frac{d}{dt} \ln(S) = \dot{R}, \quad (\text{S78})$$

430 thus S_∞ satisfies the equation

$$431 \quad \frac{\gamma}{\beta} \ln\left(\frac{S(0)}{S_\infty}\right) = R_\infty = 1 - V(0) - S_\infty, \quad (\text{S79})$$

432 OR

$$433 \quad S_\infty = -\frac{\gamma}{\beta} W_0\left(-\frac{\beta}{\gamma} S(0) e^{-\frac{\beta}{\gamma}(1-V(0))}\right). \quad (\text{S80})$$

434 We use the principle branch of the Lambert function in order to obtain solutions satisfying
 435 $S_\infty \leq S(0)$. Since $-\frac{\beta}{\gamma} S(0) e^{-\frac{\beta}{\gamma}(1-V(0))} > -\frac{\beta}{\gamma} S(0) e^{-\frac{\beta}{\gamma} S(0)}$, and W_1 is monotonically decreasing,
 436 $-\frac{\gamma}{\beta} W_1\left(-\frac{\beta}{\gamma} S(0) e^{-\frac{\beta}{\gamma}(1-V(0))}\right) > S(0)$, which does not correspond to biologically feasible
 437 solutions. In addition, we have $R_\infty = \frac{\gamma}{\beta} \ln \frac{S(0)}{S_\infty}$.

438 Since there is no vaccination except during the initial (immediate) response to the out-
 439 break, $V_\infty = V(0) = p + \phi_{\text{inst}}(1-p)(1-\alpha)$, and so

$$440 \quad \psi_p = \frac{V_\infty - p}{1-p} = \phi_{\text{inst}}(1-\alpha) \quad (\text{S81a})$$

$$441 \quad \pi_p = 1 - \psi_p - \frac{S_\infty}{1-p}. \quad (\text{S81b})$$

443 Using Equation (S79), we also have

$$444 \quad \pi_p = -\frac{\gamma}{\beta(1-p)} \ln\left(\frac{1 - \pi_p - \phi_{\text{inst}}(1-\alpha)}{(1-\alpha)(1-\phi_{\text{inst}})}\right) \quad (\text{S82})$$

445 It follows from Equation (S81b) that π_p is a decreasing function of p : from Equation (S79)
 446 we have

$$447 \quad \frac{\gamma}{\beta} \left(-\frac{1}{1-p} - \frac{\partial_p S_\infty}{S_\infty}\right) = -1 + \phi_{\text{inst}}(1-\alpha) - \partial_p S_\infty \quad (\text{S83})$$

448 and so

$$449 \quad \partial_p S_\infty = \left(-1 + \phi_{\text{inst}}(1-\alpha) + \frac{\gamma}{\beta} \frac{1}{1-p}\right) \left(\frac{\beta S_\infty}{\beta S_\infty - \gamma}\right). \quad (\text{S84})$$

450 This gives

$$451 \quad (1-p)\partial_p S_\infty + S_\infty = \frac{\beta S_\infty}{\beta S_\infty - \gamma} ((1-p)(-1 + \phi_{\text{inst}}(1-\alpha)) + S_\infty)$$

$$452 \quad = \frac{\beta S_\infty}{\gamma - \beta S_\infty} (1 - V_\infty - S_\infty), \quad (\text{S85})$$

453

454 which is positive so long as $R_\infty > 0$ (this happens when $S(0) > S_\infty$, which is true whenever
 455 $S(0)$ and $I(0)$ are not 0). It now follows that $\partial_p \pi_p = -\frac{(1-p)\partial_p S_\infty + S_\infty}{(1-p)^2} < 0$. Note that the
 456 probability of a delayer being vaccinated post-outbreak (ψ_p) is constant.

457 Lastly, note that

$$458 \quad (1-p)\partial_p S_\infty = \left(1 - V_\infty - \frac{\gamma}{\beta}\right) \left(\frac{\beta S_\infty}{\gamma - \beta S_\infty}\right), \quad (\text{S86})$$

459 which implies that S_∞ increases with p iff $(1-p)(1 - \phi_{\text{inst}}(1 - \alpha))\beta > \gamma$, or equivalently,
 460 $I(0) + S(0) = 1 - V_\infty > \frac{\gamma}{\beta}$. Compare this to the more stringent condition $S(0) > \frac{\gamma}{\beta}$ which
 461 ensures that the epidemic takes off ($I'(0) > 0$).

462 **F Maximal vaccination rate for fair comparison of mod-** 463 **els**

464 In this section, we find the fair comparison values of the vaccination efforts ϕ_{inc} and ϕ_{prev}
 465 (see § 6.2). These are defined as the levels of vaccination effort ϕ_{inc} and ϕ_{prev} that result in
 466 maximal vaccination rates equal to 0.1/day (that is, comparable to [7]).

467 **F.1 Maximal Vaccination rate when $\dot{V} = \phi_{\text{prev}}I$**

468 We begin by finding what the maximal vaccination is when $\dot{V} = \phi_{\text{prev}}I$. Because the vac-
 469 cination rate, \dot{V} , is maximal when prevalence, I , is maximal, we aim to find the maximal
 470 prevalence. Now observe that since $\dot{I} = (\beta S - \gamma)I$, incidence is maximal when $S = \gamma/\beta$.
 471 Thus, the peak prevalence is found by substituting $S = \gamma/\beta$, into Equation (S21) to obtain

$$472 \quad I_{\text{peak}} = 1 - p - \gamma/\beta + \frac{\gamma + \phi_{\text{prev}}}{\beta} \ln \left(\frac{\gamma + \phi_{\text{prev}}}{\beta S(0) + \phi_{\text{prev}}} \right) \quad (\text{S87})$$

473 (recall that $I(0) + S(0) = 1 - p$). We now wish to find at which value of p the maximal
 474 vaccination rate (over time) is largest. Observe that

$$475 \quad \frac{\partial}{\partial p} I_{\text{peak}} = -1 + \frac{(\gamma + \phi_{\text{prev}})(1 - \alpha)}{\beta S(0) + \phi_{\text{prev}}}, \quad (\text{S88})$$

$$476 \quad \frac{\partial^2}{\partial p^2} I_{\text{peak}} = \frac{\beta(\gamma + \phi_{\text{prev}})(1 - \alpha)^2}{(\beta S(0) + \phi_{\text{prev}})^2} > 0, \quad (\text{S89})$$

478 which implies that the peak prevalence (and thus the maximal vaccination rate) has a minimum
 479 in p when

$$480 \quad p_{\text{crit}} = \frac{(\beta - \gamma)(1 - \alpha) + \phi_{\text{prev}}\alpha}{\beta(1 - \alpha)} > 0 \quad (\text{S90})$$

481 ($\beta > \gamma$ because $\mathcal{R}_0 > 1$).

482 There are now two possibilities:

- 483 • If $p_{\text{crit}} \geq 1$ (which happens *iff* $\phi_{\text{prev}}\alpha \leq \gamma(1 - \alpha)$) then the maximal vaccination rate
484 is attained when $p = 0$.
- 485 • If $p_{\text{crit}} < 1$ (which happens *iff* $\phi_{\text{prev}}\alpha > \gamma(1 - \alpha)$) then, the maximal vaccination rate
486 must be attained either when $p = 0$ or when $p = 1$.

487 Noting that

$$\begin{aligned}
488 \quad I_{\text{peak}}|_{p=0} &= 1 - \gamma/\beta + \frac{\gamma + \phi_{\text{prev}}}{\beta} \ln \left(\frac{\gamma + \phi_{\text{prev}}}{\beta(1 - \alpha) + \phi_{\text{prev}}} \right) \\
489 \quad I_{\text{peak}}|_{p=1} &= -\gamma/\beta + \frac{\gamma + \phi_{\text{prev}}}{\beta} \ln \left(\frac{\gamma + \phi_{\text{prev}}}{\phi_{\text{prev}}} \right), \tag{S91} \\
490
\end{aligned}$$

491 it follows that

$$\begin{aligned}
492 \quad \dot{V}_{\text{max}} &= -\phi_{\text{prev}} \frac{\gamma}{\beta} + \phi_{\text{prev}} \max \left\{ 1 + \frac{\gamma + \phi_{\text{prev}}}{\beta} \ln \left(\frac{\gamma + \phi_{\text{prev}}}{\beta(1 - \alpha) + \phi_{\text{prev}}} \right), \right. \\
493 &\quad \left. \frac{\gamma + \phi_{\text{prev}}}{\beta} \ln \left(\frac{\gamma + \phi_{\text{prev}}}{\phi_{\text{prev}}} \right) \right\}. \tag{S92} \\
494
\end{aligned}$$

495 We also note that

$$\begin{aligned}
496 \quad \frac{\partial}{\partial \phi_{\text{prev}}} I_{\text{peak}} &= \frac{1}{\beta} \ln \left(\frac{\gamma + \phi_{\text{prev}}}{\beta S(0) + \phi_{\text{prev}}} \right) + \frac{1}{\beta} \left(1 - \frac{\gamma + \phi_{\text{prev}}}{\beta S(0) + \phi_{\text{prev}}} \right). \tag{S93} \\
497
\end{aligned}$$

497 Because $1 - x + \ln(x) < 0$ for any $0 < x \neq 1$, it follows that the peak prevalence I_{peak} , and
498 thus the peak vaccination rate, decreases with increasing vaccination effort, ϕ_{prev} (for any
499 initial coverage, p). Also, as $\phi_{\text{prev}} \rightarrow \infty$, we have

$$\begin{aligned}
500 \quad I_{\text{peak}}|_{p=0} &\rightarrow \alpha, \\
501 \quad I_{\text{peak}}|_{p=1} &\rightarrow 0. \\
502
\end{aligned}$$

503 Setting $\dot{V}_{\text{max}} = 0.1/\text{day}$ in Equation (S92) we can numerically solve for ϕ_{prev} , with $\alpha, \beta,$
504 γ , as in Tables 1 and 2 to obtain $\phi_{\text{prev}} \approx 1582/\text{day}$.

505 F.2 Maximal Vaccination rate when $\dot{V} = \phi_{\text{inc}}SI$

506 First, we will derive a formula for the maximal vaccination rate as it depends on the model
507 parameters, $\alpha, \beta, \gamma, \phi_{\text{inc}}$. We then use this formula to calculate the appropriate range for
508 ϕ_{inc} , given the estimates of the other parameters cited in Tables 1 and 2.

509 Differentiating Equation (S46d), we have

$$\begin{aligned}
510 \quad \ddot{V} &= \phi_{\text{inc}}(\dot{S}I + S\dot{I}) = \phi_{\text{inc}}SI(\beta S - \gamma - (\beta + \phi_{\text{inc}})I).
\end{aligned}$$

511 Thus, critical points of \dot{V} (excluding those for which $\dot{V} = 0$) occur when $\beta S - \gamma = (\beta + \phi_{\text{inc}})I$.
512 Using Equation (S48) and simplifying, this is equivalent to

$$\begin{aligned}
513 \quad 2\beta S &= \gamma \ln S + \phi_{\text{inc}}(1 - p)\alpha + \beta(1 - p) + \gamma \left(1 - \ln((1 - p)(1 - \alpha)) \right),
\end{aligned}$$

514 which has two formal solutions,

$$515 \quad \hat{S}_k = -\frac{\gamma}{2\beta} W_k \left(-2\frac{\beta}{\gamma}(1-p)(1-\alpha)e^{-\frac{\beta+\phi_{\text{inc}}\alpha}{\gamma}(1-p)-1} \right),$$

516 with $k = 0$ or -1 .

517 However, it is impossible for \hat{S}_0 to be attained by $S(t)$, for all $t \geq 0$. To see this, suppose,
 518 in order to derive a contradiction, that there is some time $\hat{t}_0 \geq 0$ such that $S(\hat{t}_0) = \hat{S}_0$. Note
 519 that $-1 \leq W_0 < 0$ on the interval $[-1/e, 0)$, so $0 < \hat{S}_0 < \gamma/\beta$. Because $\hat{S}_0 < \gamma/\beta$, we have
 520 $\dot{I}(\hat{t}_0) < 0$. Since $\dot{S} < 0$ (for all time t), it follows that $\ddot{V} < 0$ when $t = \hat{t}_0$, in contradiction
 521 to the fact that by definition of \hat{S}_0 , $\ddot{V}(\hat{t}_0) = 0$. Thus, $S(t) > \hat{S}_0$ is proven¹. It follows that if
 522 there is a biologically relevant value of S at which \ddot{V} changes signs, it must be

$$523 \quad \hat{S}_{-1} = -\frac{\gamma}{2\beta} W_{-1} \left(-2\frac{\beta}{\gamma e}(1-p)(1-\alpha)e^{-\frac{\beta+\phi_{\text{inc}}\alpha}{\gamma}(1-p)} \right). \quad (\text{S94})$$

524 Note that $\hat{S}_{-1} > \frac{\gamma}{2\beta}$ because $W_{-1}(x) < -1 \quad \forall x \in [-1/e, 0]$ (but it is also possible that
 525 $\hat{S}_{-1} > S(0) = (1-p)(1-\alpha)$, which would make this critical point biologically unfeasible).

526 Because S decreases with time, we see that \ddot{V} can change signs at most once for all $t \geq 0$.
 527 Observe that since $0 < S_\infty < \gamma/\beta$, and \dot{V} decreases when $S \in (0, \gamma/\beta]$, it follows that \dot{V}
 528 eventually (*i.e.*, for large enough t) decreases with time. Hence, if $\dot{V}(0) \leq 0$, then $t = 0$ is a
 529 maximum of \dot{V} for $t \geq 0$, and if $\dot{V}(0) > 0$ then \dot{V} attains its maximum when $S(t) = \hat{S}_{-1}$. The
 530 sign of $\ddot{V}(0)$ is identical to the sign of $\beta S(0) - \gamma - (\beta + \phi_{\text{inc}})I(0)$, so the maximal vaccination
 531 rate is attained at $t = 0$ if $(1-p)((1-2\alpha)\beta - \alpha\phi_{\text{inc}}) \leq \gamma$, and when $S(t) = \hat{S}_{-1}$ otherwise.
 532 Thus,

$$533 \quad \nu(p) = \max_{t \geq 0} \dot{V} = \begin{cases} \phi_{\text{inc}}(1-p)^2(1-\alpha)\alpha & \text{if } (1-p)((1-2\alpha)\beta - \alpha\phi_{\text{inc}}) \leq \gamma, \\ \frac{\phi_{\text{inc}}}{\phi_{\text{inc}} + \beta}(\beta\hat{S}_{-1} - \gamma)\hat{S}_{-1} & \text{if } (1-p)((1-2\alpha)\beta - \alpha\phi_{\text{inc}}) > \gamma, \end{cases} \quad (\text{S95})$$

534 (where, for the second case, we used the fact that $I = \frac{\beta\hat{S}_{-1} - \gamma}{\phi_{\text{inc}} + \beta}$ when $S = \hat{S}_{-1}$).

535 To maximize ν over all $p \in [0, 1)$ (with α, β, γ and ϕ_{inc} fixed), we consider the following
 536 3 cases:

- 537 • First, if $0 < \gamma < (1-2\alpha)\beta - \alpha\phi_{\text{inc}}$, then $(1-p)((1-2\alpha)\beta - \alpha\phi_{\text{inc}}) \leq \gamma$ is equivalent to
 538 $\hat{p} = 1 - \frac{\gamma}{(1-2\alpha)\beta - \alpha\phi_{\text{inc}}} \leq p$, and $\hat{p} \in (0, 1)$. Hence, for $p \in [\hat{p}, 1)$, $\nu(p) = \phi_{\text{inc}}(1-p)^2(1-\alpha)\alpha$
 539 is a decreasing function of p , and so $\max_{p \in [\hat{p}, 1)} \nu(p) = \phi_{\text{inc}} \left(\frac{\gamma}{(1-2\alpha)\beta - \alpha\phi_{\text{inc}}} \right)^2 (1-\alpha)\alpha$, and
 540 is attained when $p = \hat{p}$.

541 When $0 \leq p < \hat{p}$, we note that because $(\beta x - \gamma)x$ is parabolic with a minimum at
 542 $x = \gamma/2\beta$, and $\hat{S}_{-1} \geq \frac{\gamma}{2\beta}$, it follows that

$$543 \quad \nu(p) = \frac{\phi_{\text{inc}}}{\phi_{\text{inc}} + \beta}(\beta\hat{S}_{-1} - \gamma)\hat{S}_{-1} \quad (\text{S96})$$

¹Consequently, $S_\infty \geq \hat{S}_0$, which is equivalent to a statement about the Lambert W function: $W_0(2x/e) \geq 2W_0(x)$, for $-1/e < x < 0$.

544 is increasing in \hat{S}_{-1} , so $\max_{p \in [0, \hat{p}]} \nu(p)$ is attained on this interval when \hat{S}_{-1} is maxi-
 545 mized. Because $-\frac{\gamma}{2\beta} W_{-1}(x)$ is monotonically increasing, it follows that $\max_{p \in [0, \hat{p}]} \nu(p)$
 546 is maximal when

$$547 \quad -2 \frac{\beta}{\gamma e} (1-p)(1-\alpha) e^{-\frac{\beta + \phi_{\text{inc}} \alpha}{\gamma} (1-p)}$$

548 is maximal. Consequently, we need to maximize $-axe^{-x}$ (with $a > 0$), with $x(p) =$
 549 $\frac{\beta + \phi_{\text{inc}} \alpha}{\gamma} (1-p)$, over the interval $0 \leq p \leq \hat{p}$. This corresponds to maximizing $-axe^{-x}$ over
 550 $[\frac{\beta + \phi_{\text{inc}} \alpha}{\gamma}, \frac{\beta + \phi_{\text{inc}} \alpha}{(1-2\alpha)\beta - \alpha\phi_{\text{inc}}}] \subset [1, \infty)$. Observe that $-axe^{-x}$ has a unique global minimum
 551 at $x = 1$, and in particular, it is increasing when $x \geq 1$. This implies that in the
 552 relevant range of x , \hat{S}_{-1} increases with x , and thus decreases in p . It follows that \hat{S}_{-1}
 553 is maximal when $p = 0$, and its value is

$$554 \quad \hat{S}_{-1}|_{p=0} = -\frac{\gamma}{2\beta} W_{-1} \left(-2 \frac{\beta}{\gamma e} (1-\alpha) e^{-\frac{\beta + \phi_{\text{inc}} \alpha}{\gamma}} \right). \quad (\text{S97})$$

555 Thus, $\max_{p \in [0, 1]} \nu(p) = \nu(0)$, and is attained when $p = 0$ (note also that $\nu(p)$ is
 556 continuous at $p = \hat{p}$).

- 557 • When $0 < (1-2\alpha)\beta - \alpha\phi_{\text{inc}} \leq \gamma$, $(1-p)((1-2\alpha)\beta - \alpha\phi_{\text{inc}}) \leq \gamma$ is equivalent to
 558 $\hat{p} = 1 - \frac{\gamma}{(1-2\alpha)\beta - \alpha\phi_{\text{inc}}} \leq p$, which is satisfied for all $p \in [0, 1]$, since $\hat{p} \leq 0$. Thus, $\nu(p)$ is
 559 a decreasing function of p for all $p \in [0, 1]$, and thus $\max_{p \in [0, 1]} \nu(p) = \phi_{\text{inc}}(1-\alpha)\alpha$ is
 560 attained at $p = 0$.
- 561 • When $\alpha\phi_{\text{inc}} \geq (1-2\alpha)\beta$, then $(1-p)((1-2\alpha)\beta - \alpha\phi_{\text{inc}}) \leq \gamma$ is always satisfied (since
 562 the left hand side is never positive, and $\gamma > 0$). Thus, $\nu(p) = \phi_{\text{inc}}(1-p)^2(1-\alpha)\alpha$,
 563 which decreases with p , so $\max_{p \in [0, 1]} \nu(p) = \phi_{\text{inc}}(1-\alpha)\alpha$, and is attained at $p = 0$.

564 Rearranging the conclusions of the preceding discussion, we see that

$$565 \quad \max_{p \in [0, 1], t \geq 0} \dot{V} = \begin{cases} \frac{\phi_{\text{inc}}}{\phi_{\text{inc}} + \beta} (\beta \hat{S}_{-1} - \gamma) \hat{S}_{-1}|_{p=0} & \text{if } 0 \leq \alpha\phi_{\text{inc}} < (1-2\alpha)\beta - \gamma, \\ \alpha\phi_{\text{inc}}(1-\alpha) & \text{if } (1-2\alpha)\beta - \gamma \leq \phi_{\text{inc}}\alpha. \end{cases} \quad (\text{S98})$$

567 When $0 \leq \alpha\phi_{\text{inc}} < (1-2\alpha)\beta - \gamma$, then $\max_{p \in [0, 1], t \geq 0} \dot{V} = \frac{\phi_{\text{inc}}}{\phi_{\text{inc}} + \beta} (\beta \hat{S}_{-1} - \gamma) \hat{S}_{-1}|_{p=0}$,
 568 which increases as $\hat{S}_{-1}|_{p=0}$ increases, as stated earlier. Since $-W_{-1}(x)$ increases with x ,
 569 and $-2 \frac{\beta}{\gamma e} (1-\alpha) e^{-\frac{\beta + \phi_{\text{inc}} \alpha}{\gamma}}$ is an increasing function of ϕ_{inc} , we conclude that in this range,
 570 $\max_{p \in [0, 1], t \geq 0} \dot{V}$ increases with ϕ_{inc} . When $(1-2\alpha)\beta - \gamma \leq \phi_{\text{inc}}\alpha$, $\max_{p \in [0, 1]} \nu(p)$ manifestly
 571 increases linearly with ϕ_{inc} . In all, $\max_{p \in [0, 1], t \geq 0} \dot{V}$ is a monotonically increasing function of
 572 ϕ_{inc} .

573 Note that at the point separating the two regimes, $\phi_{\text{inc}} = \frac{(1-2\alpha)\beta - \gamma}{\alpha} = 16570.71/\text{day}$,
 574 $\max_{p \in [0, 1], t \geq 0} \dot{V} = ((1-2\alpha)\beta - \gamma)(1-\alpha) = 0.29/\text{day}$ (with parameters as in Table 1).

575 Finally, to obtain a value of ϕ_{inc} that yields a maximal vaccination rate of $\Phi = 0.1/\text{day}$
 576 (as was estimated in [7]), we solve

$$577 \quad \frac{\phi_{\text{inc}}}{\phi_{\text{inc}} + \beta} (\beta \hat{S}_{-1} - \gamma) \hat{S}_{-1}|_{p=0} = \Phi, \quad (\text{S99})$$

578 which gives $\hat{S}_{-1}|_{p=0} = \frac{\gamma \pm \sqrt{\gamma^2 + 4\beta\Phi(1+\beta/\phi_{\text{inc}})}}{2\beta}$. We take the solution corresponding to the positive
 579 sign (the other one gives negative \hat{S}_{-1} , which is biologically absurd). Thus,

$$580 \quad W_{-1} \left(-2 \frac{\beta}{\gamma e} (1 - \alpha) e^{-\frac{\beta + \phi_{\text{inc}} \alpha}{\gamma}} \right) = -1 - \sqrt{1 + 4 \frac{\beta}{\gamma^2} \Phi(1 + \beta/\phi_{\text{inc}})}, \quad (\text{S100})$$

581 which is equivalent to

$$582 \quad \frac{2\beta(1 - \alpha)}{\gamma(1 + \sqrt{1 + 4 \frac{\beta}{\gamma^2} \Phi(1 + \beta/\phi_{\text{inc}})})} = \exp \left(\frac{\beta + \phi_{\text{inc}} \alpha}{\gamma} - \sqrt{1 + 4 \frac{\beta}{\gamma^2} \Phi(1 + \beta/\phi_{\text{inc}})} \right),$$

583 which we solve numerically for ϕ_{inc} , with parameters as in Tables 1 and 2, to get $\phi_{\text{inc}} \approx$
 584 5190/day.

585 **G The individual equilibrium**

586 In this section, we show that for each of the five models defined in § 5, the game defined
 587 in § 3 always has a unique convergently stable Nash equilibrium (defined in § 3 and abbrevi-
 588 ated CSNE). The proofs given here are constructive, *i.e.*, they also provide a method for
 589 numerically finding the individual equilibrium (p_i).

590 **G.1 Vaccination rate \propto disease prevalence**

591 In this scenario we have 3 cases to examine:

- 592 1. $p_m \geq 1$ ($\iff \alpha\phi_{\text{prev}} \geq \gamma(1 - \alpha)$)
- 593 2. $p_m \leq 1$ and $p_0 \leq 0$
- 594 3. $p_m \leq 1$ and $p_0 \in (0, 1)$

595 Recall that in the first case, we have proven (in Appendix E.1) that π_p decreases with p . As
 596 stated in § 5.1, we assume that π_p decreases with p for the other two cases as well.

597 **G.1.1 $p_m \geq 1$**

598 In this case, we have a unique convergently stable Nash equilibrium (CSNE). To see this,
 599 note that $\psi_p + \pi_p = 1$ and $\partial_p \pi_p < 0$. Thus,

$$600 \quad \begin{aligned} \Delta E &= [\pi_p + (1 - \pi_p)r - r/a]a(P - Q) \\ 601 \quad &= [\pi_p a(1 - r) - r(1 - a)](P - Q) \\ 602 \quad &= a(1 - r) \left[\pi_p - \frac{r(1 - a)}{a(1 - r)} \right] (P - Q). \end{aligned} \quad (\text{S101})$$

603

604 It is convenient to define

$$605 \quad \rho_1 = \frac{r(1-a)}{a(1-r)} = \frac{r}{1-r} \bigg/ \frac{a}{1-a}, \quad (\text{S102})$$

606 which we can interpret as an **odds ratio**, namely the odds of a bad outcome from vaccination
 607 (compared with infection) relative to the odds of an outbreak occurring. The odds ratio is
 608 well-defined and strictly positive ($\rho_1 > 0$) because $0 < r < 1$ and $0 < a < 1$. Since π_p
 609 decreases monotonically with p , there are three cases:

- 610 • If $\pi_0 \leq \rho_1$ then $\pi_p < \rho_1 \quad \forall p > 0$. It follows that $\forall \epsilon \in [0, 1) \quad \Delta E > 0 \quad \forall Q \neq P \iff$
 611 $P = 0$. Hence $p_i = 0$ is the unique Nash equilibrium. Let $0 \leq P < Q$ and fix $\epsilon \in [0, 1)$.
 612 It follows that $p > 0$, and so $\pi_p - \rho_1 < 0$. Thus $\Delta E > 0$ and p_i is convergently stable.
- 613 • If $\alpha = \pi_1 \geq \rho_1$, then $\pi_p > \rho_1 \quad \forall p < 1$. It follows that $\forall \epsilon \in [0, 1) \quad \Delta E > 0 \quad \forall Q \neq$
 614 $P \iff P = 1$. Hence $p_i = 1$ is the unique Nash equilibrium. The condition translates
 615 to $r(1-a) < \alpha a(1-r)$, or $r_v < ar_i\alpha + ar_v(1-\alpha)$. Recall that if an outbreak occurs,
 616 at the end of the epidemic individuals have either been vaccinated or have contracted
 617 the disease. Thus the right hand side is the risk to a vaccinator, and the left hand side
 618 is the minimal possible risk to a delayer (assuming no-one is infected after the initial
 619 outbreak; if there are secondary infections, then because $r_v < r_i$, the delayer's risk can
 620 only be increased). Let $1 \geq P > Q$ and fix $\epsilon \in [0, 1)$. It follows that $p < 1$, and so
 621 $\pi_p - \rho_1 > 0$. Thus $\Delta E > 0$ and p_i is convergently stable.
- 622 • If $\pi_0 > \rho_1 > \pi_1 = \alpha$ then there is a unique $\tilde{p} \in (0, 1)$ such that $\pi_p - \rho_1 > 0$ if $p < \tilde{p}$,
 623 $\pi_{\tilde{p}} = \rho_1$ and $\pi_p - \rho_1 < 0$ if $p > \tilde{p}$. Now since for any $\epsilon \in [0, 1)$, $Q < P \implies p < P$
 624 and $Q > P \implies p > P$, we have $\forall \epsilon \in [0, 1) \quad \Delta E > 0 \quad \forall Q \neq P \iff P = \tilde{p}$
 625 (for other P take Q between P and \tilde{p}). Thus, the unique Nash equilibrium p_i is the
 626 unique solution to $\pi_{p_i} = \rho_1$. Fix $\epsilon \in [0, 1)$. If $Q < P \leq p_i$, $Q \leq p < P \leq p_i$. Thus
 627 $\pi_p - \rho_1 > 0 \implies \Delta E > 0$. Similarly, If $Q > P \geq p_i$, $Q \geq p > P \geq p_i$. Thus
 628 $\pi_p - \rho_1 < 0 \implies \Delta E > 0$. Hence p_i is convergently stable.

629 Now, to find \tilde{p} : recall that $1 - \pi_p = \frac{\phi_{\text{prev}}}{\beta(1-p)} \ln \left(\frac{\beta}{\phi_{\text{prev}}} (1-\alpha)(1-p) + 1 \right)$ and thus

$$630 \quad \tilde{p} = 1 + \phi_{\text{prev}} \frac{(1-\rho_1) + (1-\alpha) W \left(-\frac{1-\rho_1}{1-\alpha} e^{-\frac{1-\rho_1}{1-\alpha}} \right)}{(1-\alpha)\beta(1-\rho_1)}. \quad (\text{S103})$$

631 Again, W is applied to a negative argument, and it is necessary to determine which
 632 branch of W to use. The principal branch gives $\tilde{p} = 1$, and $\pi_p \rightarrow \alpha$ as $p \rightarrow 1$, and
 633 $\rho_1 > \alpha$ by assumption, and so by elimination we must use W_{-1} . Interestingly, \tilde{p}
 634 depends linearly on ϕ_{prev} . Recall that $W_{-1} \leq -1$, and so $\frac{\partial \tilde{p}}{\partial \phi_{\text{prev}}} < 0$.

635 Thus, in all three cases there exists a unique CNSE.

636 **G.1.2** $p_m \leq 1$ and $p_0 \leq 0$

637 In the 2nd case, recall that $\psi_p = \frac{\phi_{\text{prev}}}{\gamma} \pi_p$, and let

$$638 \quad \rho_2 = \frac{r}{a(1 + r\phi_{\text{prev}}/\gamma)}, \quad (\text{S104})$$

639 to obtain

$$\begin{aligned}
 640 \quad \Delta E &= [\pi_p + \frac{\phi_{\text{prev}}}{\gamma} \pi_p r - r/a] a(P - Q) \\
 641 \quad &= [\pi_p a(1 + \frac{\phi_{\text{prev}}}{\gamma} r) - r](P - Q) \\
 642 \quad &= a(1 + \frac{\phi_{\text{prev}}}{\gamma} r) [\pi_p - \frac{\gamma r}{a(r\phi_{\text{prev}} + \gamma)}] (P - Q) \\
 643 \quad &= a(1 + \frac{\phi_{\text{prev}}}{\gamma} r) [\pi_p - \rho_2] (P - Q). \quad (\text{S105}) \\
 644
 \end{aligned}$$

645 Now recall that π_p decreases with p . So, an identical argument to the one in Ap-
 646 pendix G.1.1 also applies here:

- 647 • If $\pi_0 \leq \rho_2$ then $\pi_p < \rho_2 \quad \forall p > 0$. Thus, $\forall \epsilon \in [0, 1) \quad \Delta E > 0 \quad \forall Q \neq P \iff P = 0$.
 648 Hence $p_i = 0$ is the unique Nash equilibrium. Let $0 \leq Q < P < 1$ and fix $\epsilon \in [0, 1)$. It
 649 follows that $p > 0$, and so $\pi_p - \rho_2 < 0$. Thus $\Delta E > 0$ and p_i is convergently stable.
- 650 • If $\alpha = \pi_1 \geq \rho_2 \geq 0$, then $\pi_p > \rho_2 \quad \forall p < 1$. It follows that $\forall \epsilon \in [0, 1) \quad \Delta E > 0 \quad \forall Q \neq$
 651 $P \iff P = 1$. Hence $p_i = 1$ is the unique Nash equilibrium. Let $1 \geq P > Q$ and
 652 fix $\epsilon \in [0, 1)$. It follows that $p < 1$, and so $\pi_p - \rho_2 > 0$. Thus $\Delta E > 0$ and p_i is
 653 convergently stable.
- 654 • If $\pi_0 > \rho_2 > \pi_1 = \alpha$ then there is a unique $\tilde{p} \in (0, 1)$ such that $\pi_p - \rho_2 > 0$ if $p < \tilde{p}$,
 655 $\pi_{\tilde{p}} = \rho_2$ and $\pi_p - \rho_2 < 0$ if $p > \tilde{p}$. Now since for any $\epsilon \in [0, 1)$, $Q < P \implies p < P$
 656 and $Q > P \implies p > P$, we have $\forall \epsilon \in [0, 1) \quad \Delta E > 0 \quad \forall Q \neq P \iff P = \tilde{p}$.
 657 Thus, the unique Nash equilibrium p_i is the unique solution to $\pi_{p_i} = \rho_2$. Fix $\epsilon \in [0, 1)$.
 658 If $Q < P \leq p_i$, $Q \leq p < P \leq p_i$. Thus $\pi_p - \rho_2 > 0 \implies \Delta E > 0$. Similarly,
 659 If $Q > P \geq p_i$, $Q \geq p > P \geq p_i$. Thus $\pi_p - \rho_2 < 0 \implies \Delta E > 0$. Hence p_i is
 660 convergently stable.

661 To find \tilde{p} , recall that $R_\infty^1 = (1-p)\pi_p$. Furthermore, from Equation (S27), we have

$$\begin{aligned}
662 \quad R_\infty^1 &= \frac{\gamma}{\beta} \ln \left(\frac{\beta S(0) + \phi_{\text{prev}}}{\beta S_\infty^1 + \phi_{\text{prev}}} \right) \\
663 \quad S_\infty^1 &= 1 - p - \frac{\phi_{\text{prev}} + \gamma}{\gamma} R_\infty^1 \\
664 \quad &\Downarrow \\
665 \quad R_\infty^1 &= \frac{\gamma}{\beta} \ln \left(\frac{\beta S(0) + \phi_{\text{prev}}}{\beta(1-p - \frac{\phi_{\text{prev}} + \gamma}{\gamma} R_\infty^1) + \phi_{\text{prev}}} \right) \\
666 \quad &\Downarrow \\
667 \quad (1-p)\pi_p &= \frac{\gamma}{\beta} \ln \left(\frac{\beta S(0) + \phi_{\text{prev}}}{\beta(1-p) \left(1 - \frac{\phi_{\text{prev}} + \gamma}{\gamma} \pi_p \right) + \phi_{\text{prev}}} \right).
\end{aligned}$$

669 Substituting $p = \tilde{p}$, and using $\pi_{\tilde{p}} = \rho_2 = \frac{\gamma r}{a(r\phi_{\text{prev}} + \gamma)}$, we obtain after minor rearrange-
670 ment

$$671 \quad \frac{\beta(1-\tilde{p})r}{a(r\phi_{\text{prev}} + \gamma)} = \ln \left(\frac{a(\beta(1-\tilde{p})(1-\alpha) + \phi_{\text{prev}})(r\phi_{\text{prev}} + \gamma)}{\beta(1-\tilde{p})(\gamma(a-r) - (1-a)r\phi_{\text{prev}}) + \phi_{\text{prev}}a(r\phi_{\text{prev}} + \gamma)} \right). \tag{S106}$$

673 However, we have not succeeded in obtaining an analytical solution for the individual
674 equilibrium from this equation.

675 Thus, in all three cases there exists a unique CSNE.

676 **G.1.3** $p_m \leq 1$ and $p_0 \in (0, 1)$

677 Since π_p decreases with p , the argument above shows that there is a unique CSNE in each
678 of the the two intervals $[0, p_0]$ and $[p_0, 1]$, denoted $P_{I,1}$ and $P_{I,2}$, respectively. These are
679 the only candidates for Nash equilibria in the interval $[0, 1]$: Adding the two sub-intervals
680 together amounts to adding more strategies to the game. Thus, a strategy which was a
681 Nash equilibrium in one of the sub-intervals may not be a Nash equilibrium for the larger
682 strategy set (because players now have a larger strategy set to choose from). However, a
683 strategy which is a Nash equilibrium for $[0, 1]$ must be a Nash equilibrium in any sub-interval
684 of $[0, 1]$ which contains it. The situation for convergent stability is a bit more subtle, and is
685 considered below.

686 Note that when $p \neq p_0$,

$$687 \quad \text{sgn}(\Delta E) = \text{sgn}((\pi_p - \rho(p))(P - Q)) \tag{S107}$$

688 where

$$689 \quad \rho(p) = \begin{cases} \rho_1 & \text{if } p < p_0, \\ \rho_2 & \text{if } p \geq p_0. \end{cases} \tag{S108}$$

690 Note also that

$$691 \quad \pi_{p_0} = \rho_1 \iff r = \frac{\gamma a}{\phi_{\text{prev}}(1-a) + \gamma} \iff \pi_{p_0} = \rho_2. \quad (\text{S109})$$

693 This may seem slightly perplexing at first, but recall that $\pi_{p_0} = \frac{\gamma}{\phi_{\text{prev}} + \gamma}$. Thus, if it so
 694 happens that $\pi_{p_0} = \rho_1$ or $\pi_{p_0} = \rho_2$, r and a must be related so that in fact $\rho_1 = \rho_2$.

695 We must now check a number of cases:

- 696 1. $\pi_0 \leq \rho_1$ and $\pi_{p_0} < \rho_2$: In this case, $\pi_p < \rho(p) \quad \forall p \in (0, 1]$, and $P_{I,1} = 0$ and $P_{I,2} = p_0$.
 697 But if $P = p_0$ and $Q \in [0, p_0)$, $\Delta E < 0$ and so $P = p_0$ cannot be a Nash equilibrium.
 698 However, if $P = 0$, then $\forall \epsilon \in [0, 1) \quad \Delta E > 0 \quad \forall Q \neq P$, and so $P = 0$ is a Nash
 699 equilibrium. This is trivial for $Q \leq p_0$. For $p_0 < Q$, we have $\pi_p - \rho(p) < 0$ and
 700 $P - Q < 0$, so $\Delta E > 0$ as required. For convergent stability, we only need to check
 701 that if $0 < P < p_0 < Q \leq 1$, then for any $\epsilon \in [0, 1)$ we have $\Delta E > 0$. In this case,
 702 $p \in [P, Q]$, and $P - Q < 0$. Furthermore, $\pi_p - \rho_1 < 0$ for any $p \leq p_0$, and $\pi_p - \rho_2 < 0$
 703 for any $p \geq p_0$. But since $\pi_{p_0} - \rho_2 < 0$, $\pi_{p_0} - \rho_1 = 0$ is impossible and from continuity,
 704 $\pi_p - \rho_1 < 0$. Thus, $\Delta E(P, Q, \epsilon) > 0$ as required for convergent stability.
- 705 2. $\pi_0 \leq \rho_1$ and $\pi_{p_0} > \rho_2 > \pi_1 = \alpha$: In this case, $P_{I,1} = 0$ and $P_{I,2} \in (p_0, 1)$. Since
 706 $P_{I,1} = 0$ is a CSNE in $[0, p_0]$, we know that for any $\epsilon \in [0, 1)$ and $0 < P < Q \leq p_0$,
 707 $\Delta E > 0$. In particular, for $\epsilon = 0$, and $0 < P < Q = p_0$ we get $[\pi_{p_0} + \psi_{p_0}r - r/a] =$
 708 $\frac{\Delta E}{(P-Q)} < 0$. But, from $P_{I,2} \in (p_0, 1)$ we can similarly get (for $\epsilon = 0$, $p_0 = Q < P < P_{I,2}$)
 709 $[\pi_{p_0} + \psi_{p_0}r - r/a]a = \frac{\Delta E}{(P-Q)} > 0$, a contradiction.
- 710 3. $\pi_0 \leq \rho_1$ and $\alpha = \pi_1 \geq \rho_2 > 0$: Here, $P_{I,1} = 0$ and $P_{I,2} = 1$. Since $P_{I,1} = 0$ is a CSNE
 711 in $[0, p_0]$, $\forall \epsilon \in [0, 1)$ and $0 \leq P < Q \leq p_0$, $\Delta E > 0$. In particular, for $\epsilon = 0$, and
 712 $0 < P < Q = p_0$ we get $[\pi_{p_0} + \psi_{p_0}r - r/a] = \frac{\Delta E}{(P-Q)} < 0$. Similarly, since $P_{I,2} = 1$ is a
 713 CSNE in $[p_0, 1]$, $\forall \epsilon \in [0, 1)$ and $p_0 \leq Q < P \leq 1$, $\Delta E > 0$. In particular, for $\epsilon = 0$,
 714 and $p_0 = Q < P \leq 1$ we get $[\pi_{p_0} + \psi_{p_0}r - r/a] = \frac{\Delta E}{(P-Q)} > 0$, which is a contradiction.
- 715 4. $\rho_2 \geq \pi_{p_0} \geq \rho_1$: In this case, simple algebra gives $\rho_1 = \rho_2 = \pi_{p_0}$ and $P_{I,1} = p_0$ and
 716 $P_{I,2} = p_0$. Thus, it follows that p_0 is the unique CSNE in the interval $[0, 1]$.
- 717 5. $\pi_{p_0} > \rho_1$ and $\alpha = \pi_1 \geq \rho_2$: In a manner analogous to the first case, here, $p_1 = 1$ is the
 718 unique CSNE.
- 719 6. $\pi_{p_0} > \rho_1$ and $\pi_{p_0} > \rho_2 > \pi_1 = \alpha$: , $P_{I,1} = p_0$ and $P_{I,2} \in (p_0, 1)$. $P_{I,1} = p_0$ cannot
 720 be a Nash equilibrium since for $p_0 < Q < P_{I,2}$, and any $\epsilon \in [0, 1)$, $\Delta E < 0$ since $P_{I,2}$
 721 is the unique CSNE in $[p_0, 1]$. To show that $P_{I,2}$ is a Nash equilibrium, fix $\epsilon \in [0, 1)$
 722 and $P = P_{I,2}$ and let $Q < P_{I,2}$. Note that $\pi_p > \rho_1 \quad \forall p \in [0, p_0]$ and that because π_p
 723 is decreasing and $\pi_{P_{I,2}} = \rho_2$, $\pi_p > \rho_2 \quad \forall p < P_{I,2}$. Thus $\pi_p > \rho(p) \quad \forall p < P_{I,2}$ and so in
 724 particular, $\Delta E > 0$ and $P_{I,2}$ is the unique Nash equilibrium. To see that $P_{I,2}$ is also
 725 convergently stable, we must only show that for any P and Q , $0 \geq Q < p_0 < P < P_{I,2}$
 726 and $\epsilon \in [0, 1)$, we have $\Delta E > 0$. But under these conditions, $p < P_{I,2}$ and so again
 727 $\pi_p > \rho(p)$, which implies $\Delta E > 0$, as required.

728 7. $\pi_0 > \rho_1 > \pi_{p_0}$ and $\pi_{p_0} < \rho_2$: Now, $P_{I,1} \in (0, p_0)$ and $P_{I,2} = p_0$. Similar to the above
 729 case, this implies that $P_{I,1}$ is the unique CSNE (given by Equation (S103)).

730 8. $\pi_0 > \rho_1 > \pi_{p_0}$ and $\alpha = \pi_1 \geq \rho_2$: Simple algebra shows that this case is impossible:

$$731 \quad \rho_1 > \pi_{p_0} = \frac{\gamma}{\gamma + \phi_{\text{prev}}}$$

$$732 \quad r(1-a)(\gamma + \phi_{\text{prev}}) > a(1-r)\gamma$$

$$733 \quad r((1-a)\phi_{\text{prev}} + \gamma) > a\gamma$$

734 but

$$735 \quad \rho_2 \leq \pi_1 < \pi_{p_0} = \frac{\gamma}{\gamma + \phi_{\text{prev}}}$$

$$736 \quad r(\gamma + \phi_{\text{prev}}) < a(\phi_{\text{prev}}r + \gamma)$$

$$737 \quad r((1-a)\phi_{\text{prev}} + \gamma) < a\gamma.$$

738 9. $\pi_0 > \rho_1 > \pi_{p_0}$ and $\pi_{p_0} > \rho_2 > \pi_1$: The reasoning applied to show that the case above
 739 is impossible also rules this case out.

740 We conclude that in all cases there is a unique CSNE, which we denote by p_i .

743 G.2 Vaccination rate \propto incidence

744 Since $\partial_p \psi_p < 0$ and $\partial_p \pi_p < 0$, $\partial_p (\pi_p + r\psi_p) < 0$. Thus, an identical argument to the one
 745 given in Appendix G.3 allows us to show that there is always a CSNE for this model. In
 746 particular, there are three possibilities:

- 747 • If $\pi_0 + r\psi_0 \leq r/a$ then $p_i = 0$ is a unique CSNE.
- 748 • If $\alpha = \pi_1 + r\psi_1 \geq r/a$, then $p_i = 1$ is a unique CSNE.
- 749 • If $\pi_0 + r\psi_0 > r/a > \pi_1 + r\psi_1 = \alpha$ then there is a unique CSNE, $p_i \in (0, 1)$ such that
 750 $\pi_{p_i} + r\psi_{p_i} = r/a$. To simplify this last condition, we use Equations (S56) to obtain

$$751 \quad 0 = \pi_{p_i} + \psi_{p_i}r - \frac{r}{a} = \frac{a(\beta + r(1-\alpha)\phi_{\text{inc}}) - r(\beta + \phi_{\text{inc}})}{a(\beta + \phi_{\text{inc}})} - \frac{\beta + r\phi_{\text{inc}}}{\beta + \phi_{\text{inc}}} \frac{S_\infty}{1 - p_i}$$

752 which is equivalent to

$$753 \quad \frac{S_\infty}{1 - p_i} = \frac{a(\beta + \alpha\phi_{\text{inc}} + r(1-\alpha)\phi_{\text{inc}}) - r(\beta + \phi_{\text{inc}})}{a(\beta + r\phi_{\text{inc}})}. \quad (\text{S110})$$

754 Plugging Equation (S110) into Equation (S50) and rearranging gives the individual
 755 equilibrium,
 756

$$757 \quad p_i = 1 + \frac{a\gamma(\beta + r\phi_{\text{inc}})}{r(\beta + \phi_{\text{inc}})(\beta + a\alpha\phi_{\text{inc}})}$$

$$758 \quad \times \ln \left(\frac{a(\beta + \alpha\phi_{\text{inc}} + r(1-\alpha)\phi_{\text{inc}}) - r(\beta + \phi_{\text{inc}})}{a(1-\alpha)(\beta + r\phi_{\text{inc}})} \right). \quad (\text{S111})$$

760 G.3 Vaccination rate \propto proportion still susceptible

761 Recall that for this model, $\psi_p = 1 - \pi_p$ (see Appendix E.3) and that we assume π_p decreases
 762 with p (as stated in §5.2). Thus, $\partial_p(\pi_p + r\psi_p) = \partial_p((1-r)\pi_p) < 0$, and we can infer that:

- 763 • If $\pi_0 + r\psi_0 \leq r/a$ then $\pi_p + r\psi_p < r/a \quad \forall p > 0$. Hence $p_i = 0$ is the unique Nash
 764 equilibrium. From reasoning similar to that given in Appendix G.1.1, p_i is convergently
 765 stable.
- 766 • If $\alpha = \pi_1 + r\psi_1 \geq r/a$, then $\pi_p + r\psi_p > r/a \quad \forall p < 1$. It follows that $p_i = 1$ is the
 767 unique Nash equilibrium and that it is convergently stable. The condition is equivalent
 768 to $a\alpha r_i \geq r_v$, which is quite intuitive: $a\alpha r_i$ is a lower bound on the cost for a delayer
 769 (attained when there are no new cases after the initial outbreak). If even this lower
 770 bound is higher than the cost of vaccinating, then clearly everyone should vaccinate.
- 771 • If $\pi_0 + r\psi_0 > r/a > \pi_1 + r\psi_1 = \alpha$ then there is a unique CSNE, $p_i \in (0, 1)$ such that
 772 $\pi_{p_i} + r\psi_{p_i} = r/a$.

773 Thus, there is always a CSNE for this model.

774 G.4 Instantaneous vaccination of a proportion ϕ_{inst} of the popu- 775 lation

776 In this case, $\psi_p = \phi_{\text{inst}}(1 - \alpha)$ and $\pi_p = 1 - \psi_p - \frac{S_\infty}{1-p}$. Thus,

$$777 \quad \Delta E = [\pi_p + \phi_{\text{inst}}(1 - \alpha)r - r/a]a(P - Q). \quad (\text{S112})$$

778 Letting $\rho = r/a - r\phi_{\text{inst}}(1 - \alpha)$, since π_p decreases monotonically with p , we can use an
 779 argument similar to those used for the models considered above to show that:

- 780 • if $\pi_0 \leq \rho$ the $p_i = 0$ is the unique CSNE.
- 781 • if $\pi_1 \geq \rho$ then $\pi_p > \rho$ for any $p \in [0, 1)$ and so $p_i = 1$ is the unique CSNE. Rearranging
 782 the condition $\pi_1 \geq \rho$ gives $a\alpha + ra\phi_{\text{inst}}(1 - \alpha) \geq r$. This admits a simple biological
 783 interpretation: $a\alpha + ra\phi_{\text{inst}}(1 - \alpha)$ is the relative risk of delaying when the epidemic does
 784 not successfully spread (that is, one can only be infected during the initial outbreak).
 785 Thus, if the risks of delaying are greater than vaccinating even if the disease does not
 786 spread beyond the cohort initially infected in the outbreak, then it is worthwhile for
 787 individuals to vaccinate pre-emptively.
- 788 • if $\pi_0 > \rho > 0$ then there is a unique CSNE, $p_i \in (0, 1)$ such that $\pi_{p_i} = \rho$. In order to
 789 find p_i explicitly, we use Equation (S82) and substitute $\pi_{p_i} = r/a - r\phi_{\text{inst}}(1 - \alpha)$:

$$790 \quad r/a - r\phi_{\text{inst}}(1 - \alpha) = -\frac{\gamma}{\beta(1 - p_i)} \ln \left(\frac{1 - r/a - (1 - r)\phi_{\text{inst}}(1 - \alpha)}{(1 - \alpha)(1 - \phi_{\text{inst}})} \right),$$

791 and consequently,

$$792 \quad p_i = 1 + \frac{a\gamma}{r\beta(1 - a\phi_{\text{inst}}(1 - \alpha))} \ln \left(\frac{a(1 - (1 - r)\phi_{\text{inst}}(1 - \alpha)) - r}{a(1 - \alpha)(1 - \phi_{\text{inst}})} \right). \quad (\text{S113})$$

793 **G.5 constant rate vaccination**

794 As in [7], we assume π_p is a decreasing function of p . The analysis (performed originally
795 in [7]) is then identical to Appendix G.1.1, implying the existence of a unique CSNE, which
796 we denote by p_i . Using the definition of ρ_1 given in Equation (S102), we have

- 797 • if $\pi_1 \geq \rho_1$ then $p_i = 1$.
- 798 • if $\pi_1 < \rho_1 < \pi_0$ then p_i is the unique solution of $\pi_{p_i} = \rho_1$
- 799 • if $\rho_1 \geq \pi_0$ then $p_i = 0$.

800 **H The group optimum**

801 We have obtained an analytical formula for the group optimum (defined in § 4), for one
802 sub-case of one of our models. The calculation is given below.

803 **H.1 Vaccination rate \propto disease prevalence**

804 We consider the case when $p_m \geq 1$. Recall that if $p_m \geq 1$, then $S_\infty = 0$, $\psi_p = \frac{\phi_{\text{prev}}}{\beta(1-p)} \ln \left(\frac{\beta}{\phi_{\text{prev}}} S(0) + 1 \right)$,
805 and $\pi_p = 1 - \psi_p$ (see Appendix E.1). Thus,

$$\begin{aligned}
 806 \quad C(p) &= rp + (1-p)a(1 - (1-r)\psi_p) \\
 807 \quad &= rp + (1-p)a \left(1 - (1-r) \frac{\phi_{\text{prev}}}{\beta(1-p)} \ln \left(\frac{\beta}{\phi_{\text{prev}}} S(0) + 1 \right) \right) \\
 808 \quad &= rp + a \left((1-p) - (1-r) \frac{\phi_{\text{prev}}}{\beta} \ln \left(\frac{\beta}{\phi_{\text{prev}}} S(0) + 1 \right) \right) \\
 809 \quad C(0) &= a(1 - (1-r) \frac{\phi_{\text{prev}}}{\beta} \ln \left(\frac{\beta}{\phi_{\text{prev}}} (1-\alpha) + 1 \right)) \\
 810 \quad C(1) &= r \\
 811 \quad C'(p) &= r + a \left(-1 - (1-r) \frac{\phi_{\text{prev}}}{\beta} \left(-\frac{\beta}{\phi_{\text{prev}}} (1-\alpha) \right) \frac{1}{\frac{\beta}{\phi_{\text{prev}}} S(0) + 1} \right) \\
 812 \quad &= r + a \left(\frac{\phi_{\text{prev}}(1-\alpha)(1-r)}{\beta S(0) + \phi_{\text{prev}}} - 1 \right) \tag{S114} \\
 813
 \end{aligned}$$

814 Note that $C'(p)$ increases with p since $S(0)$ decreases with p and critical points of $C(p)$ are
815 minima. Thus, if there is a critical point within $[0, 1]$, then it is the global minimum; other-
816 wise, the global minimum is at $C(0)$. To find critical points, set $C'(p) = 0$ or equivalently,

$$817 \quad \frac{r}{a} = 1 - \frac{\phi_{\text{prev}}(1-\alpha)(1-r)}{\beta S(0) + \phi_{\text{prev}}} \tag{S115}$$

818 which can only happen if $r < a$ (since $\alpha < 1$ and $r < 1$), that is, the relative risk (of
819 vaccination *versus* infection) is less than the probability of an outbreak. If $r \geq a$ then

820 $C'(p) \geq 0$ throughout $[0, 1]$ and the minimal group cost $C(p)$ is attained at $p_g = 0$. This is
821 easily explained: If $r_v \geq ar_i$ then the mortality risks from vaccination are no less than those
822 of dying in an outbreak. In this case, vaccinating is not worthwhile for either the individual
823 or the group. We now solve Equation (S115) for the initial coverage p at the critical point
824 of the group cost, $C(p)$, assuming $r < a$:

$$\begin{aligned}
825 \quad \beta S(0) &= \frac{\phi_{\text{prev}}(1-\alpha)a(1-r)}{a-r} - \phi_{\text{prev}} \\
826 \quad 1-p &= \frac{\phi_{\text{prev}}}{\beta(1-\alpha)} \left(\frac{(1-\alpha)a(1-r)}{a-r} - 1 \right) \\
827 \quad p &= 1 - \frac{\phi_{\text{prev}}}{\beta(1-\alpha)} \left(\frac{(1-a)r - (1-r)\alpha a}{a-r} \right) \tag{S116} \\
828
\end{aligned}$$

829 The critical point is attained at $p \geq 1$ if and only if $(1-a)r \leq (1-r)\alpha a$, which is
830 equivalent to

$$831 \quad r_v \leq a(r_i\alpha + r_v(1-\alpha)), \tag{S117}$$

832 and in this case the group optimum is vaccinating the entire population ($p_g = 1$). Biologi-
833 cally, Equation (S117) means that more people are expected to die if, in case of an outbreak,
834 all individuals not infected initially are vaccinated (discounted by the outbreak probability,
835 a), than the number of people expected to die if the entire population is vaccinated pre-
836 emptively. To see this, note the probability of death due to vaccinating is r_v (the left hand
837 side of Equation (S117)). To interpret the right hand side of Equation (S117), consider an
838 individual who is not vaccinated pre-emptively. If there is an outbreak (represented by the
839 factor a), the first term in brackets ($r_i\alpha$) represents the probability of being in the initially in-
840 fected cohort (α), and then dying due to the disease. The second term ($r_v(1-\alpha)$), represents
841 the probability of not being in the initially infected cohort, and dying due to the vaccine side
842 effects. Note that because in this scenario $S_\infty = 0$, no delayers remain susceptible (they are
843 either infected or vaccinated). Thus, any individual who is not pre-emptively vaccinated,
844 and who is not in the initially infected cohort ($1-\alpha$) has either a probability r_v of dying
845 due to vaccine side effects, or a probability r_i of dying due to the disease. But since $r_v < r_i$,
846 the term $r_v(1-\alpha)$ is a lower bound on the probability of death for an individual who is
847 susceptible immediately after the outbreak is seeded (that is, not pre-emptively vaccinated,
848 or in the cohort initially infected at the beginning of the outbreak).

849 Note that not vaccinating anyone pre-emptively is the group optimum ($p_g \leq 0$) iff

$$850 \quad \beta(1-\alpha)(a-r) \leq \phi_{\text{prev}}((1-a)r - (1-r)\alpha a), \tag{S118}$$

851 (but this condition is difficult to interpret biologically).

852 Lastly, if $\phi_{\text{prev}} \leq \beta \frac{(a-r)(1-\alpha)}{r(1-a(1-\alpha))-a\alpha}$ and $(1-a)r > (1-r)\alpha a$, then $p_g \in [0, 1]$ and is given
853 by Equation (S116). It is thus interesting to note that p_g depends piece-wise linearly on
854 ϕ_{prev} .

855 **References**

- 856 [1] Kermack WO, McKendrick AG. A contribution to the mathematical theory of epidemics.
857 Proceedings of the Royal Society of London Series A. 1927;115:700–721.
- 858 [2] Ma J, Earn DJD. Generality of the final size formula for an epidemic of a newly invading
859 infectious disease. *Bulletin of Mathematical Biology*. 2006;68(3):679–702.
- 860 [3] Earn DJD, Andrews PW, Bolker BM. Population-level effects of suppressing fever. Proc
861 R Soc Lond B. 2014;281(1778):20132570.
- 862 [4] Weisstein EW. Lambert W-Function. From MathWorld –A Wolfram Web Resource.
863 <http://mathworld.wolfram.com/LambertW-Function.html>;
- 864 [5] Corless RM, Gonnet GH, Hare DEG, Jeffrey DJ, Knuth DE. On the Lambert W function.
865 *Advances in Computational Mathematics*. 1996;5(4):329–359.
- 866 [6] Hirsch MW, Smale S. *Differential equations, dynamical systems, and linear algebra*.
867 Academic Press, New York; 1974.
- 868 [7] Bauch CT, Galvani AP, Earn DJD. Group interest versus self-interest in smallpox vac-
869 cination policy. *PNAS – Proceedings of the National Academy of Sciences of the U.S.A.*.
870 2003;100(18):10564–10567.