

FOUR-MANIFOLDS, TWO-COMPLEXES AND THE QUADRATIC BIAS INVARIANT

IAN HAMBLETON AND JOHN NICHOLSON

ABSTRACT. Kreck and Schafer produced the first examples of stably diffeomorphic closed smooth 4-manifolds which are not homotopy equivalent. They were constructed by applying the doubling construction to 2-complexes over certain finite abelian groups of odd order. By extending their methods, we formulate a new homotopy invariant on the class of 4-manifolds arising as doubles of 2-complexes with finite fundamental group. As an application we show that, for any $k \geq 2$, there exist a family of k closed smooth 4-manifolds which are all stably diffeomorphic but are pairwise not homotopy equivalent.

1. INTRODUCTION

Two closed smooth 4-manifolds M, N are said to be *stably diffeomorphic* if there exists $r \geq 0$ and a diffeomorphism $M \# r(S^2 \times S^2) \cong N \# r(S^2 \times S^2)$. Kreck's modified surgery [41] gives techniques to classify 4-manifolds up to stable diffeomorphism, and these methods have been applied to study manifolds over a range of fundamental groups [23–26, 28, 29, 34, 38].

The existence of exotic smooth structures shows that simply-connected oriented 4-manifolds which are stably diffeomorphic need not be diffeomorphic, but it follows from results of Donaldson [16] and Wall [64] that such 4-manifolds are h -cobordant and hence homotopy equivalent.

In contrast, Kreck-Schafer [42] produced the first examples of closed smooth 4-manifolds which are stably diffeomorphic, but not even homotopy equivalent. Their examples arose from the following *doubling construction*: for a finite 2-complex X , let $M(X)$ be the boundary of a smooth regular neighbourhood of an embedding $X \hookrightarrow \mathbb{R}^5$ (see Section 5 for more details). The construction has the following properties (see [42, Section 2]):

- (i) If X and Y are finite 2-complexes such that $X \simeq Y$ are homotopy equivalent, then $M(X)$ and $M(Y)$ are h -cobordant, hence homotopy equivalent.
- (ii) If $\chi(X) = \chi(Y)$ and $\pi_1(X) \cong \pi_1(Y)$, then $M(X)$ and $M(Y)$ are stably diffeomorphic.

Their main result was that there exist pairs of finite 2-complexes X, Y with $\chi(X) = \chi(Y)$ and $M(X) \not\simeq M(Y)$ such that $\pi_1(X) \cong \pi_1(Y)$ is elementary abelian of odd order. To achieve this, they defined a homotopy invariant (in $\mathbb{Z}/2$) for doubles $M(X)$ with $\pi_1(X)$ finite of odd order.

In this paper we will define and study the *quadratic bias invariant*, which generalises the invariant of Kreck-Schafer. Let G be a finite group and X a finite 2-complex with $\pi_1(X) \cong G$ which is *minimal* in the sense that $\chi(X)$ is minimal over such complexes. The *bias invariant* was defined by Metzler [47] to be a class $\beta(X)$ in an abelian group $B(G) := (\mathbb{Z}/m)^\times / \langle \pm D(G) \rangle$ where $m = m_G$ (Definition 3.4) and $D(G)$ is the image of a certain map $\varphi: \text{Aut}(G) \rightarrow (\mathbb{Z}/m)^\times / \{\pm 1\}$ (Definition 3.17). The quadratic bias invariant will be a class $\beta_Q(M(X))$ in a quotient group $B_Q(G)$ of $B(G)$ (Definition 6.10). Let $\mathcal{M}_4(G)$ denote the set of homotopy types of 4-manifolds $M(X)$ where X is a minimal finite 2-complex with $\pi_1(X) \cong G$. We will show:

Theorem A. *The quadratic bias invariant is a homotopy invariant. In particular, for a finite group G , the quadratic bias invariant defines a map*

$$\beta_Q: \mathcal{M}_4(G) \rightarrow B_Q(G).$$

Furthermore, $\beta_Q(M(X)) = q(\beta(X))$ where $q: B(G) \twoheadrightarrow B_Q(G)$ is the natural surjection.

The extent to which the quadratic bias can be used to distinguish manifolds depends on choosing the quotient $B_Q(G)$ of $B(G)$ so that $\beta_Q(M(X))$ is both a homotopy invariant and can

Date: November 20, 2025.

2020 Mathematics Subject Classification. Primary 57K41, 57N65; Secondary 20F05, 55U15, 57R67.

also be computed in non-trivial examples. Sections 6 and 7 are directed towards this goal. A description of the subgroup $N(G) = \ker(q: B(G) \rightarrow B_Q(G))$ can be found in Remark 6.11 (ii).

If $H_2(G; \mathbb{Z})$ has a certain special form we can explicitly compute the quadratic bias obstruction group $B_Q(G)$. Recall that a finite group G is *efficient* if $\chi_{\min}(G) = 1 + d(H_2(G; \mathbb{Z}))$, where $\chi_{\min}(G)$ denotes the minimal Euler characteristic of a finite 2-complex X with $\pi_1(X) \cong G$, and $d(\cdot)$ denotes the minimal number of generators of a group [32].

Theorem B. *Let G be a finite group such that $H_2(G; \mathbb{Z}) \cong (\mathbb{Z}/m)^d$ for some $m \geq 1$, $d \geq 3$. If G is efficient, then there is an isomorphism*

$$B_Q(G) \cong \frac{(\mathbb{Z}/m)^\times}{\pm(\mathbb{Z}/m)^{\times 2} \cdot D(G)}$$

where $D(G) = \text{im}(\varphi_G: \text{Aut}(G) \rightarrow (\mathbb{Z}/m)^\times / \{\pm 1\})$. If G is not efficient, then $B_Q(G) = 0$.

It is known that finite abelian groups are efficient [57, Proposition 5], but Swan constructed a group of the form $G \cong (\mathbb{Z}/7)^3 \rtimes \mathbb{Z}/3$ which is not efficient [60, p196].

Remark 1.1. We would expect the structure of $B_Q(G)$ to be much more complicated in general. Its definition involves new ideas related to the unitary isometries of multi-scaled hyperbolic forms arising from the decomposition of $H_2(G; \mathbb{Z})$ into cyclic factors (see Proposition 4.15).

Using Theorems A and B, we are now able to effectively compare closed smooth 4-manifolds of the form $M(X)$ up to homotopy equivalence. This allows us to establish the following, which answers a recent question of Kasprowski-Powell-Ray in the affirmative [36, Question 1.5].

Theorem C. *For each $k \geq 2$, there exist closed smooth 4-manifolds M_1, M_2, \dots, M_k which are all stably diffeomorphic but not pairwise homotopy equivalent.*

These examples can be taken to be stably parallelisable and have finite abelian fundamental groups of the form $G = (\mathbb{Z}/m)^d$, for any $d \geq 3$ odd, and any $m \geq 3$ with sufficiently many distinct prime factors. This generalises the case $k = 2$ which was established by Kreck-Schafer [42] over fundamental groups of the form $(\mathbb{Z}/p)^d$ for a prime $p \equiv 1 \pmod{4}$. We also construct examples over non-elementary abelian groups $(\mathbb{Z}/m)^d \times \mathbb{Z}/t$ where $d \geq 4$ is even and certain values of m and t with $m \neq t$.

Remark 1.2. The result of Theorem C is optimal for manifolds with finite fundamental group, since by [26, Corollary 1.5] there are only finitely many homeomorphism types of closed 4-manifolds with a given finite fundamental group and Euler characteristic. It remains open whether there exists an infinite collection of such manifolds with arbitrary fundamental group.

To obtain examples over other fundamental groups, we require a pair of finite 2-complexes X, Y with finite fundamental group G and $\chi(X) = \chi(Y)$ but which are not homotopy equivalent. Such examples have previously only been known to exist when G is either a finite abelian group [47, 57] or a group with periodic cohomology [18, 50, 52]. However, if G has periodic cohomology, then $H_2(G; \mathbb{Z}) = 0$ [61, Corollary 2] and so the bias invariant contains no information.

In spite of this, we will establish the following (see Theorem 9.5). This serves to demonstrate that the quadratic bias invariant is computable for non-abelian fundamental groups.

Theorem D. *Let $G = Q_8 \times (\mathbb{Z}/p)^3$ where p is a prime such that $p \equiv 1 \pmod{8}$. Then:*

- (i) *There exist minimal finite 2-complexes X, Y with fundamental group G which are homotopically distinct.*
- (ii) *There exist closed smooth 4-manifolds M, N with fundamental group G which are stably diffeomorphic but not homotopy equivalent.*

This also gives the first example of a non-abelian finite group G which does not have periodic cohomology such that there exists homotopically distinct finite 2-complexes X, Y with fundamental group G and $\chi(X) = \chi(Y)$. Part (ii) follows from (i) by taking $M = M(X)$, $N = M(Y)$ and applying the conditions of Theorem B to show that $\beta_Q(M) \neq \beta_Q(N)$.

A number of interesting questions remain concerning the doubles $M(X)$. If $X \simeq Y$, then $M(X)$ and $M(Y)$ are h -cobordant. More generally, we ask:

Question 1.3. *If $X \simeq Y$, then are $M(X)$ and $M(Y)$ diffeomorphic?*

An important special case is when X is a point and Y is the presentation 2-complex of a potential counterexample to the Andrews-Curtis conjecture [3]. In this case we have $X \simeq Y$, $M(X) = S^4$, $M(Y) \simeq S^4$ and so Question 1.3 is equivalent to the question of whether $M(Y)$ is an exotic 4-sphere. Such examples were considered by Akbulut-Kirby in [2] and, for one such example, Question 1.3 was shown by Gompf to have an affirmative answer [21].

It was recently shown by Freedman-Krushkal-Lidman that all Seiberg-Witten invariants vanish for doubles $M(X)$ [20, Proposition 1.3]. We could also ask Question 1.3 for simple homotopy equivalent 2-complexes, since in that case $M(X)$ and $M(Y)$ are s -cobordant [42, p15].

1a. Comparison with quadratic 2-type. In [6, 26], it was shown that a closed topological 4-manifold M with $\pi_1(M)$ finite of odd order is determined up to homotopy equivalence by its quadratic 2-type

$$Q(M) = [\pi_1(M), \pi_2(M), k_M, S_M]$$

where $k_M \in H^3(\pi_1(M); \pi_2(M))$ denotes the k -invariant and $S_M: \pi_2(M) \times \pi_2(M) \rightarrow \mathbb{Z}[\pi_1(M)]$ denotes the equivariant intersection form. An isometry \cong of two such quadruples is an isomorphism of pairs π_1, π_2 respecting the k -invariant and inducing an isometry on S . For arbitrary finite fundamental groups, there is a possibly non-zero additional invariant which lies in $\text{Tors}(\Gamma(\pi_2(M)) \otimes_{\mathbb{Z}G} \mathbb{Z})$. However, we will show (see Theorem 6.16):

Theorem 1.4. *Let G be a finite group and let $M_1, M_2 \in \mathcal{M}_4(G)$. If $Q(M_1) \cong Q(M_2)$ are isometric, then $\beta_Q(M_1) = \beta_Q(M_2)$.*

By Theorem C and [6, 26, 35, 37], there are examples of stably diffeomorphic homotopy distinct 4-manifolds M_1, \dots, M_k such that $\text{Tors}(\Gamma(\pi_2(M_i)) \otimes_{\mathbb{Z}G} \mathbb{Z}) \neq 0$. It is open whether this additional invariant is needed in order to determine such manifolds up to homotopy equivalence.

Although $\beta_Q(M)$ is determined by $Q(M)$, it is not immediately clear how to compute $\beta_Q(M)$ from $Q(M)$. It is also not clear how to use $Q(M)$ directly in order to establish quantitative results such as the ones given in Theorems C and D (ii). The following is currently open:

Question 1.5. *Do there exist closed smooth 4-manifolds M, N with fundamental group G which are stably diffeomorphic but such that $\pi_2(M)$ and $\pi_2(N)$ are not $\text{Aut}(G)$ -isomorphic?*

Similarly it is not known whether such manifolds exist such that S_M and S_N are not isometric modulo the action of $\text{Aut}(G)$. See Section 2 for the definition of $\text{Aut}(G)$ -isomorphic.

1b. Results in higher dimensions. In Definition 6.10, the invariant β_Q is generalised to the doubles of finite (G, n) -complexes (see Section 2). We define the quadratic bias invariant

$$\beta_Q: \mathcal{M}_{2n}(G) \rightarrow B_Q(G, n)$$

for all $n \geq 2$ where $\mathcal{M}_{2n}(G)$ is the set of homotopy types of doubles of minimal finite (G, n) -complexes. We obtain an analogue of Theorem A (see Theorem 6.12) as well as an analogue of Theorem B (see Theorem 7.1) when G is a finite group such that $H_n(G; \mathbb{Z}) \cong (\mathbb{Z}/m)^d$ for some $m \geq 1$, $d \geq 3$ and (G, n) satisfies the *strong minimality hypothesis* (see Definition 3.1). The result is similar to Theorem B for n even, but we show that $B_Q(G, n) = 0$ for n odd.

We also obtain explicit examples in higher dimensions, complementing the results of Conway-Crowley-Powell-Sixt [11, 12] which for dimensions $4n > 4$ were either simply connected with $H_2(M) \neq 0$, or had infinite fundamental group $\pi_1(M) \cong \mathbb{Z}$. See Theorems 9.1 and 9.2.

Theorem 1.6. *For all $n \geq 2$ even, and all $k \geq 2$, there exist closed smooth $2n$ -manifolds M_1, M_2, \dots, M_k with non-trivial finite fundamental group which are all stably diffeomorphic, but not pairwise homotopy equivalent. Furthermore, for the case $k = 2$, the manifolds can be taken to have isometric equivariant intersection forms.*

Remark 1.7. This result also addresses two gaps in the paper of Kreck-Schafer [42], where this result was given in the case $k = 2$. Firstly, the proof that examples exist in the case $n > 2$ is incomplete since it relies on a formula from [57, Proposition 8] which is incorrect, as pointed out in [46, p305] (see [42, p36-38]).

Secondly, the examples constructed by [42] were claimed to have isometric equivariant intersection form (see [42, p21]). The equivariant intersection forms were shown to be hyperbolic, but the possibility that they were hyperbolic over non-isomorphic $\mathbb{Z}G$ -modules was not considered. This also affects the remark after [27, Theorem 4.1] and was mentioned in [11, p2].

In the case $n = 2$, since the examples in Theorem 1.6 are distinguished by the quadratic bias invariant (see Section 9), it follows from Theorem 1.4 that their quadratic 2-types are not isometric. Hence the closed smooth 4-manifolds M_1 and M_2 with isometric equivariant intersection form that we construct are distinguished by their k -invariants. This gives the first examples of stably diffeomorphic closed smooth 4-manifolds which are distinguished up to homotopy equivalence by their k -invariants.

Organisation of the paper. We begin by recalling the necessary background on the bias invariant. We give preliminaries on CW-complexes (Section 2), then define the bias invariant in the setting of finite (G, n) -complexes and establish its main properties (Section 3).

We next introduce the quadratic bias invariant. We give preliminaries on hermitian forms (Section 4) and the doubling construction (Section 5). In Section 6, we define the quadratic bias invariant and prove Theorem A (see Theorem 6.12) and Theorem 1.4 (see Theorem 6.16).

Finally, we evaluate the quadratic bias invariant and apply it to examples. We establish Theorem B (see Theorem 7.1), give details concerning the bias invariant for complexes (Section 8), and then prove Theorems C and D (Section 9). Some technical calculations are contained in Appendices A, B and C, which give information about surgery obstruction groups and numerical functions describing the action of $\text{Aut}(G)$ on the polarised quadratic bias invariant.

Conventions. Our rings R have identity, and we work in the category of finitely generated left R -modules and left R -module homomorphisms. We will assume all CW-complexes are connected with basepoint, and maps between CW-complexes will be cellular and basepoint-preserving. Manifolds will be assumed to be smooth, closed, oriented and connected.

Acknowledgements. IH was partially supported by an NSERC Discovery Grant. JN was supported by the Heilbronn Institute for Mathematical Research and a Rankin-Sneddon Research Fellowship from the University of Glasgow. We gratefully acknowledge that this research was partially supported by the Fields Institute for Research in Mathematical Sciences in April 2023, the Isaac Newton Institute in May 2024, the University of Glasgow in June 2024, and the Heilbronn Institute in November 2024. The authors would like to thank Diarmuid Crowley, Jens Harlander, Daniel Kasprowski and Mark Powell for helpful conversations. We would also like to thank the referee for many useful suggestions which improved the exposition of this article.

CONTENTS

1. Introduction	1
2. Preliminaries on CW-complexes	5
3. The bias invariant for (G, n) -complexes	6
4. Preliminaries on hermitian forms	10
5. The doubling construction	16
6. The quadratic bias invariant	19
7. Evaluation of the quadratic bias obstruction group	25
8. Examples of homotopy inequivalent (G, n) -complexes	30
9. Examples of homotopy inequivalent doubled (G, n) -complexes	32
Appendix A. Unitary groups and odd-dimensional L -theory	37
Appendix B. Computations of odd-dimensional L -theory of abelian groups	39
Appendix C. The numerical functions $e(d, n)$ and $s(d, n)$	41
References	43

2. PRELIMINARIES ON CW-COMPLEXES

We begin by establishing some conventions and definitions. Let G be a group. A (G, n) -complex is an n -dimensional CW-complex X such that $\pi_1(X) \cong G$ and \tilde{X} is $(n-1)$ -connected. Equivalently, it is the n -skeleton of a $K(G, 1)$ -space. Note that a $(G, 2)$ -complex is equivalently a 2-complex X with $\pi_1(X) \cong G$. We say a group G has type F_n if there exists a finite (G, n) -complex. In particular, a group has type F_2 if and only if it is finitely presented. If G is a finite group, then G has type F_n for all $n \geq 1$.

For a group G , a G -polarised space is a pair (X, ρ_X) where X is a space and $\rho_X: \pi_1(X) \xrightarrow{\cong} G$ is a group isomorphism which we often refer to as a *polarisation*. If $h: X \rightarrow Y$ is a map, then we can view $\pi_1(h) \in \text{Aut}(G)$ using the G -polarisations ρ_X and ρ_Y ; formally, we use $\pi_1(h)$ to denote $\rho_Y \circ \pi_1(h) \circ \rho_X^{-1}$. Two G -polarised spaces X and Y are said to be *polarised homotopy equivalent* if there exists a homotopy equivalence $h: X \rightarrow Y$ such that $\pi_1(h) = \text{id}_G \in \text{Aut}(G)$. We will assume all (G, n) -complexes are G -polarised.

Let $\text{HT}(G, n)$ denote the set of homotopy types of finite (G, n) -complexes and let $\text{PHT}(G, n)$ the set of polarised homotopy types of finite (G, n) -complexes.

There is an action of $\text{Aut}(G)$ on $\text{PHT}(G, n)$ where $\theta \in \text{Aut}(G)$ maps $(X, \rho) \in \text{PHT}(G, n)$ to $(X, \theta \circ \rho)$. It follows easily that

$$\text{HT}(G, n) \cong \text{PHT}(G, n) / \text{Aut}(G).$$

The following will be our algebraic model for finite (G, n) -complexes. We will view \mathbb{Z} as a $\mathbb{Z}G$ -module with a trivial G -action.

Definition 2.1. Let $n \geq 2$ and let G be a group. An *algebraic n -complex* over $\mathbb{Z}G$ is a chain complex $C = (C_*, \partial_*)$ of (finitely generated) free $\mathbb{Z}G$ -modules C_* equipped with a choice of $\mathbb{Z}G$ -module isomorphism $H_0(C_*) \cong \mathbb{Z}$ such that

- (i) $C_i = 0$ for $i < 0$ or $i > n$.
- (ii) $H_i(C_*) = 0$ for $0 < i < n$.

Let $\text{Alg}(G, n)$ denote the set of algebraic n -complexes over $\mathbb{Z}G$ considered up to the equivalence relation where $C \simeq C'$ if there exists a chain map $f: C \rightarrow C'$ such that $H_0(f) = \text{id}_{\mathbb{Z}}$ and $H_n(f)$ is a $\mathbb{Z}G$ -isomorphism. We refer to this equivalence relation as *chain homotopy equivalence*.

If $C = (C_*, \partial_*) \in \text{Alg}(G, n)$, then define $\chi(C) := \sum_{i=0}^n (-1)^i \text{rank}_{\mathbb{Z}G}(C_i)$ where $\text{rank}_{\mathbb{Z}G}(C_i)$ denotes the rank of C_i as a free $\mathbb{Z}G$ -module. This is a chain homotopy invariant and so does not depend on the choice of representative in $\text{Alg}(G, n)$.

Let $\theta \in \text{Aut}(G)$. If M is a $\mathbb{Z}G$ -module, let M_θ denote the $\mathbb{Z}G$ -module with the same underlying abelian group but with G -action given by $g \cdot m := \theta(g) \cdot m$ for $g \in G$ and $m \in M$. We say two $\mathbb{Z}G$ -modules M and N are *Aut(G)-isomorphic*, written $M \cong_{\text{Aut}(G)} N$, if there is an isomorphism $M \cong N_\theta$ for some $\theta \in \text{Aut}(G)$.

The class of algebraic n -complexes over $\mathbb{Z}G$ admit an action by $\text{Aut}(G)$ (see [53, Section 6]). If $C = (C_*, \partial_*) \in \text{Alg}(G, n)$, then define

$$C_\theta = ((C_n)_\theta \xrightarrow{\partial_n} (C_{n-1})_\theta \xrightarrow{\partial_{n-1}} \cdots \xrightarrow{\partial_2} (C_1)_\theta \xrightarrow{\partial_1} (C_0)_\theta).$$

Since each C_i is a free $\mathbb{Z}G$ -module, we have that $(C_i)_\theta \cong C_i$ (see [53, Lemma 6.1 (i)]) and so $C_\theta \in \text{Alg}(G, n)$. This action is well-defined on chain homotopy types and so induces an action of $\text{Aut}(G)$ on $\text{Alg}(G, n)$ (see [53, Section 6]).

If X is a finite CW-complex, then $C_*(\tilde{X})$ is a chain complex over $\mathbb{Z}[\pi_1(X)]$. Given a G -polarisation $\rho: \pi_1(X) \rightarrow G$, we can then convert this to a chain complex over $\mathbb{Z}G$. We will denote this by $C_*(\tilde{X}, \rho)$ when we want to emphasise the choice of polarisation.

The following two propositions are standard and show that, in order to study finite (G, n) -complexes up to homotopy equivalence, it suffices to study algebraic n -complexes over $\mathbb{Z}G$ up to chain homotopy equivalence, and the action of $\text{Aut}(G)$ on this class. For a convenient reference, see [53, Proposition 5.1 & Lemma 6.2].

Proposition 2.2. *Let $n \geq 2$ and let G be a group of type F_n . Then:*

- (i) If X is a finite (G, n) -complex, and $\rho: \pi_1(X) \rightarrow G$ is a polarisation, then $C_*(\tilde{X}, \rho)$ is an algebraic n -complex over $\mathbb{Z}G$. Furthermore, $\chi(C_*(\tilde{X}, \rho)) = \chi(X)$.
- (ii) If X, Y are finite (G, n) -complexes and $\theta \in \text{Aut}(G)$, then there exists a homotopy equivalence $f: X \rightarrow Y$ such that $\pi_1(f) = \theta$ if and only if there exists a $\mathbb{Z}G$ -chain homotopy equivalence $h: C_*(\tilde{X}) \rightarrow C_*(\tilde{Y})_\theta$.

Proposition 2.3. Let $n \geq 2$ and let G be a group of type F_n . Then the map

$$\mathcal{C}: \text{PHT}(G, n) \rightarrow \text{Alg}(G, n), \quad (X, \rho) \mapsto C_*(\tilde{X}, \rho)$$

is injective. Furthermore, it induces an injective map $\overline{\mathcal{C}}: \text{HT}(G, n) \rightarrow \text{Alg}(G, n)/\text{Aut}(G)$ where the action of $\text{Aut}(G)$ on $\text{Alg}(G, n)$ is as defined above.

Definition 2.4. Let $n \geq 2$. If G has type F_n , define

$$\chi_{\min}(G, n) = \min\{(-1)^n \chi(X) : X \text{ a finite } (G, n)\text{-complex}\}.$$

This value always exists (see, for example, [51, Proposition 2.3 (ii)]). We say that a finite (G, n) -complex X is *minimal* if $(-1)^n \chi(X) = \chi_{\min}(G, n)$, and we let $\text{HT}_{\min}(G, n)$ denote the set of homotopy types of minimal finite (G, n) -complexes. Similarly, let $\text{Alg}_{\min}(G, n)$ denote the set of chain homotopy types of algebraic n -complexes C over $\mathbb{Z}G$ such that $(-1)^n \chi(C) = \chi_{\min}(G, n)$.

3. THE BIAS INVARIANT FOR (G, n) -COMPLEXES

Throughout this section, we will fix an integer $n \geq 2$ and a finite group G . All algebraic n -complexes will be assumed to be over $\mathbb{Z}G$. All group homology groups will be taken to have coefficients in \mathbb{Z} (with the trivial group action) unless otherwise mentioned. We let $d(G)$ denote the minimal number of generators of a group G .

Throughout, we will make use of 0th Tate cohomology. For a finite group G and a $\mathbb{Z}G$ -module A , recall that the corresponding 0th Tate cohomology group is defined to be

$$\hat{H}^0(G; A) := A^G / (N \cdot A)$$

where $N := \sum_{g \in G} g \in \mathbb{Z}G$ denotes the group norm and $N \cdot A := \{N \cdot a \mid a \in A\} \leq A^G$. Then $(\cdot)^\wedge := \hat{H}^0(G; -): \mathbb{Z}G\text{-mod} \rightarrow \text{Ab}$ is a functor from $\mathbb{Z}G$ -modules to abelian groups, and factors through the functor $(\cdot)^G: \mathbb{Z}G\text{-mod} \rightarrow \text{Ab}$.

3a. The strong minimality hypothesis. Since G is finite, $H_n(G)$ is a finite abelian group and so $d(H_n(G)) < \infty$. The following is the geometric analogue of the minimality hypothesis [57].

Definition 3.1. Let $n \geq 2$ and let G be a finite group. We say that the pair (G, n) satisfies the *strong minimality hypothesis* if $\chi_{\min}(G, n) = (-1)^n + d(H_n(G))$.

This can be viewed as a higher dimensional generalisation of efficiency, restricted to the case of finite groups. Recall that a finitely presented group G is *efficient* if

$$\chi_{\min}(G) = 1 - r(H_1(G)) + d(H_2(G)),$$

where $r(A)$ denotes the torsion free rank of A [32, p166]. If G is finite, then $r(H_1(G)) = 0$ and so $(G, 2)$ satisfies the strong minimality hypothesis if and only if G is efficient.

Remark 3.2. For $n \geq 2$ and a finite group G , the pair (G, n) satisfies the *minimality hypothesis* if $\chi_{\min}^{\text{alg}}(G, n) = (-1)^n + d(H_n(G))$, where $\chi_{\min}^{\text{alg}}(G, n) = \min\{(-1)^n \chi(C) : C \in \text{Alg}(G, n)\}$ (see [57, Section 3]). If $n > 2$, then Wall [65, Theorem E] shows that $\chi_{\min}(G, n) = \chi_{\min}^{\text{alg}}(G, n)$ and so the minimality hypothesis and strong minimality hypothesis coincide in this case. If $n = 2$, then $\chi_{\min}^{\text{alg}}(G, 2) \leq \chi_{\min}(G, 2)$ and equality holds if G has the D2 property [65, Section 3], [22], [33]. It is currently not known whether or not $\chi_{\min}^{\text{alg}}(G, 2) = \chi_{\min}(G, 2)$ for all finitely presented groups G .

Note that $\chi_{\min}(G, 2) \geq \chi_{\min}^{\text{alg}}(G, 2) \geq 1 + d(H_2(G))$. Thus, if $(G, 2)$ satisfies the strong minimality hypothesis, then $(G, 2)$ satisfies the minimality hypothesis.

It is known that finite abelian groups and finite p -groups satisfy the minimality hypothesis in all dimensions [57, Proposition 5], but finite p -groups are not known to be efficient in general.

Moreover, Swan [60, p196] constructed a group of the form $G \cong (\mathbb{Z}/7)^3 \rtimes \mathbb{Z}/3$ such that $(G, 2)$ does not satisfy the minimality hypothesis, and so G is not efficient.

We will now give an alternate formulation of the strong minimality hypothesis, which will be useful in defining the bias invariant in Section 3. Define the *invariant rank* to be $r_{(G,n)} := \text{rank}_{\mathbb{Z}}(L^G)$ where $L = \pi_n(X)$ for X any minimal (G, n) -complex. This does not depend on the choice of X . The following can be proven similarly to [57, Proposition 4].

Proposition 3.3. *Let $n \geq 2$ and let G be a finite group. Then $\chi_{\min}(G, n) = (-1)^n + r_{(G,n)}$. In particular, (G, n) satisfies the strong minimality hypothesis if and only if $r_{(G,n)} = d(H_n(G))$.*

If X is a minimal (G, n) -complex and $L = \pi_n(X)$, then it is a consequence of dimension shifting that $H_n(G) \cong \hat{L}$, and so the natural surjection $L^G \twoheadrightarrow \hat{L}$ implies that $r_{(G,n)} \geq d(H_n(G))$.

Definition 3.4. Let $n \geq 2$, let G be a finite group and let $H_n(G) \cong \mathbb{Z}/m_1 \oplus \cdots \oplus \mathbb{Z}/m_r$ where $m_i \mid m_{i+1}$ for all $i \geq 1$ and $r = r_{(G,n)}$. Such a decomposition exists since $r_{(G,n)} \geq d(H_n(G))$, but we need not have $m_i \neq 1$ for all i . Define the *modulus* to be $m_{(G,n)} = m_1$.

By the classification of finite abelian groups, $m_{(G,n)}$ does not depend on the choice of identification of $H_n(G)$. As usual, when $n = 2$, we will write $r_G := r_{(G,2)}$ and $m_G := m_{(G,2)}$.

Remark 3.5. It follows by comparing Definitions 3.1 and 3.4 that $m_{(G,n)} \neq 1$ if and only if (G, n) satisfies the strong minimality hypothesis.

3b. The bias invariant for algebraic n -complexes. The aim of this section will be to define the bias invariant, which was introduced by Metzler in [47]. We will formulate our definitions so that they are analogous to those made in the definition of the quadratic bias invariant in Section 6. Our treatment resembles the one given by Sieradski-Dyer [57, p202] and is the version implicitly used by Kreck-Schafer [42, p34]. A convenient reference is [55, Section 2].

Observe that, if two algebraic n -complexes C_* and D_* are $\mathbb{Z}G$ -chain homotopy equivalent, then $\chi(C_*) = \chi(D_*)$. We will therefore restrict to the case of chain complexes with equal Euler characteristic. The following is [55, Lemma 1 (§2)] specialised to the case $j = 0$ and $G' = G$.

Proposition 3.6. *Let C_* , D_* be algebraic n -complexes such that $\chi(C_*) = \chi(D_*)$ and let $h: C_* \rightarrow D_*$ be a chain map such that $H_0(h) = \text{id}_{\mathbb{Z}}$. Then the map*

$$H_n(h)^\wedge: H_n(C_*)^\wedge \rightarrow H_n(D_*)^\wedge$$

is an isomorphism and is independent of the choice of h . Let this be denoted by $\sigma(C_, D_*)$.*

Since C_* and D_* are algebraic n -complexes, it follows from standard homological algebra that a chain map such that $H_0(h) = \text{id}_{\mathbb{Z}}$ always exists. Hence $\sigma(C_*, D_*)$ is always defined.

Definition 3.7. Fix an algebraic n -complex \overline{C}_* , which we will refer to as the *reference complex*. Let $L := H_n(\overline{C}_*)$, let $\hat{L} = H_n(\overline{C}_*)^\wedge$ and let $\psi: L^G \twoheadrightarrow \hat{L}$ denote the quotient map.

Let C_* be an algebraic n -complex such that $\chi(C_*) = \chi(\overline{C}_*)$ and let

$$\psi_{C_*}: H_n(C_*)^G \twoheadrightarrow H_n(C_*)^\wedge$$

denote the quotient map. Since $L^G \cong H_n(C_*)^G$ and $\hat{L} \cong H_n(C_*)^\wedge$ are isomorphic as abelian groups, we can find isomorphisms $\bar{\tau}_{C_*}$ and τ_{C_*} such that $\psi_{C_*} \circ \bar{\tau}_{C_*} = \tau_{C_*} \circ \psi$. We do this by choosing bases for L^G and $H_n(C_*)^G$ whose images under the quotient maps give generating sets for invariant factor decompositions of the finite abelian groups \hat{L} and $H_n(C_*)^\wedge$ (see Remark 3.12 for more details). The reference maps $\bar{\tau}_{C_*}$ and τ_{C_*} should be regarded as fixed once and for all.

Before defining the bias, we will first need the following definition.

Definition 3.8. For abelian groups A_1 and A_2 , we will let $\text{Hom}(A_1, A_2)$ (resp. $\text{Iso}(A_1, A_2)$) denote the set of homomorphisms (resp. isomorphisms) from A_1 to A_2 .

Let $\psi_1: A_1 \twoheadrightarrow B_1$ and $\psi_2: A_2 \twoheadrightarrow B_2$ be surjective homomorphisms for abelian groups B_1 and B_2 . We say a homomorphism $\varphi \in \text{Hom}(A_1, A_2)$ is a (ψ_1, ψ_2) -homomorphism if

$\varphi(\ker(\psi_1)) \subseteq \ker(\psi_2)$. Let $\text{Hom}_{\psi_1, \psi_2}(A_1, A_2) \subseteq \text{Hom}(A_1, A_2)$ denote the subset consisting of (ψ_1, ψ_2) -homomorphisms. There is an induced function

$$(\psi_1, \psi_2)_*: \text{Hom}_{\psi_1, \psi_2}(A_1, A_2) \rightarrow \text{Hom}(B_1, B_2), \quad \varphi \mapsto (x \mapsto \psi_2(\varphi(\tilde{x})))$$

where $\tilde{x} \in A_1$ is any lift of $x \in B_1$, i.e. $\psi_1(\tilde{x}) = x$.

Similarly, an isomorphism $\varphi \in \text{Iso}(A_1, A_2)$ is a (ψ_1, ψ_2) -isomorphism if $\varphi(\ker(\psi_1)) = \ker(\psi_2)$, and we let $\text{Iso}_{\psi_1, \psi_2}(A_1, A_2) \subseteq \text{Iso}(A_1, A_2)$ denote the subset consisting of (ψ_1, ψ_2) -isomorphisms. The function $(\psi_1, \psi_2)_*$ defined above restricts to a function $\text{Iso}_{\psi_1, \psi_2}(A_1, A_2) \rightarrow \text{Iso}(B_1, B_2)$.

In the case where $A_1 = A_2 =: A$, $B_1 = B_2 =: B$ and $\psi_1 = \psi_2 =: \psi$, a (ψ, ψ) -isomorphism will be referred to as a ψ -automorphism. The set of ψ -automorphisms forms a subgroup $\text{Aut}_\psi(A) \leq \text{Aut}(A)$ and the induced function ψ_* is a group homomorphism $\psi_*: \text{Aut}_\psi(A) \rightarrow \text{Aut}(B)$.

Definition 3.9. Fix a reference complex \overline{C}_* with $(-1)^n \chi(\overline{C}_*) = \chi$ and let $L = H_n(\overline{C}_*)$. Define the *polarised bias obstruction group* to be

$$PB(G, n, \chi) := \frac{\text{Aut}(\widehat{L})}{\text{Aut}_\psi(L^G)}.$$

This depends on χ but, up to isomorphism, does not depend on the choice of reference complex \overline{C}_* (see Definition 3.7). If $\chi = \chi_{\min}(G, n)$, then we define $PB(G, n) := PB(G, n, \chi)$.

Let C_*, D_* be algebraic n -complexes such that $(-1)^n \chi(C_*) = (-1)^n \chi(D_*) = \chi$. Define the *bias invariant* to be:

$$\beta(C_*, D_*) := [\tau_{D_*}^{-1} \circ \sigma(C_*, D_*) \circ \tau_{C_*}] \in PB(G, n, \chi).$$

where $[\cdot]: \text{Aut}(\widehat{L}) \twoheadrightarrow PB(G, n, \chi)$ is the quotient map and τ_{C_*}, τ_{D_*} are as above.

It follows from Proposition 3.6 that the vanishing of $\beta(C_*, D_*)$ in its respective obstruction group does not depend on the choice of reference complex.

We will now establish the following, where $m_{(G, n)}$ is as defined in Definition 3.4. This implies that the bias invariant contains no information in the non-minimal case.

Proposition 3.10.

- (i) *There is an isomorphism $PB(G, n) \cong (\mathbb{Z}/m)^\times / \{\pm 1\}$ where $m = m_{(G, n)}$.*
- (ii) *If $\chi > \chi_{\min}(G, n)$, then $PB(G, n, \chi) = 0$.*

This is a consequence of the following, which was pointed out by Webb [69, Corollary 3.2]. In what follows, we write $a \mid b$ for integers $a, b \geq 1$ if a divides b . We write $m_1 \mid \cdots \mid m_d$ for integers $m_1, \dots, m_d \geq 1$ if $m_i \mid m_{i+1}$ for all $1 \leq i \leq d-1$.

Lemma 3.11. *Let A be a finite abelian group, let $d \geq 1$ and let $\psi: \mathbb{Z}^d \rightarrow A$ be a surjective homomorphism. Suppose A has invariant factors $m_1 \mid \cdots \mid m_d$ (possibly with some $m_i = 1$), i.e. $A \cong \mathbb{Z}/m_1 \times \cdots \times \mathbb{Z}/m_d$. Consider the following composition:*

$$\rho: \text{Aut}(A) \rightarrow \text{Aut}(\mathbb{Z}/m_1 \otimes_{\mathbb{Z}} A) \xrightarrow{\det} (\mathbb{Z}/m_1)^\times \rightarrow (\mathbb{Z}/m_1)^\times / \{\pm 1\}.$$

Then ρ is surjective and $\text{im}(\psi_: \text{Aut}_\psi(\mathbb{Z}^d) \rightarrow \text{Aut}(A)) = \ker(\rho)$. In particular, we have that $\text{im}(\text{Aut}_\psi(\mathbb{Z}^d)) \trianglelefteq \text{Aut}(A)$ is a normal subgroup and ρ induces an isomorphism*

$$\rho_*: \frac{\text{Aut}(A)}{\text{Aut}_\psi(\mathbb{Z}^d)} \rightarrow (\mathbb{Z}/m_1)^\times / \{\pm 1\}.$$

Note that ρ only depends on A and d , and not on the choice of ψ .

Proof of Proposition 3.10. Let \overline{C}_* be a reference complex with $(-1)^n \chi(\overline{C}_*) = \chi$, let $L = H_n(\overline{C}_*)$, let $\psi: L^G \rightarrow \widehat{L}$ denote the quotient map, let $d = \text{rank}_{\mathbb{Z}}(L^G)$ and suppose \widehat{L} has invariant factors $m_1 \mid \cdots \mid m_d$ (possibly with some $m_i = 1$). Then Lemma 3.11 implies that $PB(G, n, \chi) \cong (\mathbb{Z}/m_1)^\times / \{\pm 1\}$.

(i) If $\chi = \chi_{\min}(G, n)$, then $r_{(G, n)} = \text{rank}_{\mathbb{Z}}(L^G)$ and so $m_1 = m_{(G, n)}$ (see Definition 3.4).

(ii) Suppose $\chi > \chi_{\min}(G, n)$, so that $r := \chi - \chi_{\min}(G, n) > 0$. Let \overline{D}_* be a reference complex with $(-1)^n \chi(\overline{D}_*) = \chi_{\min}(G, n)$, and let $L_0 = H_n(\overline{D}_*)$. Since L^G and \widehat{L} depend only on χ and not

on \overline{C}_* (see Definition 3.7), we have that $L^G \cong (L_0 \oplus \mathbb{Z}G^r)^G \cong L_0^G \oplus \mathbb{Z}^r$ and $\widehat{L} \cong (L_0 \oplus \mathbb{Z}G^r)^\wedge \cong \widehat{L}_0$. Since $\text{rank}_{\mathbb{Z}}(L_0^G) \geq d(\widehat{L}_0)$, it follows that $\text{rank}_{\mathbb{Z}}(L^G) \geq d(\widehat{L}) + r > d(\widehat{L})$. Hence $m_1 = 1$. \square

Remark 3.12. The quotient map $\psi: L^G \twoheadrightarrow \widehat{L}$ is induced by the identification $\widehat{L} \cong L^G/(N \cdot L)$. If $d = \text{rank}_{\mathbb{Z}}(L^G)$, then we have isomorphisms $L^G \cong \mathbb{Z}^d$ and $\widehat{L} \cong \mathbb{Z}/m_1 \times \cdots \times \mathbb{Z}/m_d$ for some m_i such that $m_i \mid m_{i+1}$ for all i . In fact, we can choose these isomorphisms such that the induced map $\psi: \mathbb{Z}^d \twoheadrightarrow \mathbb{Z}/m_1 \times \cdots \times \mathbb{Z}/m_d$ is a direct sum of quotient maps $\mathbb{Z} \twoheadrightarrow \mathbb{Z}/m_i$ for $1 \leq i \leq d$. To see this, note that since \widehat{L} is a finite group, the kernel $\ker \psi$ of the surjection $\psi: L^G \twoheadrightarrow \widehat{L}$ must be of the form $\ker \psi \cong \mathbb{Z}^d$.

Since \mathbb{Z} is a principal ideal domain, we can choose bases of $\ker \psi$ and L^G so that the inclusion map $\ker \psi \hookrightarrow L^G$ is given by a $d \times d$ matrix with non-zero diagonal entries n_1, \dots, n_d such that $n_i \mid n_{i+1}$ for all i . By comparing cokernels, it follows that $n_i = m_i$ for all i . By taking the induced generating set for \widehat{L} , we obtain identifications such that ψ is as required.

In light of Proposition 3.10 (ii), we will now restrict to the case of minimal complexes. The bias invariant has the following two basic properties. The first can be extracted from [43, Theorem 1.13], and the second is a consequence of the independence of the choice of chain map.

Proposition 3.13. *The bias invariant is a chain homotopy invariant. In particular, if D_* is an algebraic n -complex with $(-1)^n \chi(D_*) = \chi_{\min}(G, n)$, and $m = m_{(G, n)}$, then the bias invariant defines a map*

$$\beta: \text{Alg}_{\min}(G, n) \rightarrow (\mathbb{Z}/m)^\times / \{\pm 1\}, \quad C_* \mapsto \beta(C_*, D_*).$$

Proposition 3.14. *Let $C_*, D_*, E_* \in \text{Alg}_{\min}(G, n)$ and $m = m_{(G, n)}$, then*

$$\beta(C_*, E_*) = \beta(C_*, D_*) \cdot \beta(D_*, E_*) \in (\mathbb{Z}/m)^\times / \{\pm 1\}.$$

We conclude this section by noting that Metzler's original formulation differs to the one presented above. We will present it here since it will be useful for the explicit computations in Section 9b. A convenient reference for this formulation is [43].

Observe that, if $C_*, D_* \in \text{Alg}_{\min}(G, n)$, then $H_n(\mathbb{Z} \otimes_{\mathbb{Z}G} C_*)$ and $H_n(\mathbb{Z} \otimes_{\mathbb{Z}G} D_*)$ are free abelian groups of the same rank. The following is a consequence of [43, Lemma 1.8 & Exercise 1.10] and [55, p163], and gives another formulation of the bias invariant.

We use the identification $PB(G, n) \cong (\mathbb{Z}/m)^\times / \{\pm 1\}$ given in Proposition 3.10.

Proposition 3.15. *Let C_*, D_* be algebraic n -complexes such that $(-1)^n \chi(C_*) = (-1)^n \chi(D_*) = \chi_{\min}(G, n)$ and let $h: C_* \rightarrow D_*$ be a chain map such that $H_0(h) = \text{id}_{\mathbb{Z}}$. Fix identifications $H_n(\mathbb{Z} \otimes_{\mathbb{Z}G} C_*) \cong \mathbb{Z}^r$ and $H_n(\mathbb{Z} \otimes_{\mathbb{Z}G} D_*) \cong \mathbb{Z}^r$, and view $H_n(\text{id}_{\mathbb{Z}} \otimes h): H_n(\mathbb{Z} \otimes_{\mathbb{Z}G} C_*) \rightarrow H_n(\mathbb{Z} \otimes_{\mathbb{Z}G} D_*)$ as an element of $M_r(\mathbb{Z})$. Let $m = m_{(G, n)}$ and let $\cdot: \mathbb{Z} \rightarrow (\mathbb{Z}/m) / \{\pm 1\}$ denote the quotient map. Then:*

$$\beta(C_*, D_*) = \overline{\det(H_n(\text{id}_{\mathbb{Z}} \otimes h))} \in (\mathbb{Z}/m)^\times / \{\pm 1\}.$$

3c. The bias invariant for (G, n) -complexes. The following is implicit in [43, Definition 1.11]. The proof is a consequence of Proposition 3.14 and is omitted for brevity.

Proposition 3.16. *Let G be a finite group and let $n \geq 2$. If X is a finite (G, n) -complex with $(-1)^n \chi(X) = \chi$, then the map*

$$\varphi_{(G, n, \chi)}: \text{Aut}(G) \rightarrow PB(G, n, \chi), \quad \theta \mapsto \beta(C_*(\tilde{X}), C_*(\tilde{X})_\theta)$$

is a group homomorphism and is independent of the choice of X . If $\chi = \chi_{\min}(G, n)$, then we define $\varphi_{(G, n)} := \varphi_{(G, n, \chi)}$.

We will now use Proposition 2.2 to adapt the notion of bias to finite (G, n) -complexes.

Definition 3.17 (Bias invariant for (G, n) -complexes). Let $\varphi_{(G, n, \chi)}$ be the group homomorphism given in Proposition 3.16. Define

$$D(G, n, \chi) := \text{im}(\varphi_{(G, n, \chi)}: \text{Aut}(G) \rightarrow PB(G, n, \chi)) \leq PB(G, n, \chi)$$

and define the *bias obstruction group* for (G, n) -complexes to be

$$B(G, n, \chi) := \frac{PB(G, n, \chi)}{D(G, n, \chi)}.$$

If $\chi = \chi_{\min}(G, n)$, then we define $D(G, n) := D(G, n, \chi)$ and $B(G, n) := B(G, n, \chi)$. When $n = 2$, we write $B(G) := B(G, 2)$ and $D(G) := D(G, 2)$.

Let X, Y be finite (G, n) -complexes with $(-1)^n \chi(X) = (-1)^n \chi(Y) = \chi$. Define the *bias invariant* to be

$$\beta(X, Y) := [\beta(C_*(\tilde{X}), C_*(\tilde{Y}))] \in B(G, n, \chi)$$

where $[\cdot]: PB(G, n, \chi) \rightarrow B(G, n, \chi)$ is the quotient map.

By Proposition 3.10 (ii), we have that $B(G, n, \chi) = 0$ for $\chi > \chi_{\min}(G, n)$. In particular, if X and Y are finite (G, n) -complexes with $(-1)^n \chi(X) = (-1)^n \chi(Y) > \chi_{\min}(G, n)$, then $\beta(X, Y) = 0$. We will therefore primarily be interested in *minimal* finite (G, n) -complexes X , for which $(-1)^n \chi(X) = \chi_{\min}(G, n)$ has the minimum possible value (see Definition 2.4). We have:

Proposition 3.18. *If \bar{X} is a reference minimal (G, n) -complex, then the bias defines a map*

$$\beta: \text{HT}_{\min}(G, n) \rightarrow B(G, n) \cong \frac{(\mathbb{Z}/m)^\times}{\pm D(G, n)}, \quad X \mapsto \beta(X, \bar{X}).$$

Thus, β is an invariant of minimal (G, n) -complexes up to homotopy equivalence.

It follows from work of Metzler [47], Sieradski [58], Sieradski-Dyer [57], Browning [9, 10] and Linnell [46] that, for any finite abelian group G , the bias invariant completely classifies minimal (G, n) -complexes up to homotopy equivalence (Theorem 8.1). More details on the computation of the bias invariant, particularly for finite abelian groups, can be found in Section 8.

Remark 3.19. If (G, n) does not satisfy the strong minimality hypothesis (Definition 3.1), then Remark 3.5 implies that $m_{(G, n)} = 1$ and so $B(G, n) = 0$. In particular, the strong minimality hypothesis is a necessary condition for the non-vanishing of the bias invariant.

Remark 3.20. It is currently open whether or not there exists a finite group G and finite (G, n) -complexes X, Y with $(-1)^n \chi(X) = (-1)^n \chi(Y) > \chi_{\min}(G, n)$ but $X \not\cong Y$ (see [53, Question 7.4]). The above shows that the bias invariant cannot be used to distinguish such examples if they exist. It is known that such examples do not exist if $(-1)^n \chi(X) = (-1)^n \chi(Y) > \chi_{\min}(G, n) + 1$ [19, 71] or if n is even [10].

4. PRELIMINARIES ON HERMITIAN FORMS

4a. Hermitian forms on R -modules. A convenient reference is Scharlau [56] (see also [63, Section 6.2]). Let R be a ring with involution and let L be an R -module. We will mostly be concerned with commutative rings $R = \mathbb{Z}$ or \mathbb{Z}/m , with trivial involution, or $R = \mathbb{Z}G$, with involution $g \mapsto g^{-1}$, for $g \in G$ a finite group. A *sesquilinear form* on L is a bilinear form

$$h: L \times L \rightarrow R$$

such that $h(a \cdot m, b \cdot n) = a \cdot h(m, n) \cdot \bar{b} \in R$ for all $a, b \in R$ and $m, n \in L$. Define $\text{Sesq}(L)$ to be the set of sesquilinear forms on L . This is an abelian group under pointwise addition of functions. The term *quadratic form* and notation (L, q) for $q \in \text{Sesq}(L)$ is standard in the algebraic theory.

Let $T_\varepsilon: \text{Sesq}(L) \rightarrow \text{Sesq}(L)$, where $\varepsilon \in \{\pm 1\} \subseteq R$, denote the transpose operator, defined by $(Th)(m, n) = \varepsilon \overline{h(n, m)}$, for all $m, n \in L$. An ε -*hermitian form* on L is a sesquilinear form h such that $T_\varepsilon h = h$, or $h(m, n) = \varepsilon \overline{h(n, m)}$ for all $m, n \in L$. Define $\text{Herm}_\varepsilon(L) = \ker(1 - T_\varepsilon)$ to be the set of ε -hermitian forms on L , which is a subgroup of $\text{Sesq}(L)$. The standard terminology is *symmetric* for $\varepsilon = +1$ and *skew-symmetric* for $\varepsilon = -1$. When $\varepsilon = +1$, we will just write $\text{Herm}(L)$ to simplify the notation.

Let $\text{ad}: \text{Sesq}(L) \rightarrow \text{Hom}_R(L, L^*)$, $h \mapsto (m \mapsto h(-, m))$, denote the *adjoint* map, where we consider L^* as a left R -module via the involution on R . We say that an ε -hermitian form $h \in \text{Herm}_\varepsilon(L)$ is *non-singular* if $\text{ad}(h)$ is an R -isomorphism.

Definition 4.1. Let L be an R -module and let $h \in \text{Herm}_\varepsilon(L)$. Then the associated *metabolic form* is

$$\text{Met}_\varepsilon(L, h): (L^* \oplus L) \times (L^* \oplus L) \rightarrow R, \quad (\varphi_1, m_1), (\varphi_2, m_2) \mapsto \varepsilon \overline{\varphi_1(m_2)} + \varphi_2(m_1) + h(m_1, m_2).$$

We have $\text{Met}_\varepsilon(L, h) \in \text{Herm}_\varepsilon(L^* \oplus L)$. In matrix notation, we can write this as $\text{Met}_\varepsilon(L, h) = \begin{pmatrix} 0 & 1 \\ \varepsilon & h \end{pmatrix}$ where the off-diagonal maps are understood to denote the maps induced by evaluation $(\varphi, m) \mapsto \varphi(m)$. The *hyperbolic form* on L is defined as $H_\varepsilon(L) := \text{Met}_\varepsilon(L, 0)$.

A form $h \in \text{Herm}_\varepsilon(L)$ is called *even* if $h = q + T_\varepsilon q \in \text{im}(1 + T_\varepsilon)$ for some $q \in \text{Sesq}(L)$. In this case, we say the h admits a quadratic refinement q . A form $h \in \text{Herm}_\varepsilon(L)$ is called *weakly even* if $h(m, m) = a + \varepsilon \bar{a}$, for some $a \in R$ and all $m \in L$. These notions are equivalent if L is a projective R -module, but not in general.

In the other direction, a form $q \in \text{Sesq}(L)$ is called a *non-singular ε -quadratic form* if its associated hermitian form $h = q + T_\varepsilon q$ is non-singular.

4b. Hermitian forms on $\mathbb{Z}G$ -modules. Let G be a finite group and let L be a $\mathbb{Z}G$ -module. We will now define an alternative notion of dual module. For a $\mathbb{Z}G$ -module L , define $L^* := \text{Hom}_\mathbb{Z}(L, \mathbb{Z})$ to be the $\mathbb{Z}G$ -module where, for $r \in \mathbb{Z}G$ and $\varphi \in \text{Hom}_\mathbb{Z}(L, \mathbb{Z})$, we let $(r \cdot \varphi)(m) := \varphi(\bar{r} \cdot m)$ for all $m \in L$. This coincides with the notion of dual module defined previously via the $\mathbb{Z}G$ -isomorphism:

$$\psi: \text{Hom}_\mathbb{Z}(L, \mathbb{Z}) \rightarrow \text{Hom}_{\mathbb{Z}G}(L, \mathbb{Z}G), \quad \varphi \mapsto \hat{\varphi}, \quad \text{with} \quad \hat{\varphi}(m) = \sum_{g \in G} \varphi(g^{-1} \cdot m)g$$

(see, for example, [8, VI.3.4]). The inverse is given by $\hat{\varphi} \mapsto \varepsilon_1 \circ \hat{\varphi}$, where for $h \in G$, we let $\varepsilon_h: \mathbb{Z}G \rightarrow \mathbb{Z}$ denote the map $\sum_{g \in G} n_g g \mapsto n_h$.

Definition 4.2. Let $\text{Sym}_\varepsilon(L)$ denote the set of homomorphisms

$$\text{Sym}_\varepsilon(L) = \{\varphi \in \text{Hom}_\mathbb{Z}(L \otimes_\mathbb{Z} L, \mathbb{Z})^G : \varphi(m \otimes n) = \varepsilon \varphi(n \otimes m), \text{ for all } m, n \in L\}$$

where the action of G on $\text{Hom}_\mathbb{Z}(L \otimes_\mathbb{Z} L, \mathbb{Z})$ is defined by $(g \cdot \varphi)(m, n) = \varphi(gm, gn)$.

In other words, $\text{Sym}_\varepsilon(L)$ is the set of ε -symmetric bilinear forms $h: L \times L \rightarrow \mathbb{Z}$ such that $h(g \cdot m, g \cdot n) = h(m, n)$ for all $g \in G$ and $m, n \in L$.

There is a canonical isomorphism of abelian groups (with inverse induced by ψ):

$$\Xi: \text{Sesq}(L) \xrightarrow{\cong} \text{Hom}_\mathbb{Z}(L \otimes_\mathbb{Z} L, \mathbb{Z})^G, \quad h \mapsto (m \otimes n \mapsto \varepsilon_1(h(m, n))),$$

given by composing with $\varepsilon_1: \mathbb{Z}G \rightarrow \mathbb{Z}$. The following is a consequence of the definition of Ξ on restriction to $\text{Herm}_\varepsilon(L) \subseteq \text{Sesq}(L)$.

Proposition 4.3. *The map Ξ induces a natural correspondence $\text{Herm}_\varepsilon(L) \cong \text{Sym}_\varepsilon(L)$.*

Convention 4.4. From now on, unless stated otherwise, we will use the natural correspondence $\Xi: \text{Herm}_\varepsilon(L) \cong \text{Sym}_\varepsilon(L)$ to work with \mathbb{Z} -valued forms rather than $\mathbb{Z}G$ -valued forms. In particular, $\text{Met}_\varepsilon(L, h)$ and $H_\varepsilon(L)$ will mean the \mathbb{Z} -valued equivalent metabolic or hyperbolic forms.

Let R be a commutative ring and let RG denote the group ring with involution $g \mapsto g^{-1}$. Let $\mathfrak{S}_\varepsilon(G, R)$ denote the category whose objects are forms $h \in \text{Sym}_\varepsilon(L)$, for some RG -module L , with morphisms from (L, h) to (L', h') given by RG -module homomorphisms $f: L \rightarrow L'$ such that $h'(f(m), f(n)) = h(m, n)$, for all $m, n \in L$. If $G = 1$ is the trivial group, we will write $\mathfrak{S}_\varepsilon(R) := \mathfrak{S}_\varepsilon(1, R)$. We will later be interested in $\mathfrak{S}_\varepsilon(R)$ in the case where $R = \mathbb{Z}$ or \mathbb{Z}/m for $m \in \mathbb{Z}$ (see Definition 4.7).

We will now restrict to the case where $R = \mathbb{Z}$. There is an action of $\text{Aut}(G)$ on $\mathfrak{S}_\varepsilon(G, \mathbb{Z})$ defined as follows. Let $\theta \in \text{Aut}(G)$ and $h \in \text{Sym}_\varepsilon(L)$. Let L_θ be the $\mathbb{Z}G$ -module whose underlying abelian group is L and with G -action defined by $g \cdot m := \theta(g) \cdot m$, for all $g \in G$ and $m \in L$. Then $h_\theta \in \text{Sym}_\varepsilon(L_\theta)$ is the induced form on L_θ .

Example 4.5. Let G be a finite group. If M is a closed oriented 4-dimensional Poincaré complex with $\pi_1(M) \cong G$, then the equivariant intersection form S_M is a hermitian form on $\pi_2(M)$. That is, $S_M \in \text{Herm}(\pi_2(M))$. Using the convention above, we will view this as a bilinear form

$$S_M: \pi_2(M) \times \pi_2(M) \rightarrow \mathbb{Z}$$

which is G -invariant and symmetric, so that $S_M \in \text{Sym}(\pi_2(M))$. The hermitian form S_M satisfies the Bredon condition [7, Theorem 7.4] that $S_M(\tau \cdot m, m) \equiv 0 \pmod{2}$ for all $m \in \pi_2(M)$ and $\tau \in G$ of order two. This leads to the observation that the $\mathbb{Z}G$ -valued hermitian form is weakly even if the universal covering of M is a spin manifold. Note that an even metabolic form is isometric to a hyperbolic form, but this is not true for weakly even metabolic forms.

The following result is well-known. Recall that a $\mathbb{Z}G$ -lattice is a $\mathbb{Z}G$ -module which is finitely generated and torsion free as an abelian group.

Proposition 4.6. *Let L be a $\mathbb{Z}G$ -lattice, where G is a finite group of odd order. If (L, h) is a weakly even form, then (L, h) is even and the metabolic form $\text{Met}_\varepsilon(L, h)$ is hyperbolic.*

Proof. A metabolic form $\text{Met}_\varepsilon(L, h)$ is weakly even if the form (L, h) is a weakly even ε -symmetric form on the $\Lambda = \mathbb{Z}G$ module L , where L is a $\mathbb{Z}G$ -lattice. Since h is ε -symmetric, it defines a class

$$[h] \in \hat{H}^0(\mathbb{Z}/2; \text{Sesq}(L))$$

which vanishes if and only if $h = q + T_\varepsilon q$ for some $q \in \text{Sesq}(L)$. Since the obstruction group is an \mathbb{F}_2 vector space, and the reduction mod 2 map

$$\hat{H}^0(\mathbb{Z}/2; \text{Sesq}(L)) \rightarrow \hat{H}^0(\mathbb{Z}/2; \text{Sesq}(L/2))$$

is an injection (since $\text{Sesq}(L)$ is torsion free as an abelian group), it follows that (L, h) is an even form if and only if its reduction $(L/2, \bar{h})$ is an even form with values in $\Lambda/2 = \mathbb{F}_2[G]$.

If G has odd order, the ring $\mathbb{F}_2[G]$ is semisimple [13, Theorem 3.14], so the $\Lambda/2$ -module $L/2$ is projective. Hence $(L/2, \bar{h})$ is an even form by [5, Proposition 3.4], and so is h . \square

Definition 4.7. The fixed-point functor $\mathfrak{S}_\varepsilon(G, \mathbb{Z}) \rightarrow \mathfrak{S}_\varepsilon(\mathbb{Z})$ is defined on objects by $h \mapsto h^G$, where $h^G \in \text{Sym}_\varepsilon(L^G)$ is the restriction of the form h to the fixed set $L^G \subset L$. The Tate functor $\mathfrak{S}_\varepsilon(G, \mathbb{Z}) \rightarrow \mathfrak{S}_\varepsilon(\mathbb{Z}/|G|)$ is defined on objects by $h \mapsto \hat{h}$ where $\hat{h}(\bar{m}, \bar{n}) = [h^G(m, n)] \in \mathbb{Z}/|G|$ for all $m, n \in L^G$, and where $\bar{\cdot}$ denotes the quotient map $L^G \rightarrow L^G/(N \cdot L) = \hat{H}^0(G; L)$ and $[\cdot]$ denotes the quotient map $\mathbb{Z} \rightarrow \mathbb{Z}/|G|$. The functors can be defined on the morphisms in the natural way.

Note that $\hat{\mathbb{Z}} = \hat{H}^0(G; \mathbb{Z}) \cong \mathbb{Z}/|G|$ is a ring and, if L is a $\mathbb{Z}G$ -module, then \hat{L} is a $\hat{\mathbb{Z}}$ module in a natural way. For example, we can take the action induced by the cup product

$$\hat{H}^0(G; \mathbb{Z}) \times \hat{H}^0(G; L) \xrightarrow{-\cup-} \hat{H}^0(G; \mathbb{Z} \otimes_{\mathbb{Z}} L) \xrightarrow{\cong} \hat{H}^0(G; L).$$

The following is [8, Exercise VI.7.3].

Proposition 4.8. *Let L be a $\mathbb{Z}G$ -lattice. Then the cup product and the evaluation map $L^* \otimes_{\mathbb{Z}} L \rightarrow \mathbb{Z}$, $\varphi \otimes m \mapsto \varphi(m)$ induce a non-singular duality pairing*

$$\hat{H}^0(G; L^*) \times \hat{H}^0(G; L) \xrightarrow{-\cup-} \hat{H}^0(G; L^* \otimes_{\mathbb{Z}} L) \rightarrow \hat{H}^0(G; \mathbb{Z}).$$

In particular, there is an isomorphism of abelian groups $\hat{H}^0(G; L^) \cong \text{Hom}_{\mathbb{Z}}(\hat{H}^0(G; L), \mathbb{Z}/|G|)$.*

If A is a $\mathbb{Z}/|G|$ -module, then we will write $A^* := \text{Hom}_{\mathbb{Z}}(A, \mathbb{Z}/|G|)$. The above shows that, if L is a $\mathbb{Z}G$ -lattice, then there is a canonical identification $\theta: (L^*)^\wedge \cong (\hat{L})^*$. If $f: L_1 \rightarrow L_2$ is an isomorphism of $\mathbb{Z}G$ -lattices, then the duality pairing implies that $(f^*)^{-1} \oplus f: L_1^* \oplus L_1 \rightarrow L_2^* \oplus L_2$ induces an isomorphism commuting with the identifications $(L_i^*)^\wedge \cong (\hat{L}_i)^*$, for $i = 1, 2$.

Proposition 4.9. *Let $h \in \text{Sym}_\varepsilon(L)$ be a \mathbb{Z} -valued hermitian form on a $\mathbb{Z}G$ -module L . Then the form $\hat{h} \in \text{Sym}_\varepsilon(\hat{H}^0(G; L))$ defined in Definition 4.7 coincides with the form induced by the cup product on Tate cohomology:*

$$\hat{H}^0(G; L) \times \hat{H}^0(G; L) \xrightarrow{\cup} \hat{H}^0(G; L \otimes_{\mathbb{Z}} L) \xrightarrow{h} \hat{H}^0(G; \mathbb{Z}) \cong \mathbb{Z}/|G|.$$

If L is a $\mathbb{Z}G$ -lattice and h is non-singular, then h^G is non-degenerate and \hat{h} is non-singular.

Proof. The cup product is induced by the composite $L^G \otimes L^G \subseteq (L \otimes L)^G \xrightarrow{h} \mathbb{Z}$, and the form \hat{h} is induced by the restriction $h^G: L^G \otimes L^G \rightarrow \mathbb{Z}$ of h to the fixed subspaces. Hence the form \hat{h} coincides with the form induced by the cup product.

If L is a $\mathbb{Z}G$ -lattice and h is non-singular, then for any non-zero $x \in L^G$, there exists $y \in L$ such that $h(x, y) \neq 0$, implying that $h(x, Ny) = h(Nx, y) = |G| \cdot h(x, y) \neq 0$. Hence h^G is non-degenerate. To show that $\widehat{h}: \widehat{L} \times \widehat{L} \rightarrow \mathbb{Z}/|G|$ is non-singular, we consider its adjoint $\text{ad } \widehat{h}: \widehat{L} \rightarrow (\widehat{L})^*$. By Proposition 4.8, there is an isomorphism $\theta: (L^*)^\wedge \xrightarrow{\cong} (\widehat{L})^*$, such that $\text{ad } \widehat{h} = \theta \circ (\text{ad } h)^\wedge$. To check this formula, let $i: L^G \rightarrow L$ denote the inclusion, and note that $\theta(\phi)(\bar{y}) = (i^*\phi)(y) \in \mathbb{Z}/|G|$, for any $\phi \in (L^*)^G$ and $y \in L^G$, where $i^*: (L^*)^G \rightarrow (L^G)^*$ is the induced restriction map. Then for $x \in L^G$, we set $\phi := \text{ad } h(x) \in (L^*)^G$ and see that $(i^* \text{ad } h)(x)(y) = h(x, y)$, implying the formula above.

Since h is non-singular, $\text{ad } h$ is an isomorphism, and hence $(\text{ad } h)^\wedge$ is also an isomorphism. Therefore $\text{ad } \widehat{h} = \theta \circ (\text{ad } h)^\wedge$ is an isomorphism and \widehat{h} is non-singular. \square

4c. Evaluation forms. Let G be a finite group and let L be a $\mathbb{Z}G$ -module. The *evaluation form* on L is the bilinear form

$$e_L: (L^* \oplus L) \times (L^* \oplus L) \rightarrow \mathbb{Z}, \quad (\varphi_1, m_1), (\varphi_2, m_2) \mapsto \varepsilon \varphi_1(m_2) + \varphi_2(m_1),$$

where $L^* = \text{Hom}_{\mathbb{Z}}(L, \mathbb{Z})$ and $e_L \in \text{Sym}_{\varepsilon}(L^* \oplus L)$, for $\varepsilon \in \{\pm 1\}$ is an ε -symmetric form. When the module L is understood from the context, we will often write $e = e_L$. This coincides with the hyperbolic form previously defined, so that e_L is isometric to $H_{\varepsilon}(L)$ and $\text{Met}_{\varepsilon}(L, \phi) \cong \begin{pmatrix} 0 & 1 \\ \varepsilon & 0 \end{pmatrix}$, where $\phi \in \text{Sym}_{\varepsilon}(L)$. We will now define two related evaluation forms.

Definition 4.10. The restriction of the evaluation form e_L to the fixed set $(L^* \oplus L)^G = H^0(G; L^* \oplus L)$ induces a bilinear form:

$$e_L^G: ((L^*)^G \oplus L^G) \times ((L^*)^G \oplus L^G) \rightarrow \mathbb{Z}.$$

This is an ε -symmetric form over the ring \mathbb{Z} with trivial involution, and coincides with h^G for $h = e_L$ from Definition 4.7. We write $e_L^G \in \text{Sym}_{\varepsilon}((L^*)^G \oplus L^G)$.

Definition 4.11. By functoriality of $\widehat{H}^0(G; -)$, and the isomorphism $(L^*)^\wedge \cong (\widehat{L})^*$, there is an induced bilinear form:

$$\widehat{e}_L: ((L^*)^\wedge \oplus \widehat{L}) \times ((L^*)^\wedge \oplus \widehat{L}) \rightarrow \mathbb{Z}/|G|.$$

This is a non-singular ε -symmetric form of $\mathbb{Z}/|G|$ -modules. We write $\widehat{e}_L \in \text{Sym}_{\varepsilon}(\widehat{L}^* \oplus \widehat{L})$.

The form \widehat{e}_L in the definition above coincides with \widehat{h} for $h = e_L$ from Definition 4.7. This is a consequence of Proposition 4.9.

The following will be useful later (see Definition 4.7).

Proposition 4.12. Let $\phi \in \text{Sym}_{\varepsilon}(L)$ be an ε -symmetric form such that $\phi^G = 0$. If $h = \text{Met}_{\varepsilon}(L, \phi)$ is the corresponding metabolic form, then $h^G \cong e_L^G$ and $\widehat{h} \cong \widehat{e}_L$.

Proof. The first statement is immediate from the assumption that $\phi^G = 0$. It follows from Definition 4.7 that \widehat{h} is the reduction of h^G , and so we also have $\widehat{h} \cong \widehat{e}_L$. \square

The following summary is the basis for the explicit models for e_L^G and \widehat{e}_L we give in Proposition 4.15.

Lemma 4.13. Under the inclusion map $(L^*)^G \subseteq (L^G)^*$ induced by restriction, e_L^G is the restriction of the evaluation form

$$e_{L^G}: ((L^G)^* \oplus L^G) \times ((L^G)^* \oplus L^G) \rightarrow \mathbb{Z}, \quad (\varphi_1, m_1), (\varphi_2, m_2) \mapsto \varphi_1(m_2) + \varphi_2(m_1).$$

Under the canonical identification $(L^*)^\wedge \oplus \widehat{L} \cong (\widehat{L})^* \oplus \widehat{L}$, the form \widehat{e}_L corresponds to the evaluation form

$$e_{\widehat{L}}: ((\widehat{L})^* \oplus \widehat{L}) \times ((\widehat{L})^* \oplus \widehat{L}) \rightarrow \mathbb{Z}, \quad (\varphi_1, m_1), (\varphi_2, m_2) \mapsto \varphi_1(m_2) + \varphi_2(m_1).$$

Proof. This follows directly from the definitions above and Proposition 4.9. \square

4d. Matrix representations for the evaluation forms. Let G be a finite group, let L be a $\mathbb{Z}G$ -lattice and let $e = e_L$ be the non-singular evaluation form. The following is a consequence of [42, Propositions III.1 & III.2].

Lemma 4.14. Suppose $L^G \cong \mathbb{Z}^d$ for some $d \geq 0$ and that \widehat{L} has invariant factors $n_d \mid \cdots \mid n_1$ (possibly with some $n_i = 1$), i.e. $\widehat{L} \cong \mathbb{Z}/n_1 \times \cdots \times \mathbb{Z}/n_d$. Then there is an exact sequence of abelian groups:

$$0 \rightarrow (L^*)^G \xrightarrow{\text{rest}} (L^G)^* \rightarrow \mathbb{Z}/\beta_1 \times \cdots \times \mathbb{Z}/\beta_d \rightarrow 0$$

where $\text{rest}: \varphi \mapsto \varphi|_{L^G}$ is the restriction map and $\beta_i := |G|/n_i$ for $1 \leq i \leq d$ so that $\beta_1 \mid \cdots \mid \beta_d$.

By combining this with Lemma 4.13, we now have the following (which can be found in the discussion on [42, p24]).

Proposition 4.15. Let n_1, \dots, n_d and β_1, \dots, β_d be as in Lemma 4.14. Then, by choosing bases for $(L^*)^G$ and $(L^G)^*$ such that $(L^*)^G = \beta_1 \mathbb{Z} \oplus \cdots \beta_d \mathbb{Z} \subseteq \mathbb{Z}^d = (L^G)^*$, the bilinear form e^G has the matrix representation:

$$e^G: \underbrace{(\mathbb{Z}^d \oplus \mathbb{Z}^d)}_{\cong (L^*)^G \oplus L^G} \times \underbrace{(\mathbb{Z}^d \oplus \mathbb{Z}^d)}_{\cong (L^*)^G \oplus L^G} \rightarrow \mathbb{Z}, \quad e^G = \left(\begin{array}{c|c} 0 & \begin{smallmatrix} \beta_1 & & \\ & \ddots & \\ & & \beta_d \end{smallmatrix} \\ \hline \begin{smallmatrix} \beta_1 & & \\ & \ddots & \\ & & \beta_d \end{smallmatrix} & 0 \end{array} \right)$$

We can choose bases for $(L^*)^\wedge$ and \widehat{L} such that \widehat{e} has matrix representation:

$$\widehat{e}: \underbrace{(\bigoplus_{i=1}^d \mathbb{Z}/n_i \oplus \bigoplus_{i=1}^d \mathbb{Z}/n_i)}_{\cong (L^*)^\wedge \oplus \widehat{L}} \times \underbrace{(\bigoplus_{i=1}^d \mathbb{Z}/n_i \oplus \bigoplus_{i=1}^d \mathbb{Z}/n_i)}_{\cong (L^*)^\wedge \oplus \widehat{L}} \rightarrow \mathbb{Z}/|G|,$$

$$\widehat{e} = \left(\begin{array}{c|c} 0 & \begin{smallmatrix} \beta_1 & & \\ & \ddots & \\ & & \beta_d \end{smallmatrix} \\ \hline \begin{smallmatrix} \beta_1 & & \\ & \ddots & \\ & & \beta_d \end{smallmatrix} & 0 \end{array} \right).$$

Remark 4.16. (a) That \widehat{e} is non-singular follows from [42, Theorem III.1] and the fact that e is non-singular. However, it also follows directly from Lemma 4.13.

(b) In the matrix representation for \widehat{e} , the non-zero entries β_i correspond to maps $\mathbb{Z}/n_i \times \mathbb{Z}/n_i \rightarrow \mathbb{Z}/|G|$, $(x, y) \mapsto \beta_i xy$. This is well-defined since $\beta_i = |G|/n_i$ and so, if $x \equiv x' \pmod{n_i}$ and $y \equiv y' \pmod{n_i}$, then $\beta_i xy \equiv \beta_i x'y' \pmod{|G|}$.

4e. Isometries of evaluation forms. The following definitions are motivated by the definitions made in Section 3b. The first is analogous to Definition 3.8. Even more generally, we could make both definitions for homomorphisms instead of automorphisms. Many of the results we need extend to this setting (see, for example, [55, Lemma 3 (§2)]).

Definition 4.17. For $i = 1, 2$, let L_i be a $\mathbb{Z}G$ -lattice, let $e_i = e_{L_i}$ be its evaluation form and let $\psi_i: L_i^G \rightarrow \widehat{L}_i$ and $\psi'_i: (L_i^*)^G \rightarrow (L_i^*)^\wedge$ denote the canonical reduction maps. Let $\Psi_i = \psi'_i \oplus \psi_i$.

We say an isometry $\varphi \in \text{Isom}(e_1^G, e_2^G)$ is a (Ψ_1, Ψ_2) -isometry if $\varphi(\ker(\Psi_1)) = \ker(\Psi_2)$. Let $\text{Isom}_{\Psi_1, \Psi_2}(e_1^G, e_2^G) \subseteq \text{Isom}(e_1^G, e_2^G)$ denote the subset consisting of (Ψ_1, Ψ_2) -isometries. There is an induced function

$$(\Psi_1, \Psi_2)_*: \text{Isom}_{\Psi_1, \Psi_2}(e_1^G, e_2^G) \rightarrow \text{Isom}(\widehat{e}_1, \widehat{e}_2), \quad \varphi \mapsto (x \mapsto \Psi_2(\varphi(\widetilde{x})))$$

where $\widetilde{x} \in (L_1^*)^G \oplus L_1^G$ is any lift of $x \in (L_1^*)^\wedge \oplus \widehat{L}_1$, i.e. $\Psi_1(\widetilde{x}) = x$.

In the case where $L_1 = L_2 =: L$, write $e = e_i$, $\psi = \psi_i$ and $\Psi = \Psi_i$ for $i = 1, 2$. A (Ψ, Ψ) -isometry will be referred to as a Ψ -isometry. The set of Ψ -isometries forms a subgroup $\text{Isom}_\Psi(e^G) \leq \text{Isom}(e^G)$ and the induced function $(\Psi, \Psi)_*$ is a group homomorphism $\Psi_*: \text{Isom}_\Psi(e^G) \rightarrow \text{Isom}(\widehat{e})$.

Before making the next definition, we will start by establishing the following.

Lemma 4.18. *For $i = 1, 2$, let L_i be a $\mathbb{Z}G$ -lattice and let $\psi_i: L_i^G \rightarrow \widehat{L}_i$ and $\psi'_i: (L_i^*)^G \rightarrow (L_i^*)^\wedge$ denote the canonical reduction maps. Let $f \in \text{Hom}_{\psi_1, \psi_2}(L_1^G, L_2^G)$. Then there exists a unique map $g \in \text{Hom}_{\psi'_1, \psi'_2}((L_2^*)^G, (L_1^*)^G)$ such that we have a commutative diagram*

$$\begin{array}{ccc} (L_2^*)^G & \xrightarrow{\text{rest}} & (L_2^G)^* \\ g \downarrow & & f^* \downarrow \\ (L_1^*)^G & \xrightarrow{\text{rest}} & (L_1^G)^* \end{array}$$

where rest denotes the restriction maps as used in Lemma 4.14. If $f \in \text{Iso}_{\psi_1, \psi_2}(L_1^G, L_2^G)$, then $g \in \text{Iso}_{\psi'_1, \psi'_2}((L_2^*)^G, (L_1^*)^G)$.

Proof. This is a consequence of the proof of [42, Proposition III.6]. Simply note that dualising the commutative diagram given there gives the commutative diagram we require. This works since all modules involved are $\mathbb{Z}G$ -lattices and so double dualising returns the original module. \square

From now on, let e_i, Ψ_i be as defined above for $i = 1, 2$. The following is defined in [42, p30-31].

Definition 4.19. An isometry $\rho \in \text{Isom}(\widehat{e}_1, \widehat{e}_2)$ is called *diagonal* if there exists an isomorphism of abelian groups $f: \widehat{L}_1 \rightarrow \widehat{L}_2$ such that $\rho = (f^*)^{-1} \oplus f$. Here $f^*: (\widehat{L}_2)^* \rightarrow (\widehat{L}_1)^*$ is viewed as a map $(L_2^*)^\wedge \rightarrow (L_1^*)^\wedge$ via the canonical identifications $(L_i^*)^\wedge \cong (\widehat{L}_i)^*$. We write $\text{Diag}(\widehat{e}_1, \widehat{e}_2)$ for the set of diagonal isometries from \widehat{e}_1 to \widehat{e}_2 , which we will view as a subset of $\text{Isom}(\widehat{e}_1, \widehat{e}_2)$. When $e_1 = e_2 =: e$, this defines a subgroup $\text{Diag}(\widehat{e}) \leq \text{Isom}(\widehat{e})$.

An isometry $\rho \in \text{Isom}(e_1^G, e_2^G)$ is called *diagonal* if there exists an isomorphism of abelian groups $f: L_1^G \rightarrow L_2^G$ such that $\rho = (f^*|_{(L_2^*)^G})^{-1} \oplus f$. Here we note that f^* restricts to an isomorphism $f^*|_{(L_2^*)^G}: (L_2^*)^G \rightarrow (L_1^*)^G$ by Lemma 4.18. We write $\text{Diag}(e_1^G, e_2^G)$ for the set of diagonal isometries from e_1^G to e_2^G , which we will view as a subset of $\text{Isom}(e_1^G, e_2^G)$. When $e_1 = e_2 =: e$, this defines a subgroup $\text{Diag}(e^G) \leq \text{Isom}(e^G)$.

In the notation of Definition 4.17, we write $\text{Diag}_{\Psi_1, \Psi_2}(e_1^G, e_2^G)$ to denote

$$\text{Diag}(e_1^G, e_2^G) \cap \text{Isom}_{\Psi_1, \Psi_2}(e_1^G, e_2^G).$$

The map $(\Psi_1, \Psi_2)_*$ defined in Definition 4.17 restricts to a function

$$(\Psi_1, \Psi_2)_*: \text{Diag}_{\Psi_1, \Psi_2}(e_1^G, e_2^G) \rightarrow \text{Diag}(\widehat{e}_1, \widehat{e}_2).$$

Remark 4.20. As a check: if $\rho = (f^*|_{(L_2^*)^G})^{-1} \oplus f \in \text{Diag}(e_1^G, e_2^G)$, then $\rho \in \text{Isom}_{\Psi_1, \Psi_2}(e_1^G, e_2^G)$ if and only if $f \in \text{Iso}_{\psi_1, \psi_2}(L_1^G, L_2^G)$, by Lemma 4.18. In addition, the commutative diagram in Lemma 4.18 shows that the Tate reductions of $f^*|_{(L_2^*)^G}$ and f^* agree, via the canonical identifications $(L_i^*)^\wedge \cong (\widehat{L}_i)^*$. Hence under the restriction map $(\Psi_1, \Psi_2)_*(\rho) \in \text{Diag}(\widehat{e}_1, \widehat{e}_2)$.

In the case where $e_1 = e_2 =: e$, we write this as $\text{Diag}_\Psi(e^G)$. The map Ψ_* restricts to a group homomorphism

$$\Psi_*: \text{Diag}_\Psi(e^G) \rightarrow \text{Diag}(\widehat{e}).$$

Note that we actually gave a definition of $\text{Diag}(\widehat{e}_1, \widehat{e}_2)$ which is a priori stronger than the one given in [42, p30-31]. We will now show that these definitions are equivalent subject to the hypothesis that the $\mathbb{Z}G$ -lattices L_1 and L_2 are $|G|$ -locally isomorphic, meaning that $\mathbb{Z}_{(p)} \otimes_{\mathbb{Z}} L_1 \cong \mathbb{Z}_{(p)} \otimes_{\mathbb{Z}} L_2$ as $\mathbb{Z}_{(p)}G$ -modules for all primes $p \mid |G|$, where $\mathbb{Z}_{(p)}$ is the localisation of \mathbb{Z} at the prime ideal (p) .

Proposition 4.21. *Let $\rho \in \text{Isom}(\widehat{e}_1, \widehat{e}_2)$. If L_1 and L_2 are $|G|$ -locally isomorphic, then $\rho \in \text{Diag}(\widehat{e}_1, \widehat{e}_2)$ if and only if there exist maps $f \in \text{Hom}_{\psi_1, \psi_2}(L_1^G, L_2^G)$ and $g \in \text{Hom}_{\psi_2, \psi_1}(L_2^G, L_1^G)$ such that $\rho = (g^*|_{(L_1^*)^G}) \oplus \bar{f}$, under the Tate reduction maps $\bar{\cdot} = (\psi'_1, \psi'_2)_*(\cdot)$ and $\bar{\cdot} = (\psi_1, \psi_2)_*(\cdot)$ in the two cases respectively.*

Proof. Firstly note that, since L_1 and L_2 are $|G|$ -locally isomorphic, it follows directly that $\text{rank}_{\mathbb{Z}} L_1^G = \text{rank}_{\mathbb{Z}} L_2^G$ and $\widehat{L}_1 \cong \widehat{L}_2$ (see, for example, the proof of [42, Proposition III.5]).

If $\rho \in \text{Diag}(\widehat{e}_1, \widehat{e}_2)$, then $\rho = (h^*)^{-1} \oplus h$ for some $h \in \text{Iso}(\widehat{L}_1, \widehat{L}_2)$. By Lemma 4.14, for $i = 1, 2$ we have two presentations

$$0 \rightarrow (L_i^*)^G \xrightarrow{\text{rest}} (L_i^G)^* \rightarrow Z/\beta_1 \times \cdots \times Z/\beta_d \rightarrow 0$$

of the form $0 \rightarrow \mathbb{Z}^d \rightarrow \mathbb{Z}^d \rightarrow A \rightarrow 0$, where $\text{rank}_{\mathbb{Z}} L_i^G = d$ and $A = \widehat{L}_1 \cong \widehat{L}_2$ is a finite abelian group. By the structure theorem for finitely generated abelian groups, any isomorphism $h: \widehat{L}_1 \rightarrow \widehat{L}_2$ can be lifted to a map $f \in \text{Hom}_{\psi_1, \psi_2}(L_1^G, L_2^G)$ (see Remark 3.12). Similarly, we have a lift $s: (L_1^*)^G \rightarrow (L_2^*)^G$ such that $\bar{s} = (f^*)^{-1}$. By Lemma 4.18, there exists a homomorphism $g: L_2^G \rightarrow L_1^G$ for which $g^*|_{(L_1^*)^G} = s$ (compare the last paragraph of the proof of [42, Proposition III.5]). Hence $\rho = (g^*|_{(L_1^*)^G}) \oplus \bar{f}$.

Conversely, suppose $f \in \text{Hom}_{\psi_1, \psi_2}(L_1^G, L_2^G)$ and $g \in \text{Hom}_{\psi_2, \psi_1}(L_2^G, L_1^G)$ are such that $\rho = (g^*|_{(L_1^*)^G}) \oplus \bar{f} \in \text{Isom}(\widehat{e}_1, \widehat{e}_2)$. Then $h = \bar{f} \in \text{Iso}(\widehat{L}_1, \widehat{L}_2)$ and $t = (g^*|_{(L_1^*)^G}) \in \text{Iso}((\widehat{L}_1)^*, (\widehat{L}_2)^*)$. It follows from [42, Proposition III.6] that there exists a unique $h' \in \text{Iso}((\widehat{L}_1)^*, (\widehat{L}_2)^*)$ for which $h' \oplus h \in \text{Isom}(\widehat{e}_1, \widehat{e}_2)$. Since $(h^*)^{-1} \in \text{Iso}((\widehat{L}_1)^*, (\widehat{L}_2)^*)$ and we have $(h^*)^{-1} \oplus h \in \text{Isom}(\widehat{e}_1, \widehat{e}_2)$, it follows that $t = (h^*)^{-1}$, and so $\rho \in \text{Diag}(\widehat{e}_1, \widehat{e}_2)$. \square

The following is immediate from Definition 4.19.

Proposition 4.22. *There are bijections*

$$\begin{aligned} A: \text{Diag}(\widehat{e}_1, \widehat{e}_2) &\rightarrow \text{Iso}(\widehat{L}_1, \widehat{L}_2), & (f^*)^{-1} \oplus f &\mapsto f \\ B: \text{Diag}(e_1^G, e_2^G) &\rightarrow \text{Iso}(L_1^G, L_2^G), & (f^*|_{(L_1^*)^G})^{-1} \oplus f &\mapsto f \end{aligned}$$

such that

- (i) B restricts to a bijection $B': \text{Diag}_{\Psi_1, \Psi_2}(e_1^G, e_2^G) \rightarrow \text{Iso}_{\psi_1, \psi_2}(L_1^G, L_2^G)$.
- (ii) There is a commutative diagram

$$\begin{array}{ccc} \text{Diag}_{\Psi_1, \Psi_2}(e_1^G, e_2^G) & \xrightarrow{(\Psi_1, \Psi_2)_*} & \text{Diag}(\widehat{e}_1, \widehat{e}_2) \\ B' \downarrow & & \downarrow A \\ \text{Iso}_{\psi_1, \psi_2}(L_1^G, L_2^G) & \xrightarrow{(\psi_1, \psi_2)_*} & \text{Iso}(\widehat{L}_1, \widehat{L}_2). \end{array}$$

- (iii) In the case where $e_1 = e_2$, A , B and B' are isomorphisms of abelian groups.

In particular, this shows that $\text{im}(\text{Diag}_{\Psi}(e^G)) \trianglelefteq \text{Diag}(\widehat{e})$ is a normal subgroup and there are isomorphisms of abelian groups

$$(4.23) \quad \frac{\text{Diag}(\widehat{e})}{\text{Diag}_{\Psi}(e^G)} \xrightarrow{A} \frac{\text{Aut}(\widehat{L})}{\text{Aut}_{\psi}(L^G)} \xrightarrow{\rho_*} (\mathbb{Z}/m)^{\times} / \{\pm 1\}$$

where ρ_* is the map defined in Lemma 3.11 and $m = m_{(G, n)}$. We will also need to consider non-diagonal isometries. The following special cases will suffice.

Definition 4.24. An isometry $\rho \in \text{Isom}(\widehat{e})$ is called (*upper*) *triangular* if there exists a homomorphism $f: \widehat{L} \rightarrow (\widehat{L})^*$ such that $\rho = \begin{pmatrix} \text{id} & f \\ 0 & \text{id} \end{pmatrix}: (\widehat{L})^* \oplus \widehat{L} \rightarrow (\widehat{L})^* \oplus \widehat{L}$. The set of triangular isometries defines a subgroup $\text{Tri}(\widehat{e}) \leq \text{Isom}(\widehat{e})$.

Note that $\text{Tri}(\widehat{e})$ is abelian since $\begin{pmatrix} \text{id} & f \\ 0 & \text{id} \end{pmatrix} \begin{pmatrix} \text{id} & g \\ 0 & \text{id} \end{pmatrix} = \begin{pmatrix} \text{id} & f+g \\ 0 & \text{id} \end{pmatrix}$ for homomorphisms $f, g: \widehat{L} \rightarrow (\widehat{L})^*$.

5. THE DOUBLING CONSTRUCTION

We will begin by defining the doubling construction for finite (G, n) -complexes. Details can be found in [42, Section II].

Let $n \geq 2$ and let G be a group of type F_n . If X is a finite (G, n) -complex, then there exists an embedding $i: X \hookrightarrow \mathbb{R}^{2n+1}$. Let $N(X) \subseteq \mathbb{R}^{2n+1}$ be a smooth regular neighbourhood of this embedding (unique up to concordance, by [66, p76]) and define $M(X) := \partial N(X)$. This is a closed oriented smooth stably parallelisable $2n$ -manifold (see, for example, [42, p15]). If X is well defined up to homotopy equivalence, then $M(X)$ is well-defined up to h -cobordism and, in

particular, does not depend on the choice of embedding or smooth regular neighbourhood. A polarisation $\pi_1(X) \cong G$ induces a polarisation $\pi_1(M(X)) \cong G$ and, from now on, we will assume that the manifolds $M(X)$ come equipped with a polarisation of this form.

We will refer to $M(X)$ as the *double* of X and the *doubling construction* as the function

$$\mathcal{D}: \text{HT}(G, n) \rightarrow \{\text{closed oriented smooth } 2n\text{-manifolds}\} / \simeq, \quad X \mapsto M(X).$$

We refer to the manifolds arising via this construction as *doubled (G, n) -complexes* and denote the set of all such manifolds up to homotopy equivalence by $d\text{HT}(G, n) := \text{im}(\mathcal{D})$. The doubles of minimal finite (G, n) -complexes could then be denoted by $d\text{HT}_{\min}(G, n)$, but instead we will write $\mathcal{M}_{2n}(G) := \mathcal{D}(\text{HT}_{\min}(G, n))$ to simplify the notation, as in the Introduction.

This map is referred to as the doubling construction since it has the following equivalent form. By [42, Proposition II.2], there exists a $2n$ -thickening $L(X)$ of X (which need not embed in \mathbb{R}^{2n}) such that

$$M(X) \cong_{\text{hCob}} \partial(L(X) \times [0, 1]) \cong L(X) \cup -L(X)$$

where \cong_{hCob} denotes h -cobordism and $L(X) \cup -L(X)$ is the double of $L(X)$ along its boundary. Note that more general notions of double exist (see, for example, [29, 49]); our notion is sometimes referred to as a trivial double elsewhere in the literature.

The following is [42, Proposition I.1]. Since it is illuminating, we include a short proof below.

Proposition 5.1. *If X and Y are finite (G, n) -complexes such that $\chi(X) = \chi(Y)$, then $M(X)$ and $M(Y)$ are stably diffeomorphic.*

Proof. Since $\chi(X) = \chi(Y)$, there exists $r \geq 0$ such that $X' = X \vee rS^n \simeq Y' = Y \vee rS^n$ are simple homotopy equivalent (see [70]). Then by [66, Corollary 2.1], $N(Y')$ embeds in $N(X')$, and the region $W = N(X') - N(Y')$ is an s -cobordism between $M(X')$ and $M(Y')$, which implies that $M(X')$ and $M(Y')$ are stably diffeomorphic (or diffeomorphic if $n > 2$). But $M(X') \cong M(X) \# r(S^n \times S^n)$ and $M(Y') \cong M(Y) \# r(S^n \times S^n)$ by construction, and so $M(X)$ and $M(Y)$ are stably diffeomorphic. \square

We will now restrict to the case where G is a finite group. In this case, we can identify the equivariant intersection form $S_{M(X)}$ as follows.

Proposition 5.2. *Let G be a finite group and $\varepsilon = (-1)^n$. If X is a finite (G, n) -complex, then*

$$\pi_n(M(X)) \cong \pi_n(X)^* \oplus \pi_n(X)$$

and $S_{M(X)} \cong \text{Met}_\varepsilon(\pi_n(X), \phi)$ are isometric for some $\phi \in \text{Sym}_\varepsilon(\pi_n(X))$ with $\phi^G = 0$.

This is a consequence of [42, Proposition II.2]. The property that $\phi^G = 0$ is not stated explicitly and so we include further details on this below. Recall that $\text{Sym}_\varepsilon(L)$ denotes the ε -symmetric forms on a $\mathbb{Z}G$ -module L , which admit elements $g \in G$ as isometries (see Definition 4.2).

Proof. By [42, Proposition II.4], the equivariant intersection form

$$S_{M(X)} \cong \text{Met}(\pi_n(X), \phi),$$

restricted to the summand $0 \oplus \pi_n(X)$, defines a form $\phi \in \text{Sym}_\varepsilon(\pi_n(X))$. It remains to show that $\phi^G = 0$. The $2n$ -thickening $L(X)$ can be constructed explicitly as follows (see the proof of [42, Proposition II.2 (i)]). Let $N(X^{(n-1)}) \subseteq \mathbb{R}^{2n}$ be a $2n$ -thickening of the $(n-1)$ -skeleton of X . For each n -cell of X , the attaching map $f_i: S^{n-1} \rightarrow X^{(n-1)}$ can be shown to induce an embedding $g_i: S^{n-1} \times D^n \hookrightarrow \partial N(X^{(n-1)})$. Then form $L(X)$ by attaching n -handles to $N(X^{(n-1)})$ along the embeddings g_i for each n -cell of X . It follows that $N(X) = L(X) \times I$ and $M(X) = L(X) \cup -L(X)$ (compare [29, Lemma 3.3] for uniqueness of the thickenings).

By an appropriate choice of these embeddings, one can ensure that the ordinary intersection form

$$S_X: H_n(L(X); \mathbb{Z}) \otimes H_n(L(X); \mathbb{Z}) \rightarrow \mathbb{Z}$$

is zero. This part of the proof is carefully explained in [42, Proposition II.4(ii), pp17-18]. By the transfer map on rational homology, we have

$$H_n(M(X); \mathbb{Z}G)^G \otimes \mathbb{Q} = H_n(\widetilde{M(X)}; \mathbb{Q})^G = H_n(M(X); \mathbb{Q}) = H_n(L(X); \mathbb{Q}) \oplus H_n(-L(X); \mathbb{Q}).$$

Since the intersection forms respect this splitting, the restriction ϕ of $S_{M(X)}$ has the property that $\phi^G \otimes \mathbb{Q} = S_X \otimes \mathbb{Q} = 0$. It follows that $\phi^G = 0$, as required. \square

The following will lead to our algebraic model for doubled (G, n) -complexes. A similar model was given by Kreck-Schafer in [42, Section III.1].

Definition 5.3. For $n \geq 2$ and $\varepsilon = (-1)^n$, let $\mathfrak{C}_n^{\text{alg}}(G)$ denote the category whose objects consist of pairs (C, ϕ) , where $C \in \text{Alg}(G, n)$ and $\phi \in \text{Sym}_\varepsilon(H_n(C))$, such that $\phi^G = 0$. A morphism $(C, \phi) \rightarrow (C', \phi')$ is a pair of chain maps $f: C \rightarrow C'$ and $g: C' \rightarrow C$ inducing the identity on H_0 .

The objects of $\mathfrak{C}_n^{\text{alg}}(G)$ admit an $\text{Aut}(G)$ -action where, for $\theta \in \text{Aut}(G)$ and $(C, \phi) \in \mathfrak{C}_n^{\text{alg}}(G)$, we define $(C, \phi)_\theta = (C_\theta, \phi_\theta)$ where C_θ and ϕ_θ are as defined in Sections 2 and 4b respectively.

Definition 5.4. For $n \geq 2$ and $\varepsilon = (-1)^n$, let $\mathfrak{M}_{2n}^{\text{alg}}(G)$ denote the category whose objects are pairs (D, Φ) where $D = (D_*, \partial_*)$ is a chain complex of (finitely generated) free $\mathbb{Z}G$ -modules D_* equipped with choices of $\mathbb{Z}G$ -module isomorphisms $H_0(D) \cong \mathbb{Z}$ and $H_{2n}(D) \cong \mathbb{Z}$ such that

- (i) $D_i = 0$ for $i < 0$ or $i > 2n$.
- (ii) $H_i(D) = 0$ for $0 < i < n$ and $n < i < 2n$.
- (iii) $\Phi \in \text{Sym}_\varepsilon(H_n(D))$ is a non-singular ε -symmetric form.

A morphism $(D, \Phi) \rightarrow (D', \Phi')$ is a chain map $h: D \rightarrow D'$ such that $H_0(h) = \text{id}_{\mathbb{Z}}$ and $H_{2n}(h) = \text{id}_{\mathbb{Z}}$, and $H_n(h)$ induces an isometry $H_n(h)^\wedge: \hat{\Phi} \rightarrow \hat{\Phi}'$ of the induced Tate forms $(\hat{H}^0(G; H_n(D)), \hat{\Phi})$ and $(\hat{H}^0(G; H_n(D')), \hat{\Phi}')$.

Two objects $(D, \Phi), (D', \Phi') \in \mathfrak{M}_{2n}^{\text{alg}}(G)$ are said to be *homotopy equivalent* if there exists a chain homotopy equivalence $h: D \rightarrow D'$ such that $H_0(h) = \text{id}_{\mathbb{Z}}$ and $H_{2n}(h) = \text{id}_{\mathbb{Z}}$, and $H_n(h)$ defines an isometry $\Phi \rightarrow \Phi'$.

The objects in $\mathfrak{M}_{2n}^{\text{alg}}(G)$ admit an $\text{Aut}(G)$ -action where, for $\theta \in \text{Aut}(G)$ and $(D, \Phi) \in \mathfrak{M}_{2n}^{\text{alg}}(G)$, we define $(D, \Phi)_\theta = (D_\theta, \Phi_\theta)$. Here, if $D = (D_*, \partial_*)$, then $D_\theta = ((D_*)_\theta, \partial_*)$ similarly to the $\text{Aut}(G)$ -action on $\text{Alg}(G, n)$. The form Φ_θ is as defined in Section 4b.

Definition 5.5. For an object $(C, \phi) \in \mathfrak{C}_n^{\text{alg}}(G)$, we define a chain complex $D_* = M(C_*)$, called the *algebraic 2n-double*, as follows

- (i) $D_i = C_i$ for $0 \leq i \leq n-1$, $D_i = C_{2n-i}^*$ for $n+1 \leq i \leq 2n$, and $D_n = C_n^* \oplus C_n$.
- (ii) $\partial_i^D = \partial_i^C$ for $0 \leq i \leq n-1$, $\partial_i^D = \partial_{2n-i+1}^*$ for $n+2 \leq i \leq 2n$,
 $\partial_n^D = \begin{pmatrix} 0 \\ \pm \partial_n^C \end{pmatrix}: C_n^* \oplus C_n \rightarrow C_{n-1}$, and $\partial_{n+1}^D = (\pm \partial_n^*, 0): C_{n-1}^* \rightarrow C_n^* \oplus C_n$.
- (iii) The identifications $H_0(D) \cong \mathbb{Z}$ and $H_{2n}(D) \cong \mathbb{Z}$ are induced by the identification $H_0(C) \cong \mathbb{Z}$.

For $n \geq 2$ and $\varepsilon = (-1)^n$, we define a functor $M: \mathfrak{C}_n^{\text{alg}}(G) \rightarrow \mathfrak{M}_{2n}^{\text{alg}}(G)$ given on objects by $M(C, \phi) := (D, \Phi)$, where $D_* = M(C_*)$ and $\Phi = \text{Met}_\varepsilon(H_n(C), \phi)$. If $f: C \rightarrow C'$ and $g: C' \rightarrow C$ are morphisms in $\mathfrak{C}_n^{\text{alg}}(G)$, then $M(f, g) := h$ where $h_i = f_i$, $0 \leq i < n$; $h_i = g_i^*$, $n+1 \leq i \leq n$ and $h_n = g_n^* \oplus f_n$. See Proposition 6.1(i) for the proof that this defines a morphism in $\mathfrak{M}_{2n}^{\text{alg}}(G)$.

Definition 5.6. Define $d\text{Alg}(G, n)$ to be the set of homotopy types of algebraic 2n-doubles $M(C, \phi)$ for $(C, \phi) \in \mathfrak{C}_n^{\text{alg}}(G)$. Define the *algebraic doubling construction* to be the map

$$\mathcal{D}^{\text{alg}}: \text{Alg}(G, n) \rightarrow d\text{Alg}(G, n), \quad C \mapsto M(C, 0) = (M(C), H_\varepsilon(H_n(C)))$$

where $H_\varepsilon(H_n(C))$ denotes the hyperbolic form on the module $H_n(C)^* \oplus H_n(C)$.

We also define $\mathcal{M}_{2n}^{\text{alg}}(G)$ to be the set of homotopy types of algebraic 2n-doubles $M(C, \phi)$ for $(C, \phi) \in \mathfrak{C}_n^{\text{alg}}(G)$ such that $(-1)^n \chi(C) = \chi_{\min}(G, n)$; that is, $C \in \text{Alg}_{\min}(G, n)$. Note that we could alternatively write this as $d\text{Alg}_{\min}(G, n)$, but we use $\mathcal{M}_{2n}^{\text{alg}}(G)$ to simplify notation.

We will now introduce two new equivalence relations on the objects in $\mathfrak{M}_{2n}^{\text{alg}}(G)$. Both equivalence relations refine homotopy equivalence, so induce a priori weaker equivalence relations on the set of algebraic 2n-doubles $d\text{Alg}(G, n)$.

Definition 5.7. Let $(D, \Phi), (D', \Phi') \in \mathfrak{M}_{2n}^{\text{alg}}(G)$ be two objects. A morphism $\Phi \rightarrow \Phi'$ is said to be an *integral isometry* if it induces an isometry $\Phi^G \rightarrow (\Phi')^G$, and a *Tate isometry* if it induces an isometry $\hat{\Phi} \rightarrow \hat{\Phi}'$. Note that integral isometries are Tate isometries.

If there exists a chain homotopy equivalence $h: D \rightarrow D'$ such that $H_0(h) = \text{id}_{\mathbb{Z}}$, $H_{2n}(h) = \text{id}_{\mathbb{Z}}$, and $H_n(h)$ defines a morphism $\Phi \rightarrow \Phi'$ which is an integral isometry (resp. Tate isometry), then we write $(D, \Phi) \simeq_{\mathbb{Z}} (D', \Phi')$ (resp. $(D, \Phi) \simeq_{\widehat{\mathbb{Z}}} (D', \Phi')$).

For $(C, \phi) \in \mathfrak{C}_n^{\text{alg}}(G)$, it can be shown that $M((C, \phi)_{\theta}) = M(C, \phi)_{\theta}$. Note that $(M_{\theta})^* \cong (M^*)_{\theta}$ if M is a $\mathbb{Z}G$ -lattice (see, for example, [53, Section 6.1]). In particular, \mathcal{D}^{alg} induces a map on the orbits under the $\text{Aut}(G)$ -actions

$$\mathcal{D}^{\text{alg}}: \text{Alg}(G, n) / \text{Aut}(G) \rightarrow d\text{Alg}(G, n) / \text{Aut}(G).$$

Recall that, by Proposition 2.3, there is an injective map $\mathcal{C}: \text{HT}(G, n) \hookrightarrow \text{Alg}(G, n) / \text{Aut}(G)$ given by $X \mapsto C_*(\widetilde{X})$. Similarly, by [42, Proposition II.3], we have a map

$$d(\mathcal{C}): d\text{HT}(G, n) \rightarrow d\text{Alg}(G, n) / \text{Aut}(G), \quad M \mapsto (C_*(\widetilde{M}), S_M).$$

The action of $\text{Aut}(G)$ on $d\text{Alg}(G, n)$ induces an action on $d\text{Alg}(G, n) / \simeq_{\mathbb{Z}}$, so there is a bijection

$$(d\text{Alg}(G, n) / \text{Aut}(G)) / \simeq_{\mathbb{Z}} \rightarrow (d\text{Alg}(G, n) / \simeq_{\mathbb{Z}}) / \text{Aut}(G).$$

The following observation will be crucial in our definition of the quadratic bias in Section 6.

Proposition 5.8. *Let G be a finite group. If X is a finite (G, n) -complex, then*

$$(C_*(\widetilde{M(X)}), S_{M(X)}) \simeq_{\mathbb{Z}} (C_*(\widetilde{M(X)}), H_{\varepsilon}(\pi_n(X))).$$

That is, there is a commutative diagram:

$$\begin{array}{ccc} \text{HT}(G, n) & \xrightarrow{\mathcal{C}} & \text{Alg}(G, n) / \text{Aut}(G) \\ \downarrow \mathcal{D} & & \downarrow \mathcal{D}^{\text{alg}} \\ d\text{HT}(G, n) & \xrightarrow{d(\mathcal{C})} & (d\text{Alg}(G, n) / \text{Aut}(G)) / \simeq_{\mathbb{Z}}. \end{array}$$

Proof. For $X \in \text{HT}(G, n)$, we have $(d(\mathcal{C}) \circ \mathcal{D})(X) = (C_*(\widetilde{M(X)}), S_{M(X)})$. By Proposition 5.2, there is an isometry $S_{M(X)} \cong \text{Met}_{\varepsilon}(\pi_n(X), \phi)$ for some $\phi \in \text{Sym}_{\varepsilon}(\pi_n(X))$ with $\phi^G = 0$. Next, we have $(\mathcal{D}^{\text{alg}} \circ \mathcal{C})(X) = (M(C_*(\widetilde{X})), H_{\varepsilon}(\pi_n(X)))$. By [42, Proposition II.3], we have that $C_*(\widetilde{M(X)}) \cong M(C_*(\widetilde{X}))$ are chain isomorphic. Let $\Phi = \text{Met}_{\varepsilon}(\pi_n(X), \phi)$ and $L = \pi_n(X)$, so that $H_{\varepsilon}(\pi_n(X)) \cong e_L$. Since $\phi^G = 0$, Proposition 4.12 implies that $\Phi^G \cong e_L^G$. Hence $(C_*(\widetilde{M(X)}), \Phi) \simeq_{\mathbb{Z}} (C_*(\widetilde{M(X)}), e_L)$ via the identity map on $C_*(\widetilde{M(X)})$, as required. \square

6. THE QUADRATIC BIAS INVARIANT

Throughout this section, we will fix $n \geq 2$ and a finite group G . We will now introduce the quadratic bias invariant, which is a homotopy invariant for the class of doubles $M(X)$ for X a finite (G, n) -complex.

In Section 3, we defined the bias invariant for an arbitrary finite (G, n) -complex X (see Definition 3.17). It followed from Proposition 3.10 that the bias invariant vanishes if X is non-minimal; in fact, the obstruction group $B(G, n, \chi)$ is trivial in this case. The construction of the quadratic bias invariant factors through the bias invariant and so, in the general setting, has an obstruction group $B_Q(G, n, \chi)$ which is a quotient of $B(G, n, \chi)$. This consequently vanishes for $\chi > \chi_{\min}(G, n)$. For notational simplicity, we will restrict to *minimal* finite (G, n) -complexes from now on (see Definition 2.4), and define the quadratic bias invariant only in this case.

We will begin with a polarised version of the invariant in the more general setting of algebraic $2n$ -doubles $M(C, \phi) \in \mathfrak{M}_{2n}^{\text{alg}}(G)$. In Section 6b, we return to manifolds and prove Theorem A.

6a. The quadratic bias for algebraic $2n$ -doubles. For a $\mathbb{Z}G$ -module L , recall that there is a canonical identification $(L^*)^{\wedge} \cong (\widehat{L})^*$. For $(C, \phi) \in \mathfrak{C}_n^{\text{alg}}(G)$, we let $L = H_n(C)$ and denote the evaluation form on $L^* \oplus L$ by $e := e_L$. Recall that, if $(D, \Phi) = M(C, \phi)$, then Proposition 4.12

implies that $\Phi^G \cong e^G$ and $\widehat{\Phi} \cong \widehat{e}$. For the remainder of this section, we will use the identifications $(L^*)^\wedge \cong (\widehat{L})^*$, $\Phi^G \cong e^G$ and $\widehat{\Phi} \cong \widehat{e}$ without further mention.

The following is a slight extension of [42, Propositions III.3 & III.4]. We can view it as the analogue of Proposition 3.6. The notation $\text{Diag}(\widehat{e}_1, \widehat{e}_2)$ for the set of *diagonal isometries* is given in Definition 4.19.

Proposition 6.1. *For $i = 1, 2$, let $(D_i, \Phi_i) = M(C_i, \phi_i)$, for $(C_i, \phi_i) \in \mathfrak{C}_n^{\text{alg}}(G)$ such that $\chi(C_1) = \chi(C_2)$. Let $L_i = H_n(C_i)$ and $e_i = e_{L_i}$, so that $\widehat{\Phi}_i \cong \widehat{e}_i$ for $i = 1, 2$. Then:*

- (i) *There exists a chain map $h: D_1 \rightarrow D_2$ such that $H_0(h) = \text{id}_{\mathbb{Z}}$, $H_{2n}(h) = \text{id}_{\mathbb{Z}}$ and*

$$H_n(h)^\wedge: (L_1^*)^\wedge \oplus \widehat{L}_1 \rightarrow (L_2^*)^\wedge \oplus \widehat{L}_2$$

is a diagonal isometry from \widehat{e}_1 to \widehat{e}_2 . Furthermore, we can take $h = M(f, g)$ where $f: C_1 \rightarrow C_2$ and $g: C_2 \rightarrow C_1$ are any chain maps such that $H_0(f) = \text{id}_{\mathbb{Z}}$ and $H_0(g) = \text{id}_{\mathbb{Z}}$.

- (ii) *Let $h: D_1 \rightarrow D_2$ be a chain map such that $H_0(h) = \text{id}_{\mathbb{Z}}$, $H_{2n}(h) = \text{id}_{\mathbb{Z}}$ and $H_n(h)^\wedge \in \text{Diag}(\widehat{e}_1, \widehat{e}_2)$. Then $H_n(h)^\wedge$ is independent of the choice of $h: D_1 \rightarrow D_2$. We will write this as $I(D_1, D_2) = (\nu(D_1, D_2)^*)^{-1} \oplus \nu(D_1, D_2) \in \text{Diag}(\widehat{e}_1, \widehat{e}_2)$ where $\nu(D_1, D_2) \in \text{Iso}(\widehat{L}_1, \widehat{L}_2)$.*
- (iii) *More generally, let $h: D_1 \rightarrow D_2$ be a chain map such that $H_0(h) = \text{id}_{\mathbb{Z}}$, $H_{2n}(h) = \text{id}_{\mathbb{Z}}$ and $H_n(h)^\wedge \in \text{Isom}(\widehat{e}_1, \widehat{e}_2)$. Then*

$$H_n(h)^\wedge = I(D_1, D_2) \circ \begin{pmatrix} \text{id} & \alpha \\ 0 & \text{id} \end{pmatrix}$$

for some $\alpha: \widehat{L}_1 \rightarrow (\widehat{L}_1)^$, where $\begin{pmatrix} \text{id} & \alpha \\ 0 & \text{id} \end{pmatrix}: (L_1^*)^\wedge \oplus \widehat{L}_1 \rightarrow (L_1^*)^\wedge \oplus \widehat{L}_1$, $(x, y) \mapsto (x + \alpha(y), y)$.*

Proof. (i) We combine parts of the arguments in [42, Propositions III.3 & III.4] to verify these statements. Since $D_i = M(C_i, \phi_i)$, there exists a chain map $f: C_1 \rightarrow C_2$ such that $H_0(f) = \text{id}_{\mathbb{Z}}$, which induces an isomorphism $f_*: H_n(C_1)^\wedge \rightarrow H_n(C_2)^\wedge$. Similarly, there exists a chain map $g: C_2 \rightarrow C_1$ such that $H_0(g) = \text{id}_{\mathbb{Z}}$, which induces an isomorphism $g_*: H_n(C_2)^\wedge \rightarrow H_n(C_1)^\wedge$. The chain map $h = M(f, g)$ has the required properties. In particular, $H_n(h)^\wedge \in \text{Diag}(\widehat{e}_1, \widehat{e}_2)$, so that $h: (D_1, \Phi_1) \rightarrow (D_2, \Phi_2)$ gives a morphism in the category $\mathfrak{M}_{2n}^{\text{alg}}(G)$.

(ii)/(iii) Let $h: D_1 \rightarrow D_2$ be a chain map such that $H_0(h) = \text{id}_{\mathbb{Z}}$, $H_{2n}(h) = \text{id}_{\mathbb{Z}}$ and $H_n(h)^\wedge \in \text{Isom}(\widehat{e}_1, \widehat{e}_2)$. By (i), there exists a chain map $h_0: D_1 \rightarrow D_2$ such that $H_0(h_0) = \text{id}_{\mathbb{Z}}$, $H_{2n}(h_0) = \text{id}_{\mathbb{Z}}$ and $I := H_n(h_0)^\wedge \in \text{Diag}(\widehat{e}_1, \widehat{e}_2)$. It follows from the arguments given in [42, Proposition III.4] that $H_n(h)^\wedge = I \circ \begin{pmatrix} \text{id} & \alpha \\ 0 & \text{id} \end{pmatrix}$ for some $\alpha: \widehat{L}_1 \rightarrow (\widehat{L}_1)^*$. The argument applies since the equivariant intersection form $\Phi_i = \text{Met}_\varepsilon(H_n(C_i), \phi_i)$ has $\phi_i^G = 0$, implying that the Tate forms $\widehat{\Phi}_i$ are hyperbolic, i.e. $\widehat{\Phi}_i = \widehat{e}_i$.

To prove (ii), suppose that $H_n(h)^\wedge \in \text{Diag}(\widehat{e}_1, \widehat{e}_2)$. If $I = (\nu^*)^{-1} \oplus \nu = \begin{pmatrix} (\nu^*)^{-1} & 0 \\ 0 & \nu \end{pmatrix}$ for some $\nu \in \text{Iso}(\widehat{L}_1, \widehat{L}_2)$, then $H_n(h)^\wedge = \begin{pmatrix} (\nu^*)^{-1} & (\nu^*)^{-1} \circ \alpha \\ 0 & \nu \end{pmatrix}$. Since this is diagonal, we have $(\nu^*)^{-1} \circ \alpha = 0$ and so $\alpha = 0$. Hence $H_n(h)^\wedge = I$ and so $I = I(D_1, D_2)$ is independent of the choice of h . (iii) now follows immediately by returning to the general case. \square

The following is a consequence of [42, Proposition III.5]. Note that the definition of $\text{Diag}(\widehat{e}_1, \widehat{e}_2)$ given in Section 4e is a priori different to the one given in [42]. However, since the lattices L_1 and L_2 are $|G|$ -locally isomorphic (see, for example, the proof of [42, Proposition III.5]), it follows from Proposition 4.21 that these definitions are equivalent.

Lemma 6.2. *Under the assumptions in Proposition 6.1, there exists a diagonal isometry $\varphi: e_1^G \rightarrow e_2^G$ inducing a diagonal isometry $\widehat{\varphi}: \widehat{e}_1 \rightarrow \widehat{e}_2$. That is, $\text{im}(\text{Diag}_{\Psi_1, \Psi_2}(e_1^G, e_2^G)) \cap \text{Diag}(\widehat{e}_1, \widehat{e}_2) \neq \emptyset$.*

We will now restrict to the case of minimal complexes. Fix $X \in \text{HT}_{\min}(G, n)$, $L = \pi_n(X)$ and $\mathcal{D}^{\text{alg}}(C_*(\widetilde{X})) = (D, e)$ where $D = C_*(\widetilde{M}(X))$ and $e = e_L \cong H_\varepsilon(L)$. We will refer to $(D, e) \in \mathcal{M}_{2n}^{\text{alg}}(G)$ as the *reference minimal algebraic $2n$ -double*. See Definition 5.6 for the definition of $\mathcal{M}_{2n}^{\text{alg}}(G)$. We often write this as (D, L, e) when L is not clear from the context.

For $i = 1, 2$, let (D_i, L_i, e_i) be as in Proposition 6.1 and such that $\chi(D_1) = \chi(D_2) = \chi(D)$. By Lemma 6.2, there exists $\bar{\tau}_{D_i} \in \text{Diag}(e^G, e_i^G)$ inducing $\tau_{D_i} \in \text{Diag}(\widehat{e}, \widehat{e}_i)$. Fix these reference isometries once and for all.

We are now ready to define the quadratic bias invariant, first in the setting of algebraic $2n$ -doubles. Recall from Section 4e that $\text{Diag}(\widehat{e}) \leq \text{Isom}(\widehat{e})$ denotes the set of diagonal isometries and $\text{Tri}(\widehat{e}) \leq \text{Isom}(\widehat{e})$ the set of triangular isometries. For subgroups $A, B \leq C$, we define $A \cdot B = \{ab : a \in A, b \in B\} \subseteq C$.

Definition 6.3 (Quadratic bias invariant for algebraic $2n$ -doubles). Fix a reference minimal doubled complex (D, e) . Define the *polarised quadratic bias obstruction group* to be

$$PB_Q(G, n) := \frac{\text{Diag}(\widehat{e})}{[\text{im}(\text{Isom}_\Psi(e^G)) \cdot \text{Tri}(\widehat{e})] \cap \text{Diag}(\widehat{e})}.$$

When $n = 2$, we write $P_Q(G) := P_Q(G, 2)$.

For $i = 1, 2$, let $(D_i, e_i) \in \mathcal{M}_{2n}^{\text{alg}}(G)$ such that $e_i = e_{L_i}$ for some module L_i . Define the *quadratic bias invariant* to be:

$$\beta_Q((D_1, e_1), (D_2, e_2)) := [\tau_{D_2}^{-1} \circ I(D_1, D_2) \circ \tau_{D_1}] \in PB_Q(G, n)$$

where the τ_{D_i} are as defined above and $[\cdot] : \text{Diag}(\widehat{e}) \rightarrow PB_Q(G, n)$ is the quotient map.

We will now establish the following two propositions.

Proposition 6.4. $[\text{im}(\text{Isom}_\Psi(e^G)) \cdot \text{Tri}(\widehat{e})] \cap \text{Diag}(\widehat{e})$ is a normal subgroup of $\text{Diag}(\widehat{e})$. In particular, $PB_Q(G, n)$ is a well-defined abelian group.

Proposition 6.5. $\beta_Q((D_1, e_1), (D_2, e_2)) \in PB_Q(G, n)$ does not depend on the choice of representatives $(D_i, e_i) \in \mathcal{M}_{2n}^{\text{alg}}(G)$ and isometries $\tau_{D_i} \in \text{Diag}(\widehat{e}, \widehat{e}_i)$. In particular, if $(\overline{D}, \overline{e})$ is a reference minimal algebraic $2n$ -double, then the quadratic bias invariant defines a map

$$\beta_Q : \{(D, e) \in \mathcal{M}_{2n}^{\text{alg}}(G) : \chi(D) = \chi(\overline{D})\} \rightarrow PB_Q(G, n), \quad (D, e) \mapsto \beta_Q((D, e), (\overline{D}, \overline{e})).$$

Furthermore, β_Q is an invariant of algebraic $2n$ -doubles up to the equivalence relation $\simeq_{\mathbb{Z}}$. That is, if $(D_1, e_1) \simeq_{\mathbb{Z}} (D_2, e_2)$, then $\beta_Q((D_1, e_1)) = \beta_Q((D_2, e_2)) \in PB_Q(G, n)$.

We will begin with the proof of Proposition 6.4. Recall that two subgroups $A, B \leq C$ commute if $A \cdot B = B \cdot A$. If so, then it follows that $A \cdot B$ is a subgroup of C .

Lemma 6.6. In the notation above, we have:

$$[\text{im}(\text{Isom}_\Psi(e^G)) \cdot \text{Tri}(\widehat{e})] \cap \text{Diag}(\widehat{e}) = [((\text{Diag}(\widehat{e}) \cdot \text{Tri}(\widehat{e})) \cap \text{im}(\text{Isom}_\Psi(e^G))) \cdot \text{Tri}(\widehat{e})] \cap \text{Diag}(\widehat{e}).$$

Proof. The inclusion \supseteq is clear and so it suffices to prove \subseteq . Let $\varphi \in [\text{im}(\text{Isom}_\Psi(e^G)) \cdot \text{Tri}(\widehat{e})] \cap \text{Diag}(\widehat{e})$. Then $\varphi \in \text{Diag}(\widehat{e})$ and $\varphi = \rho_1 \circ \rho_2$ for some $\rho_1 \in \text{im}(\text{Isom}_\Psi(e^G))$ and $\rho_2 \in \text{Tri}(\widehat{e})$. We have $\rho_1 = \varphi \circ \rho_2^{-1} \in \text{Diag}(\widehat{e}) \cdot \text{Tri}(\widehat{e})$ which implies that $\rho_1 \in (\text{Diag}(\widehat{e}) \cdot \text{Tri}(\widehat{e})) \cap \text{im}(\text{Isom}_\Psi(e^G))$, and so $\varphi \in ((\text{Diag}(\widehat{e}) \cdot \text{Tri}(\widehat{e})) \cap \text{im}(\text{Isom}_\Psi(e^G))) \cdot \text{Tri}(\widehat{e})$, which completes the proof. \square

In order to prove this equivalent form is a subgroup of $\text{Diag}(\widehat{e})$, we will use:

Lemma 6.7. If $\varphi \in \text{Diag}(\widehat{e})$, then $\varphi \cdot \text{Tri}(\widehat{e}) = \text{Tri}(\widehat{e}) \cdot \varphi$. It follows that:

- (i) $\text{Diag}(\widehat{e})$ normalises $\text{Tri}(\widehat{e})$ in $\text{Isom}(\widehat{e})$, and so $\text{Diag}(\widehat{e}) \cdot \text{Tri}(\widehat{e}) \leq \text{Isom}(\widehat{e})$ is a subgroup.
- (ii) If $H \leq \text{Diag}(\widehat{e}) \cdot \text{Tri}(\widehat{e})$ is a subgroup, then $H \cdot \text{Tri}(\widehat{e}) \leq \text{Isom}(\widehat{e})$ is a subgroup.

Proof. Let $\varphi \in \text{Diag}(\widehat{e})$ and $\varphi_T \in \text{Tri}(\widehat{e})$. Then there exists an isomorphism $f : \widehat{L} \rightarrow \widehat{L}$ and a homomorphism $h : \widehat{L} \rightarrow (\widehat{L})^*$ such that $\varphi = \begin{pmatrix} (f^*)^{-1} & 0 \\ 0 & f \end{pmatrix}$ and $\varphi_T = \begin{pmatrix} \text{id} & h \\ 0 & \text{id} \end{pmatrix}$. Then we have

$$\varphi \circ \varphi_T = \begin{pmatrix} (f^*)^{-1} & 0 \\ 0 & f \end{pmatrix} \begin{pmatrix} \text{id} & h \\ 0 & \text{id} \end{pmatrix} = \begin{pmatrix} \text{id} & (f^*)^{-1} \circ h \circ f^{-1} \\ 0 & \text{id} \end{pmatrix} \begin{pmatrix} (f^*)^{-1} & 0 \\ 0 & f \end{pmatrix} = \varphi'_T \circ \varphi$$

where $\varphi'_T \in \text{Tri}(\widehat{e})$. Hence $\varphi \cdot \text{Tri}(\widehat{e}) \cdot \varphi^{-1} \subseteq \text{Tri}(\widehat{e})$, and equality follows by applying this to φ^{-1} .

Part (i) now follows immediately. To see part (ii), consider the normaliser subgroup

$$N_{\text{Isom}(\widehat{e})}(\text{Tri}(\widehat{e})) = \{\varphi \in \text{Isom}(\widehat{e}) : \varphi \cdot \text{Tri}(\widehat{e}) = \text{Tri}(\widehat{e}) \cdot \varphi\}.$$

Since $\text{Diag}(\widehat{e}), \text{Tri}(\widehat{e}) \leq N_{\text{Isom}(\widehat{e})}(\text{Tri}(\widehat{e}))$, we have $\text{Diag}(\widehat{e}) \cdot \text{Tri}(\widehat{e}) \leq N_{\text{Isom}(\widehat{e})}(\text{Tri}(\widehat{e}))$. \square

Proof of Proposition 6.4. We will start by showing that $[\text{im}(\text{Isom}_\Psi(e^G)) \cdot \text{Tri}(\widehat{e})] \cap \text{Diag}(\widehat{e})$ is a subgroup of $\text{Diag}(\widehat{e})$. By Lemma 6.7 (i), $\text{Diag}(\widehat{e}) \cdot \text{Tri}(\widehat{e})$ is a subgroup. Since $\text{im}(\text{Isom}_\Psi(e^G))$

is a subgroup, this gives that $H := (\text{Diag}(\widehat{e}) \cdot \text{Tri}(\widehat{e})) \cap \text{im}(\text{Isom}_\Psi(e^G))$ is a subgroup. Since $H \leq \text{Diag}(\widehat{e}) \cdot \text{Tri}(\widehat{e})$ is a subgroup, Lemma 6.7 (ii) now implies that $H \cdot \text{Tri}(\widehat{e})$ is a subgroup and so $(H \cdot \text{Tri}(\widehat{e})) \cap \text{Diag}(\widehat{e})$ is a subgroup. The result now follows by combining with Lemma 6.6.

To see that $K := [\text{im}(\text{Isom}_\Psi(e^G)) \cdot \text{Tri}(\widehat{e})] \cap \text{Diag}(\widehat{e})$ is a normal subgroup of $\text{Diag}(\widehat{e})$, note that $\text{im}(\text{Diag}_\Psi(e^G)) \leq K$. By combining Lemma 3.11 and Proposition 4.22 (see the discussion following Proposition 4.22), we have that $\text{im}(\text{Diag}_\Psi(e^G)) \trianglelefteq \text{Diag}(\widehat{e})$ is a normal subgroup and $\text{Diag}(\widehat{e}) / \text{im}(\text{Diag}_\Psi(e^G)) \cong (\mathbb{Z}/m)^\times / \{\pm 1\}$ is abelian, for some $m \geq 1$. This implies there is a quotient map $f: \text{Diag}(\widehat{e}) \twoheadrightarrow (\mathbb{Z}/m)^\times / \{\pm 1\}$ with $\ker(f) = \text{im}(\text{Diag}_\Psi(e^G))$.

Let $K' = f(K)$. Since $\ker(f) = \text{im}(\text{Diag}_\Psi(e^G)) \leq K$, we have that $f^{-1}(K') = K \cdot \ker(f) = K$. Since $(\mathbb{Z}/m)^\times / \{\pm 1\}$ is abelian, $K' \trianglelefteq (\mathbb{Z}/m)^\times / \{\pm 1\}$ is a normal subgroup. The preimage of a normal subgroup is normal, and so $K \trianglelefteq \text{Diag}(\widehat{e})$ is normal. Thus, $PB_Q(G, n)$ is a well-defined group. It is a quotient of $(\mathbb{Z}/m)^\times / \{\pm 1\}$ and so is abelian. \square

Proof of Proposition 6.5. We begin by noting that, given (D_1, e_1) and (D_2, e_2) , the class $[\tau_{D_2}^{-1} \circ I(D_1, D_2) \circ \tau_{D_1}] \in PB_Q(G, n)$ is independent of the choice of τ_{D_1} and τ_{D_2} . This follows from the fact that, if τ'_{D_1} and τ'_{D_2} are other choices, then the two classes would differ by multiplication by $(\tau'_{D_2})^{-1} \circ \tau_{D_2}$ and $\tau_{D_1}^{-1} \circ \tau'_{D_1}$, which are both in $\text{Isom}_\Psi(e^G) \cap \text{Diag}(\widehat{e})$.

Next suppose that, for $i = 1, 2$, there is a chain homotopy equivalence $h_i: D_i \rightarrow D'_i$ which induces an integral isometry. Thus $H_n(h_i)^\wedge \in \text{Isom}(\widehat{e}_i, (e'_i)^\wedge)$ is an isometry which lifts to an isometry $H_n(h_i)^G \in \text{Isom}(e_i^G, (e'_i)^G)$. By Proposition 6.1 (iii), we have that

$$H_n(h_i)^\wedge = I(D_i, D'_i) \circ \begin{pmatrix} \text{id} & \alpha_i \\ 0 & \text{id} \end{pmatrix}$$

for some $\alpha_i: \widehat{L}_i \rightarrow (\widehat{L}'_i)^*$ and where $I(D_i, D'_i)$ is as defined in Proposition 6.1. This implies that

$$A_i := \tau_{D'_i}^{-1} \circ I(D_i, D'_i) \circ \tau_{D_i} = (\tau_{D'_i}^{-1} \circ H_n(h_i)^\wedge \circ \tau_{D_i}) \circ (\circ \tau_{D_i}^{-1} \circ \begin{pmatrix} \text{id} & -\alpha_i \\ 0 & \text{id} \end{pmatrix} \circ \circ \tau_{D_i}).$$

We have $\tau_{D'_i}^{-1} \circ H_n(h_i)^\wedge \circ \tau_{D_i} = \Psi_*(\tau_{D'_i}^{-1} \circ H_n(h_i)^G \circ \tau_{D_i}) \in \text{im}(\text{Isom}_\Psi(e^G))$. Since $\tau_{D_i} \in \text{Diag}(\widehat{e}_i, \widehat{e})$, we have that $\tau_{D_i}^{-1} \circ \begin{pmatrix} \text{id} & -\alpha_i \\ 0 & \text{id} \end{pmatrix} \circ \tau_{D_i} \in \text{Tri}(\widehat{e})$ by the same argument as Lemma 6.7. In particular, we have that $A_i \in [\text{im}(\text{Isom}_\Psi(e^G)) \cdot \text{Tri}(\widehat{e})] \cap \text{Diag}(\widehat{e})$ for $i = 1, 2$.

It follows from Proposition 6.1 that $I(D_1, D_2) = I(D_2, D'_2)^{-1} \circ I(D'_1, D'_2) \circ I(D_1, D'_1)$. Hence we have

$$\tau_{D_2}^{-1} \circ I(D_1, D_2) \circ \tau_{D_1} = A_2^{-1} \circ (\tau_{D'_2}^{-1} \circ I(D'_1, D'_2) \circ \tau_{D'_1}) \circ A_1$$

and so $[\tau_{D_2}^{-1} \circ I(D_1, D_2) \circ \tau_{D_1}] = [\tau_{D'_2}^{-1} \circ I(D'_1, D'_2) \circ \tau_{D'_1}] \in PB_Q(G, n)$, as required. \square

Define $\widehat{\mathcal{D}}$ to be the surjective abelian group homomorphism given by the composition

$$\widehat{\mathcal{D}}: \frac{\text{Aut}(\widehat{L})}{\text{Aut}_\psi(L^G)} \xrightarrow{A^{-1}} \frac{\text{Diag}(\widehat{e})}{\text{Diag}_\Psi(e^G)} \twoheadrightarrow PB_Q(G, n)$$

where A is as defined in the discussion following Proposition 4.22. In particular, for $f \in \text{Aut}(\widehat{L})$, we have that $\widehat{\mathcal{D}}([f]) = [(f^*)^{-1} \oplus f]$ where $(f^*)^{-1} \oplus f = \begin{pmatrix} (f^*)^{-1} & 0 \\ 0 & f \end{pmatrix} \in \text{Diag}(\widehat{e})$.

We conclude this section by establishing the following relationship between the bias invariant for an algebraic n -complex C over $\mathbb{Z}G$ with $(-1)^n \chi(C) = \chi_{\min}(G, n)$ and the quadratic bias invariant for the corresponding algebraic $2n$ -double (D, e) . The following was asserted on [42, p34] though no argument was given.

Proposition 6.8. *Fix $C \in \text{Alg}_{\min}(G, n)$, let $L = \pi_n(C)$ and let $(D, e) = \mathcal{D}^{\text{alg}}(C) \in \mathcal{M}_{2n}^{\text{alg}}(G)$ denote the corresponding algebraic $2n$ -double. Then there is a commutative diagram of sets*

$$\begin{array}{ccc} \text{Alg}_{\min}(G, n) & \xrightarrow{\mathcal{D}^{\text{alg}}} & \mathcal{M}_{2n}^{\text{alg}}(G) \\ \beta(\cdot, C) \downarrow & & \downarrow \beta_Q(\cdot, (D, e)) \\ \frac{\text{Aut}(\widehat{L})}{\text{Aut}_\psi(L^G)} & \xrightarrow{\widehat{\mathcal{D}}} & PB_Q(G, n). \end{array}$$

Proof. Let $C' \in \text{Alg}_{\min}(G, n)$. We start by evaluating the bottom left composition. By the discussion in Section 3b, there exists an isomorphism $\bar{\tau}_C: H_n(C)^G \rightarrow H_n(C')^G$ inducing an isomorphism $\tau_C: H_n(C)^\wedge \rightarrow H_n(C')^\wedge$, and similarly for $\tau_{C'}$, and there exists a chain map $f: C \rightarrow C'$ such that $H_0(f) = \text{id}_{\mathbb{Z}}$. By Proposition 3.6, $H_n(f)^\wedge$ is an isomorphism and coincides with $\sigma(C, C')$. By definition, we have $\beta(C', C) = [F]$ where $F = \tau_C^{-1} \circ H_n(f)^\wedge \circ \tau_{C'}$. This implies that $\widehat{\mathcal{D}}(\beta(C', C)) = [(F^*)^{-1} \oplus F]$.

To evaluate the top right composition, first let $(D', e') = \mathcal{D}^{\text{alg}}(C')$ where $e' = e_{L'}$ for $L' = H_n(C')$. Let $\tau_D = (\tau_C^*)^{-1} \oplus \tau_C$, $\tau_{D'} = (\tau_{C'}^*)^{-1} \oplus \tau_{C'}$, and let $h = M(f, g)$ where $g: C' \rightarrow C$ is a chain map such that $H_0(g) = \text{id}_{\mathbb{Z}}$. By Proposition 6.1(i), $H_n(h)^\wedge$ is a diagonal isometry which coincides with $I(D, D')$. By definition, we have $\beta_Q((D', e'), (D, e)) = [\tau_D^{-1} \circ H_n(h)^\wedge \circ \tau_{D'}]$. We have $H_n(h)^\wedge = (H_n(g)^*)^\wedge \oplus H_n(f)^\wedge$. It is shown in [55, Lemma 1] that, if $f': C_1 \rightarrow C_2$ is a chain map with $H_0(f') = \text{id}_{\mathbb{Z}}$, then $H_n(f')^\wedge$ depends only on C_1 and C_2 and not on the map f' . It follows that, since $g \circ f: C \rightarrow C$ has $H_0(g \circ f) = \text{id}_{\mathbb{Z}}$, we have that $H_n(g \circ f)^\wedge = H_n(\text{id}_C)^\wedge = \text{id}$ and so $H_n(g)^\wedge = (H_n(f)^\wedge)^{-1}$, and similarly we can obtain $(H_n(g)^*)^\wedge = ((H_n(f)^*)^\wedge)^{-1}$. Hence $H_n(h)^\wedge = ((H_n(f)^*)^\wedge)^{-1} \oplus H_n(f)^\wedge$ and so $\beta_Q((D', e'), (D, e)) = [(F^*)^{-1} \oplus F]$, as required. \square

6b. The quadratic bias invariant for manifolds. Fix an integer $n \geq 2$ and a finite group G . The proof of Theorem A follows from Theorem 6.12 and Proposition 6.14 in this section. Recall from Section 5 that $\mathcal{M}_{2n}(G)$ denotes the set of homotopy types of doubles $M(X)$ for X a minimal finite (G, n) -complex.

We will now establish the analogue of Proposition 3.16 for the quadratic bias invariant. Recall the definitions of ρ and $\varphi_{(G, n)}$ from Lemma 3.11 and Proposition 3.16 respectively.

Proposition 6.9. *Let $n \geq 2$ and let G be a finite group. If X is a minimal finite (G, n) -complex and $(D, e) = \mathcal{D}^{\text{alg}}(C_*(\tilde{X}))$, then the map*

$$\Psi_{G, n}: \text{Aut}(G) \rightarrow PB_Q(G, n), \quad \theta \mapsto \beta_Q((D, e), (D_\theta, e_\theta))$$

is a group homomorphism and is independent of the choice of X . Furthermore, we have that $\Psi_{G, n} = \widehat{\mathcal{D}} \circ (\rho^{-1})_ \circ \varphi_{(G, n)}$.*

Proof. The equality $\Psi_{G, n} = \widehat{\mathcal{D}} \circ (\rho^{-1})_* \circ \varphi_{(G, n)}$ follows directly from Proposition 6.8 and the fact that, if $\theta \in \text{Aut}(G)$ and $C \in \text{Alg}(G, n)$ has $(D, e) = \mathcal{D}^{\text{alg}}(C)$, then $\mathcal{D}^{\text{alg}}(C_\theta) = (D_\theta, e_\theta)$. Since, by Proposition 3.16, $\varphi_{(G, n)}$ is a group homomorphism and is independent of the choice of X , the same therefore holds for $\Psi_{G, n}$. \square

Definition 6.10 (Quadratic bias invariant for doubled (G, n) -complexes). Let $n \geq 2$ and let G be a finite group. Define $D_Q(G, n)$ to be the image of the map $\Psi_{G, n}$ given in Proposition 6.9. Fix a reference minimal (G, n) -complex \bar{X} and let $(D, e) = \mathcal{D}^{\text{alg}}(C_*(\bar{X}))$. Define the *quadratic bias obstruction group* for doubled minimal (G, n) -complexes to be

$$B_Q(G, n) := \frac{PB_Q(G, n)}{D_Q(G, n)} = \frac{\text{Diag}(\widehat{e})}{([\text{im}(\text{Isom}_\Psi(e^G))] \cdot \text{Tri}(\widehat{e}) \cap \text{Diag}(\widehat{e})) \cdot D_Q(G, n)}.$$

When $n = 2$, we write $B_Q(G) := B_Q(G, 2)$.

Let X_1, X_2 be minimal finite (G, n) -complexes, let $M(X_1), M(X_2)$ be the doubles, and let $(D_1, e_1) = \mathcal{D}^{\text{alg}}(C_*(\tilde{X}_1))$, $(D_2, e_2) = \mathcal{D}^{\text{alg}}(C_*(\tilde{X}_2))$. Define the *quadratic bias invariant* to be

$$\beta_Q(M(X_1), M(X_2)) := [\beta_Q((D_1, e_1), (D_2, e_2))] \in B_Q(G, n)$$

where $[\cdot]: PB_Q(G, n) \twoheadrightarrow B_Q(G, n)$ is the quotient map and $\beta_Q((D_1, e_1), (D_2, e_2))$ denotes the quadratic bias invariant defined in the case of algebraic $2n$ -doubles in Definition 6.3.

The quotient $PB_Q(G, n)/D_Q(G, n)$ is well defined since $PB_Q(G, n)$ is an abelian group.

Remark 6.11. (i) It follows from the definition, as well as Proposition 6.9, that $D_Q(G, n)$ is the image of $D(G, n)$ under $\widehat{\mathcal{D}} \circ (\rho^{-1})_*$. In particular, there is a commutative diagram

$$\begin{array}{ccc} (\mathbb{Z}/m)^\times / \{\pm 1\} & \xrightarrow{(\rho^{-1})_*} \frac{\text{Aut}(\widehat{L})}{\text{Aut}_\psi(L^G)} & \xrightarrow{\widehat{\mathcal{D}}} PB_Q(G, n) \\ \downarrow & & \downarrow \\ B(G, n) & \xrightarrow{q} & B_Q(G, n). \end{array}$$

where $m = m_{(G, n)}$, $B(G, n) = (\mathbb{Z}/m)^\times / \pm D(G, n)$ and q is the natural quotient map.

(ii) In the Introduction, we considered (the $n = 2$ case of) the group

$$N(G, n) := \ker(q: B(G, n) \twoheadrightarrow B_Q(G, n)).$$

This is the image of $[\text{im}(\text{Isom}_\Psi(e^G)) \cdot \text{Tri}(\widehat{e})] \cap \text{Diag}(\widehat{e}) \leq \text{Diag}(\widehat{e})$ under the surjection

$$\text{Diag}(\widehat{e}) \twoheadrightarrow \frac{\text{Diag}(\widehat{e})}{\text{Diag}_\Psi(e^G)} \cong \frac{\text{Aut}(\widehat{L})}{\text{Aut}_\psi(L^G)} \cong (\mathbb{Z}/m)^\times / \{\pm 1\}$$

provided by (4.23).

Note that the quadratic bias invariant $\beta_Q(M(X_1), M(X_2))$ was defined using the algebraic $2n$ -doubles $\mathcal{D}^{\text{alg}}(C_*(\widetilde{X}_i)) = (C_*(\widetilde{M(X_i)}), H_\varepsilon(\pi_2(X_i)))$. The latter form does not necessarily coincide with the equivariant intersection form $S_{M(X_i)}$ and so is not clearly a function of the manifold $M(X_i)$ itself. In Proposition 5.8, we showed that there is an equivalence

$$d(\mathcal{C})(M(X)) = (C_*(\widetilde{M(X_i)}), S_{M(X_i)}) \simeq_{\mathbb{Z}} (C_*(\widetilde{M(X_i)}), H_\varepsilon(\pi_2(X_i))).$$

Since the quadratic bias invariant β_Q is a $\simeq_{\mathbb{Z}}$ invariant for algebraic $2n$ -doubles (Definition 5.7 and Proposition 6.5), it follows that $\beta_Q(M(X_1), M(X_2))$ depends only on $M(X_1)$ and $M(X_2)$ up to homotopy equivalence (or, more generally, up to $\simeq_{\mathbb{Z}}$).

In particular we have now established the following result which, alongside Proposition 6.14 below, implies Theorem A.

Theorem 6.12. $\beta_Q(M(X_1), M(X_2)) \in B_Q(G, n)$ depends only on the manifolds $M(X_1)$ and $M(X_2)$ up to homotopy equivalence. In particular, if \overline{X} is a reference minimal (G, n) -complex, then the quadratic bias invariant defines a map

$$\beta_Q: \mathcal{M}_{2n}(G) \rightarrow B_Q(G, n), \quad X \mapsto \beta_Q(X, \overline{X}).$$

Thus, β_Q is an invariant of doubled minimal (G, n) -complexes up to homotopy equivalence.

In fact, Proposition 6.5 actually proves the following stronger statement. Note that $\simeq_{\mathbb{Z}}$ can be viewed as an equivalence relation on manifolds via the map $d(\mathcal{C})$ defined in Section 5.

Proposition 6.13. Let $X_1, X_2 \in \text{HT}_{\min}(G, n)$. If $M(X_1) \simeq_{\mathbb{Z}} M(X_2)$, then $\beta_Q(M(X_1)) = \beta_Q(M(X_2))$. In particular, the quadratic bias invariant defines a map

$$\beta_Q: \mathcal{M}_{2n}(G) / \simeq_{\mathbb{Z}} \rightarrow B_Q(G, n).$$

The following is a consequence of Proposition 6.8. Here $q: B(G, n) \rightarrow B_Q(G, n)$ is the map defined in Remark 6.11 (i).

Proposition 6.14. There is a commutative diagram of sets

$$\begin{array}{ccc} \text{HT}_{\min}(G, n) & \xrightarrow{\mathcal{D}} & \mathcal{M}_{2n}(G) \\ \beta \downarrow & & \downarrow \beta_Q \\ B(G, n) & \xrightarrow{q} & B_Q(G, n) \end{array}$$

In particular, if $\beta: \text{HT}_{\min}(G, n) \rightarrow B(G, n)$ is surjective, then $\beta_Q: \mathcal{M}_{2n}(G) \rightarrow B_Q(G, n)$ is surjective.

Remark 6.15. If (G, n) does not satisfy the strong minimality hypothesis then, by Remark 3.19, the bias invariant is zero and hence the quadratic bias invariant is zero.

6c. Relationship between the quadratic 2-type and the quadratic bias. The quadratic 2-type of a closed oriented (smooth or topological) 4-manifold M is the quadruple

$$Q(M) = [\pi_1(M), \pi_2(M), k_M, S_M],$$

where $k_M \in H^3(\pi_1(M); \pi_2(M))$ denotes the k -invariant and $S_M: \pi_2(M) \times \pi_2(M) \rightarrow \mathbb{Z}[\pi_1(M)]$ denotes the equivariant intersection form. An isometry of two such quadruples is an isomorphism of pairs π_1, π_2 respecting the k -invariant and inducing an isometry on S (similar definitions apply for X a closed oriented Poincaré 4-complex).

The data $[\pi_1(M), \pi_2(M), k_M]$ determine the *algebraic 2-type* $B = B(M)$, which is the total space of a 2-stage Postnikov fibration $K(\pi_2(M), 2) \rightarrow B \rightarrow K(\pi_1(M), 1)$. A B -polarised oriented finite Poincaré 4-complex is a 3-equivalence $f: X \rightarrow B$. Let $\mathcal{S}_4^{PD}(B)$ denote the set of B -polarised homotopy types over B (see [26, §1] for more details).

We will now prove the following result from the Introduction (see Theorem 1.4).

Theorem 6.16. *The quadratic 2-type determines the quadratic bias invariant. More specifically, let G be a finite group and let $M_1, M_2 \in \mathcal{M}_4(G)$. If $Q(M_1) \cong Q(M_2)$, then $\beta_Q(M_1) = \beta_Q(M_2)$.*

The proof of Theorem 6.16 will be based on the following observation which is a direct consequence from the definition of β_Q . For convenience, we will work in the B -polarised setting.

Lemma 6.17. *Let G be a finite group and let $M_1, M_2 \in \mathcal{M}_4(G)$. If there is a B -polarised equivalence $(C_*(\widetilde{M}_1), S_{M_1}) \simeq (C_*(\widetilde{M}_2), S_{M_2})$, then $\beta_Q(M_1) = \beta_Q(M_2)$.*

In particular, it now suffices to prove the following, where $B = B(X)$ for some fixed finite Poincaré 4-complex X with $\pi_1(X) = G$ as reference.

Proposition 6.18. *For G a finite group, let X_1, X_2 be oriented finite B -polarised Poincaré 4-complexes with $Q(X_1) \cong Q(X_2)$. Then $(C_*(\widetilde{M}_1), S_{M_1})$ and $(C_*(\widetilde{M}_2), S_{M_2})$ are B -polarised homotopy equivalent.*

Proof. First note that we can write $X_1 \simeq K \cup_g D^4$ where K is a finite 3-complex and the attaching map $g: S^3 \rightarrow K$ is an element of $\pi_3(K)$. Since G is finite and $Q(X_1) \cong Q(X_2)$, it follows from [39, Theorem 1.5] that X_1 and X_2 differ by the action of an element $\alpha \in \text{Tors}(\mathbb{Z} \otimes_{\mathbb{Z}G} \Gamma(\pi_2(B)))$ where B is a 3-coconnected CW-complex which is 3-equivalent to both X_1 and X_2 . This action is described on [26, p89-90]. We have that $\Gamma(\pi_2(B)) \cong H_4(\widetilde{B})$. The action implies that $X_2 \simeq K \cup_{g+\alpha'} D^4$ where $\alpha' \in \pi_3(K)$ is obtained by choosing a preimage of $\alpha \in \text{Tors}(H_4(\widetilde{B}) \otimes_{\Lambda} \mathbb{Z})$ in $H_4(\widetilde{B})$ and mapping it under the composition $H_4(\widetilde{B}) \rightarrow H_4(\widetilde{B}, \widetilde{K}) \cong \pi_4(B, K) \rightarrow \pi_3(K)$.

It now suffices to show that there is an equivalence $(C_*(\widetilde{X}_1), S_{X_1}) \simeq (C_*(\widetilde{X}_2), S_{X_2})$. However, by construction the image of $\alpha' \in \pi_3(K)$ is a torsion element which vanishes in $H_3(\widetilde{K})$, and therefore does not affect the chain complexes. \square

The proof of Theorem 6.16. For a closed oriented 4-manifold M , consider the pair of invariants $(C_*(\widetilde{M}), S_M)$. For closed oriented 4-manifolds M and N , an isometry $Q(M) \cong Q(N)$ implies that $B(M) \simeq B(N)$, and we can work in the B -polarised setting with $B := B(M)$. We say that these pairs are B -polarised equivalent, which we write as $(C_*(\widetilde{M}), S_M) \simeq (C_*(\widetilde{N}), S_N)$, if there exists a chain homotopy equivalence $f: C_*(\widetilde{M}) \rightarrow C_*(\widetilde{N})$ over B such that the induced map $f_*: H_2(\widetilde{M}) \rightarrow H_2(\widetilde{N})$ is an isometry $S_M \rightarrow S_N$ under the identifications $H_2(\widetilde{M}) \cong \pi_2(M)$ and $H_2(\widetilde{N}) \cong \pi_2(N)$. This is a B -polarised homotopy invariant of 4-manifolds in $\mathcal{S}_4^{PD}(B)$. The proof now follows from Lemma 6.17 and Proposition 6.18. \square

7. EVALUATION OF THE QUADRATIC BIAS OBSTRUCTION GROUP

Let $n \geq 2$ and let G be a finite group. Recall from Section 3b that the invariant rank is defined to be $r_{(G,n)} = \text{rank}_{\mathbb{Z}}(L^G)$ where $L = \pi_n(X)$ for X any minimal (G, n) -complex. By Proposition 3.3, (G, n) satisfies the strong minimality hypothesis if and only if $r_{(G,n)} = d(H_n(G))$.

The main result of this section will be the following, which implies Theorem B.

Theorem 7.1. *Let $n \geq 2$ and let G be a finite group such that $H_n(G) \cong (\mathbb{Z}/m)^d$ for some $m \geq 1$ and $d = r_{(G,n)} \geq 3$. Then:*

(i) *If n is even, there are isomorphisms*

$$PB_Q(G, n) \cong \frac{(\mathbb{Z}/m)^\times}{\pm(\mathbb{Z}/m)^{\times 2}} \quad \text{and} \quad B_Q(G, n) \cong \frac{(\mathbb{Z}/m)^\times}{\pm(\mathbb{Z}/m)^{\times 2} \cdot D(G, n)}.$$

(ii) *If n is odd, then $PB_Q(G, n) = B_Q(G, n) = 0$.*

Remark 7.2. If $d \neq r_{(G,n)}$, then (G, n) does not satisfy the strong minimality hypothesis and Remark 3.19 implies that $B(G, n) = 0$ and so $B_Q(G, n) = 0$. Hence the above computes $B_Q(G, n)$ for all finite groups G such that $H_n(G) \cong (\mathbb{Z}/m)^d$ for some $m \geq 1, d \geq 3$.

This section will be structured as follows. In Section 7a, we will begin by giving a formulation of $PB_Q(G, n)$ in the case where $H_n(G) \cong (\mathbb{Z}/m)^d$ in terms of unitary groups (Proposition 7.3). In Section 7b, we will prove Theorem 7.1 in the case where n is even. The strategy will be to use the formulation in terms of unitary groups to establish a group homomorphism

$$\phi: PB_Q(G, n) \longrightarrow \frac{(\mathbb{Z}/m)^\times}{\pm(\mathbb{Z}/m)^{\times 2}}$$

by using the connection between unitary groups and algebraic L -theory. Evaluating the L -groups and showing that ϕ is an isomorphism then leads to the result. The proof of Theorem 7.1 in the case where n is odd is carried out in Section 7c.

Detailed background on unitary groups and algebraic L -theory can be found in Appendix A and the L -theory calculations are carried out in Appendix B.

7a. Formulation in terms of unitary groups. Let $(\Lambda, -)$ be a ring with involution, let $\varepsilon = (-1)^n$ for some n , and let $d \geq 1$. The *unitary group* $U_{2d}^\varepsilon(\Lambda)$ is the subgroup of $GL_{2d}(\Lambda)$ consisting of block matrices of the form $\sigma = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$ where $\alpha, \beta, \gamma, \delta \in M_d(\Lambda)$ are such that

$$(i) \quad \alpha\delta^* + (-1)^n\beta\gamma^* = I$$

$$(ii) \quad \alpha\beta^* \text{ and } \gamma\delta^* \text{ each have the form } \theta - (-1)^n\theta^* \text{ for some } d \times d \text{ matrix } \theta.$$

where, if $\alpha = (\alpha_{ij}) \in GL_d(\Lambda)$, then $\alpha^* = (\bar{\alpha}_{ji})$ denotes the conjugate transpose matrix. By Remark A1 the unitary group $U_{2d}^\varepsilon(\Lambda)$ is a subgroup of the *hermitian unitary group* $\text{Isom}(H_\varepsilon(\Lambda^d))$.

Define $D_{2d}(\Lambda)$ to be the subgroup of $GL_{2d}(\Lambda)$ consisting of matrices of the form

$$\begin{pmatrix} Q & 0 \\ 0 & (Q^*)^{-1} \end{pmatrix} \quad \text{where } Q = \begin{pmatrix} a & & \\ & \ddots & \\ & & 1 \end{pmatrix} \in GL_d(\Lambda) \text{ for } a \in \Lambda^\times.$$

This is a subgroup of $U_{2d}^\varepsilon(\Lambda)$ for both choices of $\varepsilon \in \{\pm 1\}$. Next recall that, in Definition 6.3, we defined

$$PB_Q(G, n) = \frac{\text{Diag}(\hat{e})}{[\text{im}(\text{Isom}_\Psi(e^G)) \cdot \text{Tri}(\hat{e})] \cap \text{Diag}(\hat{e})}$$

where $e = e_L$ is the evaluation form and $L = \pi_n(X)$ for X a reference minimal (G, n) -complex. By hypothesis, we have that $L^G \cong \mathbb{Z}^d$ and $\hat{L} \cong H_n(G) \cong (\mathbb{Z}/m)^d$. For the forms e^G and \hat{e} , this implies that $\beta_1 = \cdots = \beta_d = m^{d-1}$ and so $e^G = m^{d-1} \cdot H_\varepsilon(\mathbb{Z}^d)$ and $\hat{e} = m^{d-1} \cdot H_\varepsilon((\mathbb{Z}/m)^d)$ are just scaled ε -hyperbolic forms for $\varepsilon = (-1)^n$ by Proposition 4.15. In particular, there are isometries $e^G \cong H_\varepsilon(\mathbb{Z}^d)$ and $\hat{e} \cong H_\varepsilon((\mathbb{Z}/m)^d)$.

By Remark 3.12, we can choose identifications $L^G \cong \mathbb{Z}^d$ and $\hat{L} \cong (\mathbb{Z}/m)^d$ so that $\psi: L^G \rightarrow \hat{L}$ is reduction mod m . Since $\ker(\psi) \subseteq \mathbb{Z}^d$ is a characteristic subgroup, this implies that

$$\text{Isom}_\Psi(e^G) = \text{Isom}(e^G) \cong \text{Isom}(H_\varepsilon(\mathbb{Z}^d)).$$

In the case where $\Lambda = \mathbb{Z}$ with the trivial involution, we have $\text{Isom}(H_\varepsilon(\mathbb{Z}^d)) = U_{2d}^\varepsilon(\mathbb{Z})$ for n even (Lemma A2). However, note that $\text{Isom}(H_\varepsilon(\mathbb{Z}^d)) \neq U_{2d}^\varepsilon(\mathbb{Z})$ for n odd and any $d \geq 1$.

The remainder of this section will be devoted to establishing the following.

Proposition 7.3. *Let $n \geq 2$, let $\varepsilon = (-1)^n$ and let G be a finite group such that $H_n(G) \cong (\mathbb{Z}/m)^d$ where $m \geq 1$ and $d = r_{(G,n)}$. Then:*

(i) If n is even, there is an isomorphism

$$PB_Q(G, n) \cong \frac{D_{2d}(\mathbb{Z}/m)}{\text{im}(U_{2d}^\varepsilon(\mathbb{Z})) \cap D_{2d}(\mathbb{Z}/m)}.$$

(ii) If n is odd, there is a surjection

$$\frac{D_{2d}(\mathbb{Z}/m)}{\text{im}(U_{2d}^\varepsilon(\mathbb{Z})) \cap D_{2d}(\mathbb{Z}/m)} \twoheadrightarrow PB_Q(G, n).$$

To prove this, first note that the discussion above implies that

$$PB_Q(G, n) \cong \frac{\text{Diag}(\widehat{e})}{[\text{im}(\text{Isom}(H_\varepsilon(\mathbb{Z}^d))) \cdot \text{Tri}(\widehat{e})] \cap \text{Diag}(\widehat{e})}$$

where $\widehat{e} \cong H_\varepsilon((\mathbb{Z}/m)^d)$. It follows by the same argument as given in Proposition 6.4 that $[\text{im}(U_{2d}^\varepsilon(\mathbb{Z})) \cdot \text{Tri}(\widehat{e})] \cap \text{Diag}(\widehat{e})$ is a normal subgroup of $\text{Diag}(\widehat{e})$, where $\text{im}(\cdot)$ denotes the image under the reduction map $U_{2d}^\varepsilon(\mathbb{Z}) \rightarrow U_{2d}^\varepsilon(\mathbb{Z}/m)$. Thus we can define a group

$$\widetilde{PB}_Q(G, n) := \frac{\text{Diag}(\widehat{e})}{[\text{im}(U_{2d}^\varepsilon(\mathbb{Z})) \cdot \text{Tri}(\widehat{e})] \cap \text{Diag}(\widehat{e})}.$$

There is a surjection $\pi : \widetilde{PB}_Q(G, n) \twoheadrightarrow PB_Q(G, n)$ induced by inclusion $U_{2d}^\varepsilon(\mathbb{Z}) \leq \text{Isom}(H_\varepsilon(\mathbb{Z}^d))$. Note that, if n is even, then $U_{2d}^\varepsilon(\mathbb{Z}) = \text{Isom}(H_\varepsilon(\mathbb{Z}^d))$ and so π is an isomorphism.

The key technical step in the proof is the following, which shows that the subgroup of triangular isometries $\text{Tri}(\widehat{e})$ does not contribute to $\widetilde{PB}_Q(G, n)$.

Lemma 7.4. $[\text{im}(U_{2d}^\varepsilon(\mathbb{Z})) \cdot \text{Tri}(\widehat{e})] \cap \text{Diag}(\widehat{e}) = \text{im}(U_{2d}^\varepsilon(\mathbb{Z})) \cap \text{Diag}(\widehat{e})$.

Proof. First note that

$$\begin{aligned} \text{Diag}(\widehat{e}) &= \left\{ \begin{pmatrix} Q & 0 \\ 0 & (Q^*)^{-1} \end{pmatrix} : Q \in GL_d(\mathbb{Z}/m) \right\} \leq GL_{2d}(\mathbb{Z}/m) \\ \text{Tri}(\widehat{e}) &= \left\{ \begin{pmatrix} I & P \\ 0 & I \end{pmatrix} : P \in M_d(\mathbb{Z}/m), P^* = -(-1)^n P \right\} \leq GL_{2d}(\mathbb{Z}/m). \end{aligned}$$

We claim that, if $A \in \text{Diag}(\widehat{e})$ and $B \in \text{Tri}(\widehat{e})$, then $AB \in \text{im}(U_{2d}^\varepsilon(\mathbb{Z}))$ if and only if $A, B \in \text{im}(U_{2d}^\varepsilon(\mathbb{Z}))$. Let $A = \begin{pmatrix} Q & 0 \\ 0 & (Q^*)^{-1} \end{pmatrix}$ and $B = \begin{pmatrix} I & P \\ 0 & I \end{pmatrix}$ where $Q \in GL_d(\mathbb{Z}/m)$ and $P \in M_d(\mathbb{Z}/m)$ is such that $P^* = -(-1)^n P$. If $AB \in \text{im}(U_{2d}^\varepsilon(\mathbb{Z}))$, then $AB \in U_{2d}^\varepsilon(\mathbb{Z}/m)$ and so

$$\begin{pmatrix} Q & QP \\ 0 & (Q^*)^{-1} \end{pmatrix} = AB \in U_{2d}^\varepsilon(\mathbb{Z}/m).$$

It follows from the definition of $U_{2d}^\varepsilon(\mathbb{Z}/m)$ that $Q(QP)^* = E - (-1)^n E^*$ for some $E \in M_d(\mathbb{Z}/m)$ and so $P = F - (-1)^n F^*$ where $F = Q^{-1}E^*(Q^*)^{-1} \in M_d(\mathbb{Z}/m)$. Let $\widetilde{F} \in M_d(\mathbb{Z})$ be any integral lift of F and let $\widetilde{P} = \widetilde{F} - (-1)^n (\widetilde{F})^*$. Then $\begin{pmatrix} I & \widetilde{P} \\ 0 & I \end{pmatrix} \in U_{2d}^\varepsilon(\mathbb{Z})$ and so

$$B = \begin{pmatrix} I & P \\ 0 & I \end{pmatrix} = \psi_* \left[\begin{pmatrix} I & \widetilde{P} \\ 0 & I \end{pmatrix} \right] \in \text{im}(U_{2d}^\varepsilon(\mathbb{Z})).$$

It follows that $A \in \text{im}(U_{2d}^\varepsilon(\mathbb{Z}))$. The converse is clear. This completes the proof of the claim.

Let $N = \text{im}(U_{2d}^\varepsilon(\mathbb{Z})) \cdot \text{Tri}(\widehat{e})$. By the claim, we have $N \cap \text{Diag}(\widehat{e}) = \text{im}(U_{2d}^\varepsilon(\mathbb{Z})) \cap \text{Diag}(\widehat{e})$. In particular, suppose $AB \in N \cap \text{Diag}(\widehat{e})$ for some $A \in \text{im}(U_{2d}^\varepsilon(\mathbb{Z}))$ and $B \in \text{Tri}(\widehat{e})$. Since $AB \in \text{Diag}(\widehat{e})$, $B^{-1} \in \text{Tri}(\widehat{e})$ and $(AB)B^{-1} \in \text{im}(U_{2d}^\varepsilon(\mathbb{Z}))$, the claim implies that $AB \in \text{im}(U_{2d}^\varepsilon(\mathbb{Z}))$ and so $AB \in \text{im}(U_{2d}^\varepsilon(\mathbb{Z})) \cap \text{Diag}(\widehat{e})$. \square

Proof of Proposition 7.3. By Lemma 7.4, we now have that

$$\widetilde{PB}_Q(G, n) \cong \frac{\text{Diag}(\widehat{e})}{\text{im}(U_{2d}^\varepsilon(\mathbb{Z})) \cap \text{Diag}(\widehat{e})}.$$

By Proposition 4.22 and the discussion that followed, we have that

$$\frac{\text{Diag}(\widehat{e})}{\text{Diag}_\Psi(e^G)} \cong (\mathbb{Z}/m)^\times / \{\pm 1\}$$

and we can see directly that $\text{Diag}_\Psi(e^G) = \text{Diag}(H_\varepsilon(\mathbb{Z}^d)) \leq U_{2d}^\varepsilon(\mathbb{Z})$. The subgroup $D_{2d}(\mathbb{Z}/m) \leq \text{Diag}(\widehat{e})$, which is isomorphic to $(\mathbb{Z}/m)^\times$, maps surjectively onto $(\mathbb{Z}/m)^\times / \{\pm 1\}$. This implies

that the inclusion map induces an isomorphism

$$\widetilde{PB}_Q(G, n) \cong \frac{D_{2d}(\mathbb{Z}/m)}{\text{im}(U_{2d}^\varepsilon(\mathbb{Z})) \cap D_{2d}(\mathbb{Z}/m)}.$$

This completes the proof since there is a surjection $\pi: \widetilde{PB}_Q(G, n) \twoheadrightarrow PB_Q(G, n)$ which is an isomorphism provided n is even. \square

7b. Proof of Theorem 7.1 for n even. Let $n \geq 2$ be even. We will now evaluate $PB_Q(G, n)$ using the form established in Proposition 7.3 (i).

Proposition 7.5. *Let $n \geq 2$ be even, let G be a finite group, and suppose that $H_n(G) \cong (\mathbb{Z}/m)^d$ where $m \geq 3$ and $d = r_{(G, n)}$. Then the quotient map $D_{2d}(\mathbb{Z}/m) \rightarrow (\mathbb{Z}/m)^\times / \{\pm 1\}$ induces a homomorphism*

$$\phi: PB_Q(G, n) \cong \frac{D_{2d}(\mathbb{Z}/m)}{\text{im}(U_{2d}(\mathbb{Z})) \cap D_{2d}(\mathbb{Z}/m)} \longrightarrow \frac{(\mathbb{Z}/m)^\times}{\pm(\mathbb{Z}/m)^{\times 2}}.$$

Proof. To check that ϕ is well-defined, we must show that

$$\phi(\text{im}(U_{2d}(\mathbb{Z})) \cap D_{2d}(\mathbb{Z}/m)) \subseteq \{\pm(\mathbb{Z}/m)^{\times 2}\}.$$

Since $m \geq 3$, we have that

$$\text{im}(U_{2d}(\mathbb{Z})) \cap D_{2d}(\mathbb{Z}/m) = \text{im}(SU_{2d}(\mathbb{Z})) \cap D_{2d}(\mathbb{Z}/m).$$

To see this note that, if $A \in GL_{2d}(\mathbb{Z})$ and the reduction $\bar{A} \in GL_{2d}(\mathbb{Z}/m)$ has $\det(\bar{A}) = 1$, then $\det(A) \in \{\pm 1\}$ and $\det(A) \equiv \det(\bar{A}) \equiv 1 \pmod{m}$ which implies that $\det(A) = 1$ since $m \geq 3$.

Let $\rho_m: SU_{2d}(\mathbb{Z}) \rightarrow SU_{2d}(\mathbb{Z}/m)$ denote reduction mod m . Then we have a commutative diagram

$$\begin{array}{ccccc} \rho_m^{-1}(D_{2d}(\mathbb{Z}/m)) & \hookrightarrow & SU_{2d}(\mathbb{Z}) & \longrightarrow & SU(\mathbb{Z})/RU(\mathbb{Z}) = L_1^s(\mathbb{Z}) \\ \downarrow \rho_m & & \downarrow \rho_m & & \downarrow \rho_m \\ D_{2d}(\mathbb{Z}/m) & \hookrightarrow & SU_{2d}(\mathbb{Z}/m) & \longrightarrow & SU(\mathbb{Z}/m)/RU(\mathbb{Z}/m) = L_1^s(\mathbb{Z}/m). \end{array}$$

The image of the subgroup

$$N := \text{im}(U_{2d}(\mathbb{Z})) \cap D_{2d}(\mathbb{Z}/m) \leq \text{im}(SU_{2d}(\mathbb{Z}) \rightarrow SU_{2d}(\mathbb{Z}/m))$$

after stabilisation gives a subgroup

$$\bar{N} \subseteq \text{im}(L_1^s(\mathbb{Z}) \rightarrow L_1^s(\mathbb{Z}/m)).$$

The calculations of Wall [68, §1] show that $L_1^s(\mathbb{Z}) = L_1^h(\mathbb{Z}) = 0$, so that

$$0 = \bar{N} \subseteq \ker(L_1^s(\mathbb{Z}/m) \rightarrow L_1^h(\mathbb{Z}/m)) \cong (\mathbb{Z}/m)^\times / \pm(\mathbb{Z}/m)^{\times 2}$$

by naturality and Proposition B1 (based on the calculations of Lemmas B2 and B3). Moreover, the composite under stabilisation

$$D_{2d}(\mathbb{Z}/m) \rightarrow SU_{2d}(\mathbb{Z}/m) \rightarrow L_1^s(\mathbb{Z}/m) \cong (\mathbb{Z}/m)^\times / \pm(\mathbb{Z}/m)^{\times 2}$$

is just the quotient map used to define ϕ (see Corollary B5). Therefore $\phi(\text{im}(U_{2d}(\mathbb{Z})) \cap D_{2d}(\mathbb{Z}/m)) \subseteq \{\pm(\mathbb{Z}/m)^{\times 2}\}$, and ϕ is well-defined. \square

We now claim that the map ϕ is an isomorphism provided that, in addition, we have $d \geq 3$. We will begin by establishing the following lifting result for squares, where $(D_{2r}(\mathbb{Z}/m))^2$ denotes the subgroup $\{a^2 : a \in D_{2d}(\mathbb{Z}/m)\}$.

Proposition 7.6. *Let $m \geq 1$, $d \geq 3$, and let $\varepsilon \in \{\pm 1\}$. Then*

$$(D_{2d}(\mathbb{Z}/m))^2 \subseteq \text{im}(EU_{2d}^\varepsilon(\mathbb{Z})) \subseteq \text{im}(U_{2d}^\varepsilon(\mathbb{Z}))$$

where $\text{im}(\cdot)$ denotes the image under the mod m reduction map $U_{2d}^\varepsilon(\mathbb{Z}) \rightarrow U_{2d}^\varepsilon(\mathbb{Z}/m)$.

The proof will be based on the following well-known identities. For $r \geq 1$, let I_r denote the $n \times n$ identity matrix and \oplus denote the orthogonal block sum of matrices.

Lemma 7.7. *Let Λ be a ring and set $I = I_r$ for some $r \geq 1$.*

(i) *If $A, B \in GL_r(\Lambda)$, then*

$$A^{-1}B^{-1}AB \oplus I = ((BA)^{-1} \oplus BA)(A \oplus A^{-1})(B \oplus B^{-1}) \in GL_{2r}(\Lambda).$$

(ii) *If $A \in GL_r(\Lambda)$, then*

$$A \oplus A^{-1} = \begin{pmatrix} I & A \\ 0 & I \end{pmatrix} \begin{pmatrix} I & 0 \\ I-A^{-1} & I \end{pmatrix} \begin{pmatrix} I & -I \\ 0 & I \end{pmatrix} \begin{pmatrix} I & 0 \\ I-A & I \end{pmatrix} \in GL_{2r}(\Lambda).$$

The two identities can be verified directly by multiplication. The former appears in [67, Proof of Theorem 6.3] and the latter is the Whitehead identity.

Lemma 7.8. *Let Λ be a ring with involution. If $Q \in GL_r(\Lambda)$, then $Q \oplus Q^{-1} \oplus I_r \in EU_{3r}^\varepsilon(\Lambda)$.*

Proof. By matrix multiplication, we have

$$Q \oplus Q^{-1} \oplus I_r = (Q \oplus I_r \oplus (Q^*)^{-1}) (I_r \oplus Q \oplus (Q^*)^{-1})^{-1} \in U_{3r}^\varepsilon(\Lambda)$$

and, since $Q \in GL_r(\Lambda)$, we have $Q \oplus (Q^*)^{-1} \in EU_{2r}^\varepsilon(\Lambda)$ by definition of the elementary unitary relations. The result follows by stabilisation. \square

Proof of Proposition 7.6. It suffices to treat the case $d = 3$ since, after reordering bases, an arbitrary element of $(D_{2d}(\mathbb{Z}/m))^2$ has the form $\begin{pmatrix} a^2 & 0 \\ 0 & a^{-2} \end{pmatrix} \oplus I_{2d-2}$ for some $a \in (\mathbb{Z}/m)^\times$.

We start by applying Lemma 7.7 (i) to the matrices

$$A = \begin{pmatrix} a^{-1} & 0 \\ 0 & a \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad (BA)^{-1} = \begin{pmatrix} 0 & a \\ a^{-1} & 0 \end{pmatrix}$$

in $GL_2(\mathbb{Z}/m)$. Since $A^{-1}B^{-1}AB = \begin{pmatrix} a^2 & 0 \\ 0 & a^{-2} \end{pmatrix}$, Lemma 7.7 (i) with $r = 2$ gives that

$$\begin{pmatrix} a^2 & 0 \\ 0 & a^{-2} \end{pmatrix} \oplus I_2 = A^{-1}B^{-1}AB \oplus I_2 = ((BA)^{-1} \oplus BA)(A \oplus A^{-1})(B \oplus B^{-1}) \in GL_4(\mathbb{Z}/m).$$

It remains to show that the three matrices on the right-hand side are, after one stabilisation, in the image of $EU_6^\varepsilon(\mathbb{Z})$ under the map induced by reduction mod m .

First, it is clear that B lifts to $\tilde{B} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \in GL_2(\mathbb{Z})$. Since B is symmetric and \mathbb{Z} has trivial involution, $B \oplus B^{-1} = B \oplus (B^*)^{-1}$ lifts to $\tilde{B} \oplus (\tilde{B}^*)^{-1} \in EU_4^\varepsilon(\mathbb{Z})$. Secondly, by Lemma 7.7 (ii), we have that

$$A = \begin{pmatrix} a^{-1} & 0 \\ 0 & a \end{pmatrix} = \begin{pmatrix} 1 & a^{-1} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1-a & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1-a^{-1} & 1 \end{pmatrix}$$

and each of the matrices on the right hand side lifts to $GL_2(\mathbb{Z})$. Hence A lifts to $\tilde{A} \in GL_2(\mathbb{Z})$. Since A is symmetric and \mathbb{Z}/m has trivial involution, $A \oplus A^{-1} = A \oplus (A^*)^{-1}$ lifts to $\tilde{A} \oplus ((\tilde{A})^*)^{-1} \in EU_4(\mathbb{Z})$. Finally, BA lifts to $\tilde{B}\tilde{A} \in GL_2(\mathbb{Z})$ and so $(BA)^{-1} \oplus BA \oplus I_2$ lifts to $(\tilde{B}\tilde{A})^{-1} \oplus (\tilde{B}\tilde{A}) \oplus I_2$. By Lemma 7.8, this is contained in $EU_6^\varepsilon(\mathbb{Z}/m)$. This completes the proof. \square

Proof of Theorem 7.1 (i). By combining Proposition 7.3 (i) and Proposition 7.5, we have a map

$$PB_Q(G, n) \cong \frac{D_{2d}(\mathbb{Z}/m)}{\text{im}(U_{2d}^\varepsilon(\mathbb{Z}) \cap D_{2d}(\mathbb{Z}/m))} \xrightarrow{\phi} \frac{(\mathbb{Z}/m)^\times}{\pm(\mathbb{Z}/m)^{\times 2}}.$$

We claim that ϕ is bijective (it is clearly surjective by definition). The result holds for $m \leq 2$ since both sides are trivial. For example, if $m = 2$, then we have $D_{2d}(\mathbb{Z}/2) \cong (\mathbb{Z}/2)^\times = \{1\}$. From now on, we will assume that $m \geq 3$.

To see that ϕ is injective, suppose that $A = (a) \oplus (a^{-1}) \oplus I_{2d-2} \in D_{2d}(\mathbb{Z}/m)$ has $\phi(A) = 0$. This implies that $a \in \pm(\mathbb{Z}/m)^{\times 2}$ and so $A \in \pm(D_{2d}(\mathbb{Z}/m))^{\times 2}$. By Proposition 4.22, we have that $-I_{2d} \in \rho_m^{-1}(D_{2d}(\mathbb{Z}/m))$ where $\rho_m: SU_{2d}(\mathbb{Z}) \rightarrow SU_{2d}(\mathbb{Z}/m)$ denote reduction mod m . By Proposition 7.6, we have that $(D_{2d}(\mathbb{Z}/m))^{\times 2} \subseteq \rho_m^{-1}(D_{2d}(\mathbb{Z}/m))$. Hence the image $[A] = 0 \in PB_Q(G, n)$, and ϕ is injective. \square

7c. Proof of Theorem 7.1 for n odd. Let $n > 2$ be odd so that $\varepsilon = -1$. In order to show that $PB_Q(G, n) = 0$, it is enough to check that every element in $D_{2d}(\mathbb{Z}/m) \subseteq SU_{2d}^\varepsilon(\mathbb{Z}/m)$ can be lifted to $U_{2d}^\varepsilon(\mathbb{Z})$ (see Proposition 7.3 (ii)). From Proposition B4, we have $L_3^s(\mathbb{Z}/m) = SU^\varepsilon(\mathbb{Z}/m)/RU^\varepsilon(\mathbb{Z}/m) = 0$, and hence the image of $D_{2d}(\mathbb{Z}/m)$ after stabilisation is zero in $L_3^s(\mathbb{Z}/m)$. In other words, $D_{2d}(\mathbb{Z}/m) \subseteq RU^\varepsilon(\mathbb{Z}/m)$.

By stability results for unitary groups (see [4], [40, Chapter VI]), and the fact that the ring $\Lambda = \mathbb{Z}/m$ has stable range 1, it follows that $RU_{2d}^\varepsilon(\mathbb{Z}/m) = RU^\varepsilon(\mathbb{Z}/m)$ provided that $d \geq 3$. Hence any element in $RU_{2d}^\varepsilon(\mathbb{Z}/m)$ can be expressed as a product of elementary special unitary matrices (see the list in Appendix A, with the additional assumption $\det Q = 1$ in item (i)). We apply this observation to each element $A = \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \in D_{2d}(\mathbb{Z}/m)$. Each of the terms in this product can be lifted to $U_{2d}^\varepsilon(\mathbb{Z})$, by the methods used in the proof of Proposition 7.6, and hence $PB_Q(G, n) = 0$ for n odd.

8. EXAMPLES OF HOMOTOPY INEQUIVALENT (G, n) -COMPLEXES

The aim of this section will be to survey examples of finite (G, n) -complexes X and Y with $\pi_1(X) \cong \pi_1(Y)$ and $\chi(X) = \chi(Y)$ but which are not homotopy equivalent. The 4-manifolds which we construct in Section 9 in order to prove Theorem C will all be constructed by applying the doubling construction to the examples here.

Let β denote the bias invariant defined in Section 3. For a finite abelian group G and $r \in (\mathbb{Z}/m_{(G,n)})^\times$, define $X_{G,n}^r$ to be the finite (G, n) -complexes defined in [57, Proof of Proposition 6]. For example, when $n = 2$ and $G = \mathbb{Z}/m_1 \times \cdots \times \mathbb{Z}/m_d$ with $m_i \mid m_{i+1}$ for all i and $m_1 > 1$ (and so $m_1 = m_G$), we have that $X_{G,2}^r$ coincides with the presentation complex for the presentation:

$$\mathcal{P}_r = \langle x_1, \dots, x_d \mid x_1^{m_1}, \dots, x_d^{m_d}, [x_1^r, x_2], \{[x_i, x_j] : i < j, (i, j) \neq (1, 2)\} \rangle.$$

We will now point out the following. This is a consequence of work of Metzler [47], Sieradski [58], Sieradski-Dyer [57], Browning [9, 10] and Linnell [46]. However, as far as we know, this observation has not previously appeared in the literature for all $n \geq 2$.

Theorem 8.1. *Let $n \geq 2$, let G be a finite abelian group and let $m = m_{(G,n)}$. Then:*

- (i) *The bias invariant gives a bijection*

$$\beta : \text{HT}_{\min}(G, n) \rightarrow B(G, n).$$

Furthermore, $\beta(X_{G,n}^r) = [r]$ for all $r \in (\mathbb{Z}/m)^\times$ and so every minimal finite (G, n) -complex X has $X \simeq X_{G,n}^r$ for some $r \in (\mathbb{Z}/m)^\times$.

- (ii) *If X, Y are finite (G, n) -complexes with $(-1)^n \chi(X) = (-1)^n \chi(Y) > \chi_{\min}(G, n)$, then $X \simeq Y$.*

This is achieved using the Browning invariant, as introduced by Browning in [9]. For a finite group G and finite (G, n) -complexes X and Y , this is an invariant $B(X, Y) \in \text{Br}(G, n)$ where $\text{Br}(G, n)$ is the Browning obstruction group (see also [43, Section 2]). If G is a finite abelian group, then this is a complete homotopy invariant.

Proof. (i) We begin by noting that, if X is a minimal finite (G, n) -complex, then $X \simeq X_{G,n}^r$ for some $r \in (\mathbb{Z}/m)^\times$. The case $n = 2$ is [9, Theorem 1.7]. For the general case, the result follows from the fact that, as noted by Linnell in [46, p318], the complexes $X_{G,n}^r$ realise all the elements of the Browning obstruction group $\text{Br}(G, n)$.

Note that β is surjective since $\beta(X_{G,n}^r) = [r]$ for all $r \in (\mathbb{Z}/m)^\times$ [57, Proposition 6]. To show that β is injective, it remains to prove that the complexes $X_{G,n}^r$ are determined up to homotopy equivalence by β . The case $n = 2$ was proven by Sieradski [58] by constructing explicit homotopy equivalences. The general case is a consequence of the comparison between [57, Proposition 8], which gives a lower bound on $|\text{HT}_{\min}(G, n)|$ by computing $\text{im}(\beta)$, and [46, Theorem 1.3] which shows that $|\text{HT}_{\min}(G, n)|$ is equal to this lower bound (see also the discussion on [46, p307]).

- (ii) This follows from a result of Browning [10, Theorem 5.4] (see also [46, Theorem 1.1]). \square

Define $\gamma(G, n) = |\text{HT}_{\min}(G, n)|$. The above shows that, for $n \geq 2$ and G finite abelian, we have $\gamma(G, n) = |B(G, n)|$. Recall from Section 3c that $B(G, n) = (\mathbb{Z}/m)^\times / \pm D(G, n)$ where $D(G, n) = \text{im}(\varphi_{(G,n)} : \text{Aut}(G) \rightarrow (\mathbb{Z}/m)^\times / \{\pm 1\})$.

The following was shown by Browning [10] in the case $n = 2$, and Sieradski-Dyer [57, Proposition 8] and Linnell [46, Theorem 1.3 & Corollary 1.5] in the general case.

Proposition 8.2. *Let $n \geq 2$, let G be a finite abelian group, let $m = m_{(G,n)}$ and $d = d(G)$. Then:*

$$D(G, n) = ((\mathbb{Z}/m)^\times)^{e(d,n)}$$

where

$$e(d, n) = \begin{cases} \sum_{i=0}^{\frac{n}{2}-1} \frac{n-2i}{2} \binom{d+2i-1}{d-2}, & \text{if } n \text{ is even and } d \geq 2 \\ \sum_{i=0}^{\frac{n-1}{2}} \frac{n+1-2i}{2} \binom{d+2i-2}{d-2}, & \text{if } n \text{ is odd and } d \geq 2 \end{cases}$$

and $e(1, n) = 1$ for n even, $e(1, n) = \frac{1}{2}(n+1)$ for n odd.

In particular, we have $\gamma(G, n) = |(\mathbb{Z}/m)^\times / \pm ((\mathbb{Z}/m)^\times)^{e(d,n)}|$.

Remark 8.3. In [57, Proposition 8], Sieradski-Dyer proved that the number of minimal finite (G, n) -complexes up to homotopy equivalence was at least the bound given above. However, as pointed out in [46, p305], the formula for $e(d, n)$ given in [57, p210-211] is incorrect when $n \geq 3$.

For example, $e(d, 2) = d - 1$ for $d \geq 2$. Using elementary number theory, it is possible to evaluate $\gamma(G, n)$ precisely (see [58, p137]).

We will now consider the homotopy classification of finite (G, n) -complexes X for which $\pi_n(X)$ is a fixed $\mathbb{Z}G$ -module. For a minimal finite (G, n) -complex X , define

$$\gamma'(G, n) = |\{Y \in \text{HT}_{\min}(G, n) : \pi_n(X) \cong_{\text{Aut}(G)} \pi_n(Y)\}|.$$

Let $N = \sum_{g \in G} g \in \mathbb{Z}G$ denote the group norm and let $(N, r) = \mathbb{Z}G \cdot N + \mathbb{Z}G \cdot r \leq \mathbb{Z}G$ denote the Swan module corresponding to $r \in \mathbb{Z}$. In what follows, we write $[\cdot]$ to denote the quotient map $(\mathbb{Z}/m)^\times / \{\pm 1\} \twoheadrightarrow (\mathbb{Z}/m)^\times / \pm ((\mathbb{Z}/m)^\times)^{e(d,n)} \cong B(G, n)$.

Theorem 8.4. *Let $n \geq 2$, let p be a prime, let G be a non-cyclic abelian p -group, let $d = d(G)$ and $m = m_{(G,n)}$. Then:*

(i) *For a minimal finite (G, n) -complex X , we have*

$$\beta(\{Y \in \text{HT}_{\min}(G, n) : \pi_n(X) \cong_{\text{Aut}(G)} \pi_n(Y)\}) = \beta(X) \cdot [V^{s(d,n)}] \subseteq B(G, n)$$

where $s(d, n) = \sum_{i=0}^{n-1} (-1)^{n+i+1} \binom{d+i}{d-1}$ and

$$V = \{[r] \in (\mathbb{Z}/m)^\times : r \in \mathbb{Z}, (r, |G|) = 1 \text{ and } (N, r) \cong \mathbb{Z}G\}.$$

(ii) *$\gamma'(G, n)$ is equal to the number of elements in $[V^{s(d,n)}] \subseteq B(G, n)$. In particular, it depends only on G and n , and not on the choice of X .*

(iii) *If $p = 2$, then $\gamma'(G, n) = 1$. If p is odd, then*

$$\gamma'(G, n) = \frac{\gcd(e(d, n), \frac{1}{2}(p-1))}{\gcd(e(d, n), s(d, n), \frac{1}{2}(p-1))}.$$

Proof. This is proven by combining Theorem 8.1 with results of Linnell [46]. More specifically, (i) follows from [46, Theorem 8.4 (iii)], (ii) follows from (i), and (iii) follows from [46, Theorems 1.2 (4)]. \square

We will now use this to show the following.

Theorem 8.5. *Let $n \geq 2$. Then there exists $d \geq 3$ such that, if $p \equiv 1 \pmod{4}$ is prime and $G = C_{p^{m_1}} \times \cdots \times C_{p^{m_d}}$ with $m_i \leq m_{i+1}$ for all i and $m_1 \geq 1$, then $\gamma'(G, n) > 1$. In particular, there exist finite (G, n) -complexes X and Y such that $\pi_n(X) \cong \pi_n(Y)$ as $\mathbb{Z}G$ -modules but $X \not\cong Y$.*

This will be a consequence of combining Theorem 8.4 with the following result concerning the parity of $e(d, n)$ and $s(d, n)$. For the remainder of this section, let \equiv denote equivalence mod 2.

Proposition 8.6. *Let $n \geq 2$. Then there exists $d \geq 3$ such that $e(d, n) \equiv 0$ and $s(d, n) \equiv 1$.*

The proof is based on detailed calculations presented in Appendix C.

Proof. We refer to Appendix C for the relevant properties of the functions $e(d, n)$ and $s(d, n)$. Here is a summary of the results:

- If $n = 4k$, then $e(4t, 4k) \equiv 0$ for any $t \geq 1$ by Proposition C3. By Proposition C4, we have $s(4t, 4k) \equiv 1 + \binom{k+t}{t}$. Let $k = 2^m r$ where $m \geq 0$ and $r \geq 1$ is odd. If we take $t = 2^m$, then $s(2^{m+2}, 4k) \equiv 1 + \binom{2^m r + 2^m}{2^m} \equiv 1 + \binom{r+1}{1} \equiv r \equiv 1$ using Lemma C1.
- If $n = 4k + 1$, then $s(4t + 1, 4k + 1) \equiv 1$ for any $t \geq 1$ by Proposition C4. By Proposition C3, we have $e(4t + 1, 4k + 1) \equiv \binom{k+t}{t}$. Similarly to the $n = 4k$ case, let $k = 2^m r$ where $m \geq 0$ and $r \geq 1$ is odd. If we take $t = 2^m$, then $e(2^{m+2} + 1, 4k + 1) \equiv \binom{2^m r + 2^m}{2^m} \equiv r + 1 \equiv 0$.
- If $n = 4k + 2$, then $e(d, 4k + 2) \equiv 0$ and $s(d, 4k + 2) \equiv 1$ whenever $d \equiv 3 \pmod{4}$, by Propositions C3 and C4. For example, we can take $d = 3$.
- If $n = 4k + 3$, then $e(d, 4k + 3) \equiv 0$ and $s(d, 4k + 3) \equiv 0$ whenever $d \equiv 1 \pmod{4}$, by Propositions C3 and C4. For example, we can take $d = 5$. \square

Proof of Theorem 8.5. By Proposition 8.6, there exists $d \geq 3$ such that $e(d, n) \equiv 0$ and $s(d, n) \equiv 1$. Let p be a prime such that $p \equiv 1 \pmod{4}$ and $G = C_{p^{m_1}} \times \cdots \times C_{p^{m_d}}$ with $m_i \leq m_{i+1}$ for all i and $m_1 \geq 1$. By Theorem 8.4, we have that

$$\gamma'(G, n) = \frac{\gcd(e(d, n), \frac{1}{2}(p-1))}{\gcd(e(d, n), s(d, n), \frac{1}{2}(p-1))}.$$

Since $e(d, n) \equiv 0$, and $p \equiv 1 \pmod{4}$ implies that $\frac{1}{2}(p-1) \equiv 0$, we have $2 \mid \gcd(e(d, n), \frac{1}{2}(p-1))$. Conversely, since $s(d, n) \equiv 1$ and $2 \nmid \gcd(e(d, n), s(d, n), \frac{1}{2}(p-1))$, we have $2 \mid \gamma'(G, n)$. \square

9. EXAMPLES OF HOMOTOPY INEQUIVALENT DOUBLED (G, n) -COMPLEXES

The aim of this section will be to use the quadratic bias invariant to distinguish certain doubled (G, n) -complexes $M(X)$ up to homotopy equivalence.

The examples must take as input a finite group G and a pair of finite (G, n) -complexes X, Y with $\chi(X) = \chi(Y)$ but which are not homotopy equivalent. Such examples have previously only been known to exist when G is either a finite abelian group [47, 57] or a group with periodic cohomology [18, 50, 52]. In light of Theorem 7.1 (ii), we must restrict to the case where $n \geq 2$ is even. If G has periodic cohomology, then $H_n(G) = 0$ for all $n \geq 2$ even [61, Corollary 2] and so the quadratic bias invariant contains no information.

For this reason, we will begin by restricting to examples over finite abelian fundamental groups (Section 9a). In Section 9b, we will demonstrate that the quadratic bias invariant is computable more generally by constructing examples over the non-abelian group $Q_8 \times (\mathbb{Z}/17)^3$. This also gives the first example of a non-abelian finite group G which does not have periodic cohomology such that there exists homotopically distinct finite (G, n) -complexes X, Y with $\chi(X) = \chi(Y)$.

9a. Examples with abelian fundamental groups. The aim of this section will be to establish the following two results, which imply Theorem C and Theorem 1.6 from the Introduction.

Theorem 9.1. *Let $n \geq 2$ be even and let $k \geq 2$. Then there exist closed smooth $2n$ -manifolds M_1, M_2, \dots, M_k which are all stably diffeomorphic but not pairwise homotopy equivalent.*

Furthermore, if m is an integer with at least $1 + \log_2(k)$ prime factors, then the examples can be taken to have fundamental group $(\mathbb{Z}/m)^d$ for some $d \geq 3$ or $(\mathbb{Z}/m)^d \times \mathbb{Z}/t$ for some $d \geq 4$ and any integer $t > 1$ such that $t \mid m$ and $m, m/t$ have the same prime factors.

Theorem 9.2. *Let $n \geq 2$ be even. Then there exist closed smooth $2n$ -manifolds M and N which are stably diffeomorphic, have isometric hyperbolic equivariant intersection forms, but are not homotopy equivalent. Furthermore, for any prime p such that $p \equiv 1 \pmod{4}$, the examples can be taken to have fundamental group $(\mathbb{Z}/p)^d$ for some $d \geq 3$.*

The proofs will rely on Theorem 7.1 which computes the quadratic bias obstruction group $B_Q(G, n)$ in the case where G is a finite group such that $H_n(G) \cong (\mathbb{Z}/m)^r$ where $r = r_{(G, n)}$. We start by giving a collection of finite abelian groups which satisfy this hypothesis.

Lemma 9.3. *Let $n \geq 2$ and let $G = (\mathbb{Z}/m)^d \times \mathbb{Z}/t$ where $d \geq 3$ and $m, t \geq 1$ are such that $t \mid m$ and $m, m/t$ have the same prime factors. Then*

$$H_n(G) \cong \begin{cases} (\mathbb{Z}/m)^r, & \text{if } n \text{ is even} \\ (\mathbb{Z}/m)^r \times \mathbb{Z}/t, & \text{if } n \text{ is odd} \end{cases}$$

where $r = r_{(G,n)} \geq 3$.

In particular, the groups $G = (\mathbb{Z}/m)^d$ for $d \geq 3$ satisfy the hypothesis for all $n \geq 2$. If $n \geq 2$ is even, then the hypothesis is satisfied by the larger class of groups $G = (\mathbb{Z}/m)^d \times \mathbb{Z}/t$ where $d \geq 3$, $t \mid m$ and $m, m/t$ have the same prime factors.

Proof. Let $G = \prod_{i=1}^s G_i$ where G_i is the p_i -primary component for some prime p_i . We have $H_n(G) \cong \prod_{i=1}^s H_n(G_i)$. The hypothesis on m and t imply that $G_i \cong (\mathbb{Z}/p_i^{a_i})^d \times \mathbb{Z}/p_i^{b_i}$ where $a_i > b_i \geq 1$. By [17, Theorem 4.3], we have that

$$H_n(G_i) \cong (\mathbb{Z}/p_i^{a_i})^r \times (\mathbb{Z}/p_i^{b_i})^{\nu(n,1) - (-1)^n}$$

where $r = \sum_{k=2}^d (\nu(n,k) - (-1)^n)$ and $\nu(n,k) = \sum_{j=0}^n (-1)^{n+j} \binom{k+j-1}{j}$. We can verify directly that, if $d \geq 3$, then $r \geq 3$. Furthermore, $\nu(n,1) = 1$ if n is even and $\nu(n,1) = 0$ if n is odd. This gives the required form for $H_n(G)$ by recombining the p_i -primary components.

Finally, the fact that $r = r_{(G,n)}$ follows since G satisfies the strong minimality hypothesis (see Section 3a and [57, Proposition 5]). \square

Fix $n \geq 2$ even, $m \geq 2$, $d \geq 3$ and $t \mid m$ such that $m, m/t$ have the same prime factors. Let $G = (\mathbb{Z}/m)^d \times \mathbb{Z}/t$. By Theorem 7.1 and Lemma 9.3, there is an isomorphism

$$B_Q(G, n) \cong \frac{(\mathbb{Z}/m)^\times}{\pm(\mathbb{Z}/m)^{\times 2} \cdot D(G, n)}.$$

Let $\delta(t) = 1$ if $t > 1$ and $\delta(t) = 0$ if $t = 1$. We will write $\delta = \delta(t)$, so that $d(G) = d + \delta$.

Since G is abelian, Proposition 8.2 implies that $D(G, n) = ((\mathbb{Z}/m)^\times)^{e(d+\delta, n)}$. It follows that

$$B_Q(G, n) \cong \begin{cases} \frac{(\mathbb{Z}/m)^\times}{\pm(\mathbb{Z}/m)^{\times 2}}, & \text{if } e(d+\delta, n) \text{ is even} \\ 0, & \text{if } e(d+\delta, n) \text{ is odd.} \end{cases}$$

By Theorem 8.1, we also have that $\beta: \text{HT}_{\min}(G, n) \rightarrow B(G, n)$ is a bijection. By Proposition 6.14, this implies that $\beta_Q: \mathcal{M}_{2n}(G) \rightarrow B_Q(G, n)$ is surjective.

Proof of Theorem 9.1. By Proposition 5.1, all the manifolds in $\mathcal{M}_{2n}(G)$ are stably diffeomorphic. It therefore suffices to show that, for all $k \geq 2$, there exists $m \geq 2$, $t \geq 1$ and $d \geq 3$ such that $|\mathcal{M}_{2n}(G)| \geq k$ when $G = (\mathbb{Z}/m)^d \times \mathbb{Z}/t$.

By Proposition 8.6, there exists $s \geq 3$ such that $e(s, n)$ is even. We now choose either $d = s$ and $t = 1$, or $d = s - 1$ and any $t > 1$ such that $t \mid m$ and $m, m/t$ have the same prime factors. By the above discussion, this implies that

$$|\mathcal{M}_{2n}(G)| \geq \left| \frac{(\mathbb{Z}/m)^\times}{\pm(\mathbb{Z}/m)^{\times 2}} \right|.$$

Let $r \geq 1$. Then $|(\mathbb{Z}/2^r)^\times / ((\mathbb{Z}/2^r)^\times)^2| = 4$ and, if p is an odd prime, then we have $|(\mathbb{Z}/p^r)^\times / ((\mathbb{Z}/p^r)^\times)^2| = 2$. If $m = p_1^{\alpha_1} \cdots p_t^{\alpha_t}$ for distinct primes p_1, \dots, p_t and $\alpha_i \geq 1$, then

$$|\mathcal{M}_{2n}(G)| \geq \frac{1}{2} \left| \frac{(\mathbb{Z}/m)^\times}{(\mathbb{Z}/m)^{\times 2}} \right| = \frac{1}{2} \prod_{i=1}^t \left| \frac{(\mathbb{Z}/p_i^{\alpha_i})^\times}{((\mathbb{Z}/p_i^{\alpha_i})^\times)^2} \right| \geq 2^{t-1}.$$

Thus, if m has at least $1 + \log_2(k)$ distinct prime factors, then $|\mathcal{M}_{2n}(G)| \geq k$. \square

In order to prove Theorem 9.2, we will need the following which is a slight extension of the examples constructed in Theorem 8.5.

Lemma 9.4. *Let $n \geq 2$, let $d \geq 3$, let p be an odd prime and let $G = (\mathbb{Z}/p)^d$. Let X be a reference minimal (G, n) -complex. Then:*

$$\beta(\{Y \in \text{HT}_{\min}(G, n) : \pi_n(X) \cong_{\text{Aut}(G)} \pi_n(Y)\}) = \frac{\pm((\mathbb{Z}/p)^\times)^{\gcd(e(d,n), s(d,n))}}{\pm((\mathbb{Z}/p)^\times)^{e(d,n)}}$$

Proof. By Theorem 8.4, we have that the image is $\pm V^{s(d,n)} \cdot ((\mathbb{Z}/m)^\times)^{e(d,n)} / \pm ((\mathbb{Z}/m)^\times)^{e(d,n)}$ where $V = \{[r] \in (\mathbb{Z}/p)^\times : r \in \mathbb{Z}, (r, |G|) = 1 \text{ and } (N, r) \cong \mathbb{Z}G\}$. It is well known that (N, r) is a projective $\mathbb{Z}G$ -module. Since G is abelian and $\mathbb{Z}G$ has projective cancellation (see [46, Lemma 3.2]), we have that $V = \ker((\mathbb{Z}/p)^\times \rightarrow \widetilde{K}_0(\mathbb{Z}G))$ is the kernel of the Swan map. It is now a consequence of a result of Taylor [62, Theorem 3] (see also [14, 54.15]) that, since G is a non-cyclic p -group for p odd, we have $|V| = \frac{1}{2}(p-1)$ (see [46, Theorem 2.1]). Since $V \leq (\mathbb{Z}/p)^\times$, we therefore have $V = (\mathbb{Z}/p)^\times$. The result follows. \square

Proof of Theorem 9.2. By Proposition 8.6, there exists $d \geq 3$ be such that $e(d, n)$ is even and $s(d, n)$ is odd. Let p be a prime such that $p \equiv 1 \pmod{4}$ and let $G = (\mathbb{Z}/p)^d$. Since $p \equiv 1 \pmod{4}$, -1 is a square mod p and so $\pm(\mathbb{Z}/p)^\times = (\mathbb{Z}/p)^\times$. Since $e(d, n)$ is even, we therefore have that $B_Q(G, n) \cong (\mathbb{Z}/p)^\times / (\mathbb{Z}/p)^{\times 2}$ which has order two. Let $q: B(G, n) \rightarrow B_Q(G, n)$ denote the natural quotient map defined in Proposition 6.14.

By Lemma 9.4, we now have that

$$q(\beta(\{Y \in \text{HT}_{\min}(G, n) : \pi_n(X) \cong_{\text{Aut}(G)} \pi_n(Y)\})) = \frac{((\mathbb{Z}/p)^\times)^{\gcd(e(d,n), s(d,n))}}{(\mathbb{Z}/p)^{\times 2}} = \frac{(\mathbb{Z}/p)^\times}{(\mathbb{Z}/p)^{\times 2}}$$

since the fact that $s(d, n)$ is odd implies that $\gcd(e(d, n), s(d, n)) \equiv 1 \pmod{2}$. By Proposition 6.14, we have that $q \circ \beta = \beta_Q \circ \mathcal{D}$. Hence there exist $X, Y \in \text{HT}_{\min}(G, n)$ such that $\pi_n(X) \cong_{\text{Aut}(G)} \pi_n(Y)$ and $\beta_Q(M(X)) \neq \beta_Q(M(Y))$, which implies that $M(X) \not\cong M(Y)$. By Proposition 5.1, $M(X)$ and $M(Y)$ are stably diffeomorphic.

It remains to show that $M(X)$ and $M(Y)$ have isometric equivariant intersection forms. For simplicity, we first rechoose the identification $\pi_1(Y) \cong G$ so that $L := \pi_n(X) \cong \pi_n(Y)$ are isomorphic as $\mathbb{Z}G$ -modules. By Proposition 5.2 (proven in [42, Proposition II.2]), we have that there are isometries $S_{M(X)} \cong \text{Met}(L^* \oplus L, \phi_X)$ and $S_{M(Y)} \cong \text{Met}(L^* \oplus L, \phi_Y)$ for some $\phi_X, \phi_Y \in \text{Sym}(L)$ with $\phi_X^G = \phi_Y^G = 0$. Since $M(X)$ is stably parallelisable, $\widetilde{M(X)}$ is spin and so Example 4.5 implies that $S_{M(X)}$ is weakly even. Since $|G| = p^d$ is odd, weakly even metabolic forms over $\mathbb{Z}G$ are hyperbolic (see Proposition 4.6). Hence $S_{M(X)} \cong S_{M(Y)} \cong H(L)$. \square

9b. Examples with non-abelian fundamental groups. We will now establish the following. For simplicity, we will restrict to the case of 4-manifolds and to a small range of fundamental groups. A similar result can be obtained for manifolds of arbitrary dimension $2n \geq 4$, and for a much wider range of fundamental groups.

Theorem 9.5. *Let $G = Q_8 \times (\mathbb{Z}/p)^3$ where p is a prime such that $p \equiv 1 \pmod{8}$. Then:*

- (i) *There exist minimal finite 2-complexes X, Y with fundamental group G which are homotopically distinct.*
- (ii) *There exist closed smooth 4-manifolds M, N with fundamental group G which are stably diffeomorphic but not homotopy equivalent. More specifically, we can take $M = M(X)$ and $N = M(Y)$ for some minimal finite 2-complexes X, Y with fundamental group G .*

The proof will be broken into the following sequence of lemmas. In what follows, we will fix identifications $Q_8 = \langle x, y \mid x^2 y^{-2}, yxy^{-1}x \rangle$ and $(\mathbb{Z}/p)^3 = \langle a, b, c \mid a^p, b^p, c^p, [a, b], [a, c], [b, c] \rangle$.

Lemma 9.6. *Let $G = Q_8 \times (\mathbb{Z}/p)^3$ where p is a prime such that $p \equiv 1 \pmod{4}$. Then:*

- (i) *For each $r \in \mathbb{Z}$ with $(r, p) = 1$, we have that*

$$\mathcal{P}_r = \langle A, B, C \mid A^{2p} B^{-2p}, BAB^{-1} A^{2p-1}, C^p, [A, B^{p-1}], [A, C^r], [B, C] \rangle$$

is a presentation for G . Furthermore, we can identify $A = xa$, $B = yb$ and $C = c$.

- (ii) *G satisfies the conditions of Theorem B. More specifically, G satisfies the strong minimality hypothesis, $H_2(G) \cong (\mathbb{Z}/p)^3$, $m_G = p$ and $X_r := X_{\mathcal{P}_r} \in \text{HT}_{\min}(G)$ for each r .*

Proof. (i) Let $(\mathbb{Z}/p)^2 = \langle a, b \rangle$ and $H = Q_8 \times (\mathbb{Z}/p)^2$. Let $A = xa$, $B = yb$. Then $A^p = x$, $A^{1-p} = a$, $B^p = y$ and $B^{1-p} = b$, and so $H = \langle A, B \rangle$. By combining the given presentations for Q_8 and $(\mathbb{Z}/p)^2$ together, and adding commutators, we obtain a presentation

$$\langle A, B \mid A^{2p}B^{-2p}, B^pA^pB^{-p}A^p, A^{p(p-1)}, B^{p(p-1)}, [A^{p-1}, B^{p-1}], [A^{p-1}, B^p], [A^p, B^{p-1}] \rangle$$

where we can omit the relators $[A^p, A^{p-1}]$ and $[B^p, B^{p-1}]$ since they are trivial in $F(A, B)$.

The relators $A^{2p}B^{-2p}$ and $B^pA^pB^{-p}A^p$ imply A^{4p} and B^{4p} . Since $4 \mid p-1$, this implies $A^{p(p-1)}$ and $B^{p(p-1)}$, so these two relators can be omitted.

Using $[A^{p-1}, B^{p-1}]$, we can replace $[A^{p-1}, B^p]$ with $[A^{p-1}, B]$ and $[A^p, B^{p-1}]$ with $[A, B^{p-1}]$. Either of these relators then implies $[A^{p-1}, B^{p-1}]$ and so it can be omitted. We can also use these relators to replace $B^pA^pB^{-p}A^p$ with $BAB^{-1}A^{2p-1}$. We now have a presentation:

$$\langle A, B \mid A^{2p}B^{-2p}, BAB^{-1}A^{2p-1}, [A^{p-1}, B], [A, B^{p-1}] \rangle.$$

We now claim that $[A^{p-1}, B]$ is a consequence of the other three relators, and so can be omitted. First note that $[A, B^{p-1}] = 1$ and $BAB^{-1} = A^{1-2p}$ implies that $B^{p-1} = AB^{p-1}A^{-1} = (AB^{-1}A^{-1})^{1-p} = (B^{-1}A^{2p})^{1-p}$. Since $A^{2p} = B^{2p}$ is central, we can rewrite this as $B^{p-1} = B^{p-1}A^{2p(1-p)}$ and so $A^{2p(p-1)} = 1$. Using $BAB^{-1} = A^{1-2p}$ and $A^{2p(p-1)} = 1$, we now obtain $BA^{p-1}B^{-1} = A^{(1-2p)(p-1)} = A^{p-1}A^{2p(1-p)} = A^{p-1}$, as required. Hence we have that

$$\langle A, B \mid A^{2p}B^{-2p}, BAB^{-1}A^{2p-1}, [A, B^{p-1}] \rangle$$

is a presentation for H . By adding a generator C and relators C^p , $[A, C]$ and $[B, C]$, we clearly obtain a presentation for $G = H \times \mathbb{Z}/p$. It is straightforward to see that, if $r \in \mathbb{Z}$ with $(r, p) = 1$, then replacing $[A, C]$ with $[A, C^r] = 1$ does not change the group. Thus \mathcal{P}_r presents G .

(ii) By the Künneth formula, we have that

$$H_2(G) \cong H_2(Q_8) \oplus H_2((\mathbb{Z}/p)^3) \oplus (H_1(Q_8) \otimes_{\mathbb{Z}} H_1((\mathbb{Z}/p)^3)) \cong H_2((\mathbb{Z}/p)^3) \cong (\mathbb{Z}/p)^3$$

by Lemma 9.3 and the fact that $H_2(Q_8) = 0$, and $H_1(Q_8) \cong (\mathbb{Z}/2)^2$ and $H_1((\mathbb{Z}/p)^3) \cong (\mathbb{Z}/p)^3$ have coprime orders. Since $\chi(X_r) = 1 - 3 + 6 = 4 = 1 + d(H_2(G))$, this implies both that G has the strong minimality hypothesis and that X_r is a minimal finite 2-complex. Since $H_2(G) \cong (\mathbb{Z}/p)^3$, it now follows that $m_G = p$. \square

In what follows, we will let X_1 be the reference minimal finite 2-complex for $G = Q_8 \times (\mathbb{Z}/p)^3$. To simplify the calculation of $D(G)$, we will now restrict to the case where $p \equiv 1 \pmod{8}$. We will use the formulation for the bias invariant established in Proposition 3.15.

Lemma 9.7. *Let $G = Q_8 \times (\mathbb{Z}/p)^3$ where p is a prime such that $p \equiv 1 \pmod{8}$. Then:*

(i) *The bias invariant gives a surjection*

$$\beta: \text{HT}_{\min}(G) \rightarrow B(G).$$

Furthermore, $\beta(X_r) = [r]$ for all $r \in \mathbb{Z}$ with $(r, p) = 1$.

(ii) *$D(G) = (\mathbb{Z}/p)^{\times 2} / \{\pm 1\}$ and $B(G) = (\mathbb{Z}/p)^{\times} / \pm (\mathbb{Z}/p)^{\times 2} \cong \mathbb{Z}/2$.*

Proof. (i) Using Fox differentiation (see [59, Section 3.2]), we get that

$$C_*(\tilde{X}_r) = \left(\mathbb{Z}G^6 \xrightarrow[d_2(\tilde{X}_r)]{\begin{pmatrix} \Sigma_{2p}(xa) & -\Sigma_{2p}(xa) & 0 \\ yb+x^{-1}a\Sigma_{2p-1}(xa) & 1-x^{-1}a & 0 \\ 0 & 0 & \Sigma_p(c) \\ 1-b^{-1} & (xa-1)\Sigma_{p-1}(yb) & 0 \\ 1-c^r & 0 & (xa-1)\Sigma_r(c) \\ 0 & 1-c & yb-1 \end{pmatrix}} \mathbb{Z}G^3 \xrightarrow[d_1(\tilde{X}_r)]{\begin{pmatrix} xa-1 \\ yb-1 \\ c-1 \end{pmatrix}} \mathbb{Z}G \right)$$

where, if $u \in \mathbb{Z}G$ and $m \geq 1$, we write $\Sigma_m(u) := 1 + u + \dots + u^{m-1}$. Note that the only terms involving r are the entries $1 - c^r$ and $(xa - 1)\Sigma_r(c)$ in the 5th row of $d_2(\tilde{X}_r)$.

There is a chain map $f = (f_2, f_1, f_0) : C_*(\tilde{X}_r) \rightarrow C_*(\tilde{X}_1)$ where $f_0 = \text{id}_{\mathbb{Z}G}$, $f_1 = \text{id}_{\mathbb{Z}G^3}$ and

$$f_2 = \cdot \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & \Sigma_r(c) & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} : \mathbb{Z}G^6 \rightarrow \mathbb{Z}G^6.$$

Let $\varepsilon: \mathbb{Z}G \rightarrow \mathbb{Z}$ denote the augmentation map. If $d \in M_m(\mathbb{Z}G)$ is a matrix, $\varepsilon(d) \in M_m(\mathbb{Z})$ will denote the matrix d with ε applied to each entry. We have

$$H_2(X_r) \cong \ker(\varepsilon(d_2(\tilde{X}_r)): \mathbb{Z}^6 \rightarrow \mathbb{Z}^3) = \ker \left\{ \cdot \begin{pmatrix} 2p & -2p & 0 \\ 2p & 0 & 0 \\ 0 & 0 & p \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \right\} = \left\langle \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} \right\rangle \cong \mathbb{Z}^3.$$

With respect to this identification, the induced map $(f_2)_*: H_2(X_r) \rightarrow H_2(X_r)$ is given by $(f_2)_* = \begin{pmatrix} 1 & 0 & 0 \\ 0 & r & 0 \\ 0 & 0 & 1 \end{pmatrix}$. Hence we have $\beta(X_r) = \det((f_2)_*) = [r] \in (\mathbb{Z}/p)^\times / \pm D(G)$.

(ii) We have $D(G) = \text{im}(\varphi_G: \text{Aut}(G) \rightarrow (\mathbb{Z}/p)^\times / \{\pm 1\})$. Let $q: (\mathbb{Z}/p)^\times / \{\pm 1\} \rightarrow (\mathbb{Z}/p)^\times / \pm (\mathbb{Z}/p)^{\times 2}$ denote the natural quotient map. Since $p \equiv 1 \pmod{8}$, we can write $p = 1 + 8m$ for some m . This implies that $(\mathbb{Z}/p)^\times / \{\pm 1\} \cong \mathbb{Z}/4m$ and so we can view φ_G as a map $\varphi_G: \text{Aut}(G) \rightarrow \mathbb{Z}/4m$. Since $p \equiv 1 \pmod{4}$, $-1 \in (\mathbb{Z}/p)^{\times 2}$ and so $(\mathbb{Z}/p)^\times / \pm (\mathbb{Z}/p)^{\times 2} \cong \mathbb{Z}/2$. Thus we can view q as a map $q: \mathbb{Z}/4m \rightarrow \mathbb{Z}/2$. This is reduction mod 2 since there is a unique surjection.

Since Q_8 and $(\mathbb{Z}/p)^3$ have coprime order, we have $\text{Aut}(G) \cong \text{Aut}(Q_8) \times \text{Aut}((\mathbb{Z}/p)^3)$. We will now deal with the image of $\text{Aut}(Q_8)$ and $\text{Aut}((\mathbb{Z}/p)^3)$ under φ_G separately.

By [1, Lemma IV.6.9], $\text{Aut}(Q_8) \cong S_4$. By the fact that $(S_4)^{\text{ab}} \cong \mathbb{Z}/2$ and $\mathbb{Z}/4m$ is abelian, we have that $\varphi_G(\text{Aut}(Q_8)) \cong 0$ or $\mathbb{Z}/2$, and so $\varphi_G(\text{Aut}(Q_8)) \subseteq \{0, 2m\}$. Since q is reduction mod 2, this implies that $(q \circ \varphi_G)(\text{Aut}(Q_8)) = \{0\}$ and so $\varphi_G(\text{Aut}(Q_8)) \leq (\mathbb{Z}/p)^{\times 2} / \{\pm 1\}$.

We have $\text{Aut}((\mathbb{Z}/p)^3) \cong GL_3(\mathbb{Z}/p)$. Since \mathbb{Z}/p is a field, it follows from a 1901 theorem of Dickson [15] that $SL_3(\mathbb{Z}/p)$ is the commutator subgroup of $GL_3(\mathbb{Z}/p)$, and so there is an isomorphism $GL_3(\mathbb{Z}/p)^{\text{ab}} \cong (\mathbb{Z}/p)^\times$ induced by the determinant map. Let $\theta \in \text{Aut}((\mathbb{Z}/p)^3)$ be given by $a \mapsto a, b \mapsto b, c \mapsto c^r$ for some generator $r \in (\mathbb{Z}/p)^\times$. Then $\det(\theta) = r$ and so θ generates the abelianisation. Since $\mathbb{Z}/4m$ is abelian, $\varphi_G|_{\text{Aut}((\mathbb{Z}/p)^3)}: \text{Aut}((\mathbb{Z}/p)^3) \rightarrow \mathbb{Z}/4m$ factors through the abelianisation and so $\varphi_G(\text{Aut}((\mathbb{Z}/p)^3)) = \langle \varphi_G(\theta) \rangle$ is generated by $\varphi_G(\theta)$ as a subgroup of $\mathbb{Z}/4m$.

Now, using the isomorphisms $\mathbb{Z}G_{\theta^{-1}} \cong \mathbb{Z}G$, we obtain a chain isomorphism

$$C_*(\tilde{X}_1)_{\theta^{-1}} \cong \left(\mathbb{Z}G^6 \xrightarrow{\theta_*(d_2(\tilde{X}_1))} \mathbb{Z}G^3 \xrightarrow{\theta_*(d_1(\tilde{X}_1))} \mathbb{Z}G \right)$$

$$\cdot \begin{pmatrix} \Sigma_{2p}(xa) & -\Sigma_{2p}(xa) & 0 \\ yb+x^{-1}a\Sigma_{2p-1}(xa) & 1-x^{-1}a & 0 \\ 0 & 0 & \Sigma_p(c^r) \\ 1-b^{-1} & (xa-1)\Sigma_{p-1}(yb) & 0 \\ 1-c^r & 0 & xa-1 \\ 0 & 1-c^r & yb-1 \end{pmatrix} \cdot \begin{pmatrix} xa-1 \\ yb-1 \\ c^r-1 \end{pmatrix}$$

where, for $i = 1, 2$, $\theta_*(d_i(\tilde{X}_1))$ denotes the matrix $d_i(\tilde{X}_1)$ with the induced map $\theta_*: \mathbb{Z}G \rightarrow \mathbb{Z}G$ applied to each entry. Since $r \in (\mathbb{Z}/p)^\times$ and c has order p , we have $\Sigma_p(c^r) = \Sigma_p(c)$. In particular, the top four rows of $C_*(\tilde{X}_1)_{\theta^{-1}}$ and $C_*(\tilde{X}_1)$ coincide.

There is a chain map $g = (g_2, g_1, g_0): C_*(\tilde{X}_1)_{\theta^{-1}} \rightarrow C_*(\tilde{X}_1)$ where $g_0 = \text{id}_{\mathbb{Z}G}$ and we have

$$g_1 = \cdot \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \Sigma_r(c) \end{pmatrix}: \mathbb{Z}G^3 \rightarrow \mathbb{Z}G^3, \quad g_2 = \cdot \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & \Sigma_r(c) & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & \Sigma_r(c) \end{pmatrix}: \mathbb{Z}G^6 \rightarrow \mathbb{Z}G^6.$$

As in (i), we have that $H_2(X_r) \cong \mathbb{Z}^3$ and $H_2(\mathbb{Z} \otimes_{\mathbb{Z}G} C_*(\tilde{X}_1)_{\theta^{-1}}) \cong \mathbb{Z}^3$ are generated by the bottom three copies of \mathbb{Z} . With respect to these identifications, the induced map $(g_2)_*: H_2(\mathbb{Z} \otimes_{\mathbb{Z}G} C_*(\tilde{X}_1)_{\theta^{-1}}) \rightarrow H_2(X_r)$ is given by $(g_2)_* = \begin{pmatrix} 1 & 0 & 0 \\ 0 & r & 0 \\ 0 & 0 & r \end{pmatrix}$. This gives that

$$\varphi_G(\theta) = \beta(C_*(\tilde{X}_1), C_*(\tilde{X}_1)_\theta) = \beta(C_*(\tilde{X}_1)_{\theta^{-1}}, C_*(\tilde{X}_1)) = \det((g_2)_*) = [r^2] \in (\mathbb{Z}/p)^\times / \{\pm 1\}.$$

Hence $\varphi_G(\text{Aut}((\mathbb{Z}/p)^3)) = \langle \varphi_G(\theta) \rangle = \langle r^2 \rangle = (\mathbb{Z}/p)^{\times 2} / \{\pm 1\}$, since $r \in (\mathbb{Z}/p)^\times$ was chosen to be a generator. We established previously that $\varphi_G(\text{Aut}(Q_8)) \leq (\mathbb{Z}/p)^{\times 2} / \{\pm 1\}$, and thus we have $D(G) = \text{im}(\varphi_G) = (\mathbb{Z}/p)^{\times 2} / \{\pm 1\}$ and $B(G) = (\mathbb{Z}/p)^\times / \pm (\mathbb{Z}/p)^{\times 2} \cong \mathbb{Z}/2$. \square

Proof of Theorem 9.5. To prove (i), note that Lemma 9.7 (i) implies that there is a surjection

$$\beta: \text{HT}_{\min}(G) \twoheadrightarrow B(G) \cong \mathbb{Z}/2.$$

This implies that $|\text{HT}_{\min}(G)| > 1$, as required.

To prove (ii), note that Lemma 9.6 (ii) implies that G satisfies the conditions of Theorem B. Since $m_G = p$, we have

$$B_Q(G) \cong \frac{(\mathbb{Z}/p)^\times}{\pm(\mathbb{Z}/p)^{\times 2} \cdot D(G)}.$$

It follows from Lemma 9.7 (ii) that $D(G) = (\mathbb{Z}/p)^\times / \{\pm 1\}$, and so we obtain

$$B_Q(G) \cong (\mathbb{Z}/p)^\times / \pm (\mathbb{Z}/p)^{\times 2} \cong \mathbb{Z}/2.$$

Since $\beta: \text{HT}_{\min}(G) \rightarrow B(G)$ is surjective, Proposition 6.14 implies that

$$\beta_Q: \mathcal{M}_4(G) \rightarrow B_Q(G) \cong \mathbb{Z}/2$$

is surjective, i.e. since $\beta_Q(M(X)) = q(\beta(X))$ is the image of the bias under a surjection q . This implies that $|\mathcal{M}_4(G)| \geq 2$, which completes the proof. \square

APPENDIX A. UNITARY GROUPS AND ODD-DIMENSIONAL L -THEORY

Let Λ be a ring with involution. For $n \in \mathbb{Z}$, we will define the surgery obstruction groups $L_{2n+1}^h(\Lambda)$ and $L_{2n+1}^s(\Lambda)$ for Λ a ring with involution. These are abelian groups which only depend on the value of $n \bmod 2$. In particular, it suffices to define the groups in the cases $2n+1 \equiv 1, 3 \bmod 4$. The original definition for $\Lambda = \mathbb{Z}G$ arose from analysing the obstructions to geometric surgery problems with fundamental group G (see [67, Chapter 6]).

We will need to recall some of the details of the definition of $L_{2n+1}^h(\Lambda)$ for use in Section 7a. For $r \geq 1$, let $M_r(\Lambda)$ denote the ring of $r \times r$ matrices over Λ . If $a \mapsto \bar{a}$ denotes the involution on Λ , then $\alpha^* = (\bar{\alpha}_{ji})$ denotes the conjugate transpose matrix, for any $\alpha = (\alpha_{ij}) \in M_r(\Lambda)$. If

$$\sigma = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$$

is a $2r \times 2r$ matrix in $GL_{2r}(\Lambda)$ expressed in $r \times r$ blocks, then $U_{2r}^\varepsilon(\Lambda) \subseteq GL_{2r}(\Lambda)$ is the subgroup consisting of the matrices σ with the properties

- (i) $\alpha\delta^* + (-1)^n\beta\gamma^* = I$
- (ii) $\alpha\beta^*$ and $\gamma\delta^*$ each have the form $\theta - (-1)^n\theta^*$ for some $\theta \in M_r(\Lambda)$.

Remark A1. The unitary group $U_{2r}^\varepsilon(\Lambda)$ is a subgroup of the *hermitian* unitary group $\text{Isom}_\varepsilon(H(\Lambda^r))$, which consists of the matrices $\sigma \in GL_{2r}(\Lambda)$ which preserve the $\varepsilon = (-1)^n$ -hyperbolic form $H(\Lambda^r)$ but not necessarily the quadratic refinement. In other words, the relation

$$\sigma \begin{pmatrix} 0 & I \\ \varepsilon I & 0 \end{pmatrix} \sigma^* = \begin{pmatrix} 0 & I \\ \varepsilon I & 0 \end{pmatrix}$$

holds in $GL_{2r}(\Lambda)$, and we have $U_{2r}^\varepsilon(\Lambda) \subseteq \text{Isom}_\varepsilon(H(\Lambda^r))$. If σ is given as above, the weaker assumption is equivalent to conditions (i) and

$$(ii)' \quad (\alpha\beta^*)^* + (-1)^n\alpha\beta^* = 0 \text{ and } (\gamma\delta^*)^* + (-1)^n\gamma\delta^* = 0.$$

Condition (ii)' is implied by (ii) but is a strictly weaker condition over an arbitrary ring with involution Λ .

Lemma A2. *Let Λ be a ring with involution, let $\varepsilon = (-1)^n$ for some n and let $r \geq 1$. Then $U_{2r}^\varepsilon(\Lambda) = \text{Isom}_\varepsilon(H_\varepsilon(\Lambda^r))$ provided $\widehat{H}^{n+1}(\mathbb{Z}/2; (\Lambda, +)) = 0$, where the additive group $(\Lambda, +)$ is viewed as a $\mathbb{Z}/2$ -module under the involution. Furthermore:*

- (a) *For n even, this holds for $\Lambda = \mathbb{Z}$ or $\Lambda = \mathbb{Z}/m$ for m odd.*
- (b) *For n odd, this holds when $2 \in \Lambda^\times$, hence for $\Lambda = \mathbb{Z}/m$ for m odd (but not for $\Lambda = \mathbb{Z}$).*

Proof. Let Λ be a ring with involution such that $\widehat{H}^{n+1}(\mathbb{Z}/2; (\Lambda, +)) = 0$. We note that any matrix $X \in M_r(\Lambda)$ such that $X + \varepsilon X^* = 0$ can be expressed as $X = D + V - \varepsilon V^*$, where D is diagonal with $D + \varepsilon D^* = 0$ and V is strictly upper triangular. The condition $\widehat{H}^{n+1}(\mathbb{Z}/2; (\Lambda, +)) = 0$ applied to each diagonal entry of D shows that $D = E - \varepsilon E^*$. It follows that condition (ii)' implies condition (ii) for all $\alpha, \beta, \gamma, \delta \in M_r(\Lambda)$, and hence $U_{2r}^\varepsilon(\Lambda) = \text{Isom}_\varepsilon(H_\varepsilon(\Lambda^r))$. For the examples (a) and (b), we must show that every $x \in \Lambda$ such that $x + (-1)^n \bar{x} = 0$ is of the form $x = y - (-1)^n \bar{y}$ for some $y \in \Lambda$. This is straightforward to check in each case. For example, if $\Lambda = \mathbb{Z}$ and n is even, then the statement is simply that $2x = 0$ implies $x = 0$ for all $x \in \mathbb{Z}$. \square

A(i). **The surgery obstruction groups $L_{2n+1}^h(\Lambda)$.** We will now define the group $L_{2n+1}^h(\Lambda)$ as the quotient of the stabilised unitary group by a certain subgroup generated by elementary unitary matrices.

Definition A3. A $2r \times 2r$ matrix in $U_{2r}^\varepsilon(\Lambda)$ is called *elementary unitary* if it is a product of matrices of the following forms (see the list in [44, §1] for the case $n \equiv 1 \pmod{2}$):

- (i) $\begin{pmatrix} Q & 0 \\ 0 & (Q^*)^{-1} \end{pmatrix}$ where $Q \in GL_r(\Lambda)$
- (ii) $\begin{pmatrix} I & P \\ 0 & I \end{pmatrix}$ and $\begin{pmatrix} I & 0 \\ P & I \end{pmatrix}$ where $P = A - (-1)^n A^*$ for some $A \in M_r(\Lambda)$
- (iii) $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$ where $A = \begin{pmatrix} 0 & 0 \\ 0 & I \end{pmatrix}$, $B = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, $C = \begin{pmatrix} (-1)^n & 0 \\ 0 & 0 \end{pmatrix}$ and $D = \begin{pmatrix} 0 & 0 \\ 0 & I \end{pmatrix}$.

The subgroup in $U_{2d}^\varepsilon(\Lambda)$ of elementary unitary matrices is denoted $EU_{2r}^\varepsilon(\Lambda)$.

After orthogonal stabilisation, one defines the inclusions

$$U_2^\varepsilon(\Lambda) \subset U_4^\varepsilon(\Lambda) \subset \cdots \subset U_{2r}^\varepsilon(\Lambda) \subset U_{2r+2}^\varepsilon(\Lambda) \subset \cdots$$

whose union is the *stable unitary group* $U^\varepsilon(\Lambda)$. More precisely, the inclusion map is given by

$$i: U_{2r}(\Lambda) \rightarrow U_{2r+2}(\Lambda), \quad \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \mapsto \begin{pmatrix} \alpha' & \beta' \\ \gamma' & \delta' \end{pmatrix}$$

for $\alpha, \beta, \gamma, \delta \in M_r(\Lambda)$, and where $\alpha' = \begin{pmatrix} \alpha & 0 \\ 0 & 1 \end{pmatrix}$, $\beta' = \begin{pmatrix} \beta & 0 \\ 0 & 0 \end{pmatrix}$, $\gamma' = \begin{pmatrix} \gamma & 0 \\ 0 & 0 \end{pmatrix}$, $\delta' = \begin{pmatrix} \delta & 0 \\ 0 & 1 \end{pmatrix} \in M_{r+1}(\Lambda)$.

The union of the corresponding stabilisations $EU_{2r}^\varepsilon(\Lambda) \subset EU_{2r+2}^\varepsilon(\Lambda)$ is the subgroup $EU^\varepsilon(\Lambda) \subset U^\varepsilon(\Lambda)$ generated by all elementary unitary matrices. A key result is that $EU^\varepsilon(\Lambda)$ contains the commutator subgroup of $U^\varepsilon(\Lambda)$ (see Wall [67, Chapter 6], [54, Theorem 4.2]). The abelian quotient group

$$L_{2n+1}^h(\Lambda) = U^\varepsilon(\Lambda)/EU^\varepsilon(\Lambda)$$

provides an algebraic description of the (unbased) surgery obstruction group.

A(ii). **The surgery obstruction groups $L_{2n+1}^s(\Lambda)$.** To define the *simple* obstruction groups $L_{2n+1}^s(\Lambda)$, one must take Whitehead torsion into account and work with based free modules (see [48]). For a general ring Λ with involution, we have a natural group homomorphism by composition

$$\tau: U_{2d}^\varepsilon(\Lambda) \hookrightarrow GL_{2d}(\Lambda) \rightarrow K_1(\Lambda),$$

and we define the *special unitary group* $SU_{2r}^\varepsilon(\Lambda) = \ker \tau$. This is the group of simple automorphisms of the $(-1)^n$ -hyperbolic form

$$H(\Lambda^r) = \left(\Lambda^r \oplus \Lambda^r, \begin{pmatrix} 0 & I \\ (-1)^n I & 0 \end{pmatrix} \right)$$

on a based free module, which preserve its quadratic refinement and the preferred class of bases.

The simple surgery obstruction groups are defined as above by a certain quotient group after stabilisation. In the notation of Wall [67, Chap. 6], the group $SU^\varepsilon(\Lambda)$ is the limit of the automorphism groups $SU_{2r}^\varepsilon(\Lambda)$, where $\varepsilon = (-1)^n$, under the natural inclusions

$$SU_2^\varepsilon(\Lambda) \subset SU_4^\varepsilon(\Lambda) \subset \cdots \subset SU_{2r}^\varepsilon(\Lambda) \subset SU_{2r+2}^\varepsilon(\Lambda) \subset \cdots$$

Definition A4. We say that a $2r \times 2r$ matrix in $SU_{2r}^\varepsilon(\Lambda)$ is an *elementary special unitary* matrix if it is the product of the elementary unitary matrices in Definition A3, provided that the elements of the form

$$\begin{pmatrix} Q & 0 \\ 0 & (Q^*)^{-1} \end{pmatrix} \text{ where } Q \in GL_r(\Lambda)$$

satisfy the additional condition $\tau(Q) = 0 \in K_1(\Lambda)$, so that

$$Q \in SL_r(\Lambda) = \ker(GL_r(\Lambda) \rightarrow K_1(\Lambda)).$$

The subgroup in $SU_{2d}^\varepsilon(\Lambda)$ of elementary special unitary matrices is denoted $RU_{2r}^\varepsilon(\Lambda)$.

The corresponding union $RU^\varepsilon(\Lambda) \subset SU^\varepsilon(\Lambda)$ again contains the commutator subgroup (see [67, Chapter 6], [54, Theorem 4.2]), and the abelian quotient group

$$L_{2n+1}^s(\Lambda) = SU^\varepsilon(\Lambda)/RU^\varepsilon(\Lambda)$$

provides an algebraic description of the (based) surgery obstruction groups.

There are exact sequences relating based and unbased obstruction groups (usually called the Ranicki-Rothenberg sequences [54, Theorem 5.7]):

$$(A.5) \quad \cdots \rightarrow \hat{H}^0(\mathbb{Z}/2; \tilde{K}_1(\Lambda)) \xrightarrow{\partial} L_{2n+1}^s(\Lambda) \xrightarrow{i_*} L_{2n+1}^h(\Lambda) \xrightarrow{\tau_*} \hat{H}^1(\mathbb{Z}/2; \tilde{K}_1(\Lambda)) \rightarrow \cdots$$

where the relative terms are given by the Tate cohomology groups:

$$\hat{H}^n(\mathbb{Z}/2; \tilde{K}_1(\Lambda)) = \frac{\{A \in \tilde{K}_1(\Lambda) \mid A^* = (-1)^n A\}}{\{A + (-1)^n A^* \mid A \in \tilde{K}_1(\Lambda)\}}.$$

The map $\partial([A]) = \left[\begin{pmatrix} A & 0 \\ 0 & (A^*)^{-1} \end{pmatrix} \right]$ for $A \in GL_r(\Lambda)$ is often referred to as the *hyperbolic map* and the map i_* is induced by the inclusion $i: SU^\varepsilon(\Lambda) \hookrightarrow U^\varepsilon(\Lambda)$ which satisfies $i(RU^\varepsilon(\Lambda)) \subseteq EU^\varepsilon(\Lambda)$. The map $\tau_*: L_{2n+1}^h(\Lambda) \rightarrow \hat{H}^1(\mathbb{Z}/2; \tilde{K}_1(\Lambda))$ is induced by $\tau: U^\varepsilon(\Lambda) \rightarrow K_1(\Lambda)$. For more details see Wall [67, Chapter 6], Ranicki [54, Theorem 4.2, Theorem 5.6], Lee [44, §1], or Lees [45, §6].

A(iii). Round L -groups. In Appendix B we will use the closely related functors $L_{2n+1}^X(\Lambda)$, with torsion $X \subseteq K_1(\Lambda)$ in any involution-invariant subgroup. These L -groups are more suitable for computations since they respect products of rings with involution. The case $X = \{0\}$ is denoted $L_*^S(\Lambda)$, and $L_*^K(\Lambda)$ is defined by $X = K_1(\Lambda)$. If $X \subseteq Y$ are involution-invariant subgroups of $K_1(\Lambda)$, there is a similar Ranicki-Rothenberg sequence relating them to Tate cohomology $\hat{H}^*(\mathbb{Z}/2; K_1(\Lambda))$ (see [68, 1.1], [30], or [31, §3]).

The relation between round L -groups (corresponding to subgroups $X \subseteq K_1(\Lambda)$ with $\{\pm 1\} \in X$) and the usual L -groups (corresponding to their quotients $\tilde{X} \subseteq K_1(\Lambda)/\{\pm 1\}$) is given by the isomorphism $L_{2n}^X(\Lambda) \cong L_{2n}^{\tilde{X}}(\Lambda)$, and an exact sequence

$$(A.6) \quad 0 \rightarrow \mathbb{Z}/2 \rightarrow L_{2n+1}^X(\Lambda) \rightarrow L_{2n+1}^{\tilde{X}}(\Lambda) \rightarrow 0$$

obtained by dividing out a single $\mathbb{Z}/2$ (see [30, Proposition 3.2]).

To compare the $L_*^S(\Lambda) \rightarrow L_*^{\{\pm 1\}}(\Lambda) \rightarrow L_*^S(\Lambda)$ is a two-step process. The following braid diagram will be used in the proof of Proposition B4 and combines the exact sequences (A.6) and (A.5):

$$(A.7) \quad \begin{array}{ccccccc} & & & \beta & & & \\ & \curvearrowright & & \curvearrowright & & \curvearrowright & \\ 0 & & \mathbb{Z}/2 & & \{\pm 1\} & & 0 \\ & \searrow & \nearrow & \searrow & \nearrow & \searrow & \\ & \ker \gamma & & L_{2n+1}^{\{\pm 1\}}(\Lambda) & & \text{coker } \gamma & \\ & \nearrow & \searrow & \nearrow & \searrow & \nearrow & \\ \{\pm 1\} & & L_{2n+1}^S(\Lambda) & \xrightarrow{\alpha} & L_{2n+1}^S(\Lambda) & & 0 \\ & \curvearrowright & & \curvearrowright & & \curvearrowright & \\ & & & \gamma & & & \end{array}$$

Remark A8. For integral group rings, the round L -group associated to $L_*^S(\mathbb{Z}G)$ is $L_*^X(\mathbb{Z}G)$, where $X = \{\pm g \mid g \in G\} \subset K_1(\mathbb{Z}G)$. In particular, $L_*^K(\mathbb{Z}) = L_*^{(\pm 1)}(\mathbb{Z})$ is the round version of $L_*^S(\mathbb{Z}) = L_*^h(\mathbb{Z})$, which are the surgery obstruction groups for the trivial group $G = 1$ [68, 1.4].

APPENDIX B. COMPUTATIONS OF ODD-DIMENSIONAL L -THEORY OF ABELIAN GROUPS

In order to prove Theorem 7.1, we need to justify the L -group computations used in the proof of Proposition 7.5 by computing the boundary maps

$$\text{im}(\partial: \hat{H}^0(\mathbb{Z}/2; \tilde{K}_1(\Lambda)) \rightarrow L_{2n+1}^s(\Lambda)) = \ker(L_{2n+1}^s(\Lambda) \rightarrow L_{2n+1}^h(\Lambda))$$

in the Ranicki-Rothenberg sequences for $\Lambda = \mathbb{Z}$ and $\Lambda = \mathbb{Z}/m$. These computations use the more directly computable round L -groups and the comparison sequences (A.5) and (A.6).

Recall from Section 7a that $D_{2d}(\mathbb{Z}/m)$ is a subgroup of $SU_{2d}^\varepsilon(\mathbb{Z}/m) \subset SU^\varepsilon(\mathbb{Z}/m)$. We let

$$q: D_{2d}(\mathbb{Z}/m) \cong (\mathbb{Z}/m)^\times \rightarrow (\mathbb{Z}/m)^\times / \pm (\mathbb{Z}/m)^{\times 2}$$

denote the natural surjection induced by reduction modulo squares and $\langle -1 \rangle$.

Proposition B1. *Let $n \in \mathbb{Z}$ be even, $d \geq 1$, and $m \geq 1$. Then there is a commutative diagram*

$$\begin{array}{ccccc} D_{2d}(\mathbb{Z}/m) & \hookrightarrow & SU_{2d}(\mathbb{Z}/m) & \twoheadrightarrow & SU(\mathbb{Z}/m) \\ \downarrow & & & & \downarrow \\ (\mathbb{Z}/m)^\times / \pm (\mathbb{Z}/m)^{\times 2} & \xrightarrow{\cong} & & & L_1^S(\mathbb{Z}/m) \end{array}$$

where the lower isomorphism is induced by the hyperbolic map.

The proof will follow by first analysing $\ker(L_{2n+1}^S(\Lambda) \rightarrow L_{2n+1}^K(\Lambda))$ for $\Lambda = \mathbb{Z}$ and \mathbb{Z}/m .

Lemma B2. *Let $n \in \mathbb{Z}$. Then the hyperbolic map ∂ induces an isomorphism*

$$\ker(L_{2n+1}^S(\mathbb{Z}) \rightarrow L_{2n+1}^K(\mathbb{Z})) \cong \mathbb{Z}/2 \cdot \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}.$$

Proof. This is a consequence of a result of Wall [68, Proposition 1.4.1] (see also [42, Proposition III.10 (I)]) that there are isomorphisms

$$L_1^S(\mathbb{Z}) \cong \mathbb{Z}/2 \cdot \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \quad \text{and} \quad L_3^S(\mathbb{Z}) = \mathbb{Z}/4 \cdot [\tau]$$

where $\tau = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ and $[\tau^2] = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} = \text{im}(\partial: \widehat{H}^0(\mathbb{Z}/2; \mathbb{Z}^\times) \rightarrow L_3^S(\mathbb{Z}))$. \square

Lemma B3. *Let $m \geq 1$. Then the hyperbolic map ∂ induces isomorphisms:*

- (i) $\ker(L_1^S(\mathbb{Z}/m) \rightarrow L_1^K(\mathbb{Z}/m)) \cong \mathbb{Z}/2 \cdot \left\{ \begin{bmatrix} r_i & 0 \\ 0 & r_i^{-1} \end{bmatrix} : 1 \leq i \leq t \right\} \cong (\mathbb{Z}/m)^\times / (\mathbb{Z}/m)^{\times 2}$ where the $r_i \in (\mathbb{Z}/m)^\times$ are coset representatives for $(\mathbb{Z}/m)^\times / (\mathbb{Z}/m)^{\times 2}$.
- (ii) $\ker(L_3^S(\mathbb{Z}/m) \rightarrow L_3^K(\mathbb{Z}/m)) \cong \begin{cases} \mathbb{Z}/2 \cdot \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}, & \text{if } 8 \mid m \\ 0, & \text{otherwise.} \end{cases}$

Proof. Suppose that $m = m_1 m_2$ for $m_1, m_2 \geq 2$ coprime. Since the L^S and L^K groups respect products of rings with involution [31, §3], the isomorphism of rings $\mathbb{Z}/m \cong \mathbb{Z}/m_1 \times \mathbb{Z}/m_2$ induces a commutative diagram

$$\begin{array}{ccc} (\mathbb{Z}/m)^\times / (\mathbb{Z}/m)^{\times 2} & \xrightarrow{\cong} & (\mathbb{Z}/m_1)^\times / (\mathbb{Z}/m_1)^{\times 2} \oplus (\mathbb{Z}/m_2)^\times / (\mathbb{Z}/m_2)^{\times 2} \\ \downarrow \partial^{(m)} & & \downarrow (\partial^{(m_1)}, \partial^{(m_2)}) \\ L_r^S(\mathbb{Z}/m) & \xrightarrow{\cong} & L_r^S(\mathbb{Z}/m_1) \oplus L_r^S(\mathbb{Z}/m_2) \\ \downarrow & & \downarrow \\ L_r^K(\mathbb{Z}/m) & \xrightarrow{\cong} & L_r^K(\mathbb{Z}/m_1) \oplus L_i^K(\mathbb{Z}/m_2) \end{array}$$

for $r \equiv 0, 1, 2, 3 \pmod{4}$, where $\partial^{(m)}$, $\partial^{(m_1)}$ and $\partial^{(m_2)}$ denote the hyperbolic maps in the respective cases. We may therefore assume that $m = p^k$ is a prime power.

- (i) A starting point is $k = 1$, where Wall [68, §1.2] shows that when p is an odd prime $L_r^S(\mathbb{Z}/p) = 0, \mathbb{Z}/2, \mathbb{Z}/2, 0$, for $r \equiv 0, 1, 2, 3 \pmod{4}$, $L_1^K(\mathbb{Z}/p) = L_3^K(\mathbb{Z}/p) = 0$, and $L_0^K(\mathbb{Z}/p) = L_2^K(\mathbb{Z}/p) = \mathbb{Z}/2$. For $p = 2$, we have $L_*^S(\mathbb{Z}/2) = L_*^K(\mathbb{Z}/2) = \mathbb{Z}/2$ in each dimension.
- (ii) Since $\widehat{H}^0(\mathbb{Z}/2; (\mathbb{Z}/m)^\times) \cong (\mathbb{Z}/m)^\times / (\mathbb{Z}/m)^{\times 2}$, it suffices to show that the hyperbolic map $\partial: \widehat{H}^0(\mathbb{Z}/2; (\mathbb{Z}/m)^\times) \rightarrow L_1^S(\mathbb{Z}/m)$ is injective. By exactness, this would then imply that $\widehat{H}^0(\mathbb{Z}/2; (\mathbb{Z}/m)^\times) \cong \text{im}(\partial) = \ker(L_1^S(\mathbb{Z}/m) \rightarrow L_1^K(\mathbb{Z}/m))$. This holds since the previous map in the Rothenberg sequence is zero: $L_2^K(\mathbb{Z}/m) \cong \mathbb{Z}/2$ generated by a rank two skew-hermitian form of Arf invariant 1 with hyperbolic bilinearisation, and $\det \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = 1$.
- (iii) The kernel of the hyperbolic map $\partial: \widehat{H}^0(\mathbb{Z}/2; (\mathbb{Z}/m)^\times) \rightarrow L_3^S(\mathbb{Z}/m)$ is the image of the norm map $L_0^K(\mathbb{Z}/m) \rightarrow \widehat{H}^0(\mathbb{Z}/2; (\mathbb{Z}/m)^\times)$. If $m = p^k$ is a prime power, then the reduction map $\mathbb{Z}/p^k \rightarrow \mathbb{Z}/p$ induces isomorphisms $L_*^K(\mathbb{Z}/p^k) \cong L_*^K(\mathbb{Z}/p)$ [68, §1.2] and $\widehat{H}^r(\mathbb{Z}/2; (\mathbb{Z}/p^k)^\times) \cong \widehat{H}^r(\mathbb{Z}/2; (\mathbb{Z}/p)^\times)$ for any $r \in \mathbb{Z}$. Since $L_0^K(\mathbb{Z}/p) = \mathbb{Z}/2$ for p an odd prime, the norm map is surjective and the image of the hyperbolic map is zero in this case by naturality of the Ranicki-Rothenberg sequence.

(iv) If $m = 2^k$, the map $L_0^K(\mathbb{Z}/2^k) \rightarrow \widehat{H}^0(\mathbb{Z}/2; (\mathbb{Z}/2^k)^\times)$ is zero for $k = 1$ (trivially), but injective for $k \geq 2$. In these cases, $L_0^K(\mathbb{Z}/2^k) = \mathbb{Z}/2$ is generated by the hermitian form $h := \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$ with $\det h = 3$. Since $(\mathbb{Z}/4)^\times = \langle -1 \rangle$ and $(\mathbb{Z}/2^k)^\times = \langle -1, 3 \rangle$ if $k \geq 3$, we have $\partial\langle 3 \rangle = 0$. Therefore $\ker(L_3^S(\mathbb{Z}/2^k) \rightarrow L_3^K(\mathbb{Z}/2^k)) = 0$ for $k \leq 2$, and for $k \geq 3$ we have $\ker(L_3^S(\mathbb{Z}/2^k) \rightarrow L_3^K(\mathbb{Z}/2^k)) = \mathbb{Z}/2 \cdot \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$. \square

Proof of Proposition B1. For $\Lambda = \mathbb{Z}/m$, the natural map $L_1^S(\Lambda) \rightarrow L_1^s(\Lambda)$ factors through the comparison maps $L_1^S(\Lambda) \rightarrow L_1^{\{\pm 1\}}(\Lambda)$ and $L_1^{\{\pm 1\}}(\Lambda) \rightarrow L^s(\Lambda)$, for which the maps are explicitly described above. The isomorphisms in Lemma B2 and Lemma B3 (i) give explicit representatives for the elements of $\ker(L_1^S(\Lambda) \rightarrow L_1^K(\Lambda))$ by matrices in $D_2(\Lambda)$ for $\Lambda = \mathbb{Z}$ and $\Lambda = \mathbb{Z}/m$. Now we apply the braid diagram (A.7). The map $\alpha: L_1^S(\Lambda) \rightarrow L_1^{\{\pm 1\}}(\Lambda)$ has cokernel $\mathbb{Z}/2$, hence $\text{coker}(\gamma: L_1^S(\Lambda) \rightarrow L^s(\Lambda)) = 0$ and $\ker \gamma = \mathbb{Z}/2$. To check these results, note that by Lemma B3, $L_1^S(\Lambda) \cong (\mathbb{Z}/m)^\times / (\mathbb{Z}/m)^{\times 2}$ via the hyperbolic map, and by Lemma B2 the element $\partial(\langle -1 \rangle) \in \text{im}(L_1^S(\mathbb{Z}) \rightarrow L_1^S(\Lambda))$ generates the image of $\ker \gamma$ in $L_1^S(\Lambda)$. Therefore $L_1^s(\mathbb{Z}/m)$ is the quotient of $L_1^S(\Lambda)$ by $\partial(\langle -1 \rangle)$, and the required formula follows. \square

Proposition B4. For $m \geq 1$, $L_3^s(\mathbb{Z}/m) = L_3^h(\mathbb{Z}/m) = 0$.

Proof. If m is odd or $m \equiv 2 \pmod{4}$, part (i) of the proof of Lemma B3 shows that $L_3^K(\mathbb{Z}/m) = 0$. Therefore $L_3^h(\mathbb{Z}/m) = 0$ by (A.6). If $m = 2^k m_1$, with m_1 odd and $k \geq 3$, we have $L_3^h(\mathbb{Z}/m) = L_3^K(\mathbb{Z}/2^k) = \mathbb{Z}/2$. In the exceptional case $m = 2^k$, $k \geq 3$, the calculations of Wall [68, §1] show that the reduction map $L_3^K(\mathbb{Z}) \rightarrow L_3^K(\mathbb{Z}/2^k)$ is an isomorphism. For any $m \geq 1$, we have the commutative diagram

$$\begin{array}{ccccc} L_3^K(\mathbb{Z}) & \longrightarrow & L_3^h(\mathbb{Z}) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \\ L_3^K(\mathbb{Z}/m) & \longrightarrow & L_3^h(\mathbb{Z}/m) & \longrightarrow & 0 \end{array}$$

Since $L_3^h(\mathbb{Z}) = 0$, naturality shows that $L_3^h(\mathbb{Z}/m) = 0$, for $m \geq 1$. The Ranicki-Rothenberg sequence (A.5) combined with part (iv) of the proof of Lemma B3 shows that $L_3^s(\mathbb{Z}/m) = 0$. \square

Corollary B5. The image of $D_{2d}(\mathbb{Z}/m)$ in $SU^\varepsilon(\mathbb{Z}/m)$ is contained in $RU^\varepsilon(\mathbb{Z}/m)$, for $\varepsilon = (-1)^n$ and n odd.

APPENDIX C. THE NUMERICAL FUNCTIONS $e(d, n)$ AND $s(d, n)$

We first establish some useful facts about binomial coefficients mod 2. We will adopt the convention that $\binom{a}{b} = 0$ if $a < b$ and we will use \equiv to denote equivalence mod 2. The first property is well-known but is recalled here for convenience.

Lemma C1. Let $a, b \in \mathbb{Z}_{\geq 0}$. Then $\binom{2a}{2b+1} \equiv 0$ and $\binom{2a}{2b} \equiv \binom{2a+1}{2b} \equiv \binom{2a+1}{2b+1} \equiv \binom{a}{b}$.

Proof. Firstly, $\binom{2a}{2b+1} = \frac{2a}{2b+1} \binom{2a-1}{2b}$ implies that $\binom{2a}{2b+1} \equiv (2b+1) \binom{2a}{2b+1} \equiv (2a) \binom{2a-1}{2b} \equiv 0$.

Next note that $\binom{2a+1}{2b+1} = \frac{2a+1}{2b+1} \binom{2a}{2b}$. Clearing denominators similarly implies that $\binom{2a+1}{2b+1} \equiv \binom{2a}{2b}$. Similarly, $\binom{2a+1}{2b} = \binom{2a+1}{2(a-b)+1} \equiv \binom{2a}{2(a-b)} = \binom{2a}{2b}$ by applying the result just proven.

Finally, note that $\binom{2(a+1)}{2(b+1)} = \frac{(2a+2)(2a+1)}{(2b+2)(2b+1)} \binom{2a}{2b}$ and so $\binom{2(a+1)}{2(b+1)} \equiv \frac{a+1}{b+1} \binom{2a}{2b}$ by clearing denominators. By induction, this implies that $\binom{2a}{2b} \equiv \frac{a(a-1)\cdots(a-b)}{b(b-1)\cdots 1} \binom{2(a-b)}{0} = \binom{a}{b}$. \square

The following is often known as the hockey-stick identity.

Lemma C2. Let $a, b \in \mathbb{Z}_{\geq 0}$. Then $\sum_{i=0}^n \binom{i+k}{k} = \binom{n+k+1}{k}$.

Proof. $\sum_{i=0}^n \binom{i+k}{k} = \sum_{i=0}^n \left[\binom{i+k+1}{k+1} - \binom{i+k}{k+1} \right] = \binom{n+k+1}{k+1}$. \square

We will now establish our main results concerning the parity of $e(d, n)$ and $s(d, n)$ respectively.

Proposition C3. Let $d, n \geq 2$. Then:

(i) If $n = 4k$, then

$$e(d, n) \equiv \begin{cases} 0, & \text{if } d \equiv 0, 1, 3 \pmod{4} \\ \binom{t+k}{t+1}, & \text{if } d = 4t + 2. \end{cases}$$

(ii) If $n = 4k + 1$, then

$$e(d, n) \equiv \begin{cases} \binom{t+k}{t}, & \text{if } d = 4t \text{ or } 4t + 1 \\ \binom{t+k+1}{t+1}, & \text{if } d = 4t + 2 \text{ or } 4t + 3. \end{cases}$$

(iii) If $n = 4k + 2$, then

$$e(d, n) \equiv \begin{cases} 0, & \text{if } d \equiv 1, 3 \pmod{4} \\ \binom{t+k}{t}, & \text{if } d = 4t \\ \binom{t+k+1}{t+1}, & \text{if } d = 4t + 2. \end{cases}$$

(iv) If $n = 4k + 3$, then

$$e(d, n) \equiv \begin{cases} 0, & \text{if } d = 4t \text{ or } 4t + 1 \\ \binom{t+k+1}{t+1}, & \text{if } d = 4t + 2 \text{ or } 4t + 3. \end{cases}$$

Note that, if n is even and d is odd, then $e(d, n)$ is even. Throughout the proof, we will make repeated use of Lemmas C1 and C2.

Proof. (i) $e(d, 4k) = \sum_{i=0}^{2k-1} (2k-i) \binom{d+2i-1}{d-2} \equiv \sum_{0 \leq i \leq 2k-1, i \text{ odd}} \binom{d+2i-1}{d-2} = \sum_{i=0}^{k-1} \binom{d+4i+1}{d-2}$. If d is odd, then $\binom{d+4i+1}{d-2} \equiv 0$ for all i by Lemma C1 and so $e(d, 4k) \equiv 0$. If $d = 4t$, then $\binom{d+4i+1}{d-2} = \binom{4t+4i+1}{4t-2} \equiv \binom{2t+2i}{2t-1} \equiv 0$ by Lemma C1 and so $e(d, 4k) \equiv 0$. If $d = 4t + 2$, then $e(d, 4k) \equiv \sum_{i=0}^{k-1} \binom{4t+4i+3}{4t} \equiv \sum_{i=0}^{k-1} \binom{t+i}{t} = \binom{t+k}{t}$ by Lemmas C1 and C2.

(ii) $e(d, 4k + 1) = \sum_{i=0}^{2k} (2k-i+1) \binom{d+2i-2}{d-2} \equiv \sum_{0 \leq i \leq 2k, i \text{ even}} \binom{d+2i-2}{d-2} = \sum_{i=0}^k \binom{d+4i-2}{d-2}$. If $d = 2r$, then $e(d, 4k + 1) \equiv \sum_{i=0}^k \binom{2r+4i-2}{2r-2} \equiv \sum_{i=0}^k \binom{r+2i-1}{r-1}$ by Lemma C1. If $r = 2t$, i.e. $d = 4t$, then $e(d, 4k + 1) \equiv \sum_{i=0}^k \binom{2t+2i-1}{2t-1} \equiv \sum_{i=0}^k \binom{t+i-1}{t-1} = \binom{t+k}{t}$ by Lemmas C1 and C2. If $r = 2t + 1$, i.e. $d = 4t + 2$, then $e(d, 4k + 1) \equiv \sum_{i=0}^k \binom{2t+2i+1}{2t+1} \equiv \sum_{i=0}^k \binom{t+i}{t} = \binom{t+k+1}{t+1}$ by Lemmas C1 and C2. If $d = 2r + 1$, then $e(d, 4k + 1) \equiv \sum_{i=0}^k \binom{2r+4i-1}{2r-1} \equiv \sum_{i=0}^k \binom{r+2i-1}{r-1}$ by Lemma C1. This coincides with the $d = 2r$ case, and so $e(d, 4k + 1) \equiv \binom{t+k}{t}$ for $d = 4t + 1$ and $e(d, 4k + 1) \equiv \binom{t+k+1}{t+1}$ for $d = 4t + 3$.

(iii) $e(d, 4k + 2) = \sum_{i=0}^{2k} (2k-i+1) \binom{d+2i-1}{d-2} \equiv \sum_{0 \leq i \leq 2k, i \text{ even}} \binom{d+2i-1}{d-2} = \sum_{i=0}^k \binom{d+4i-1}{d-2}$. If d is odd, then $\binom{d+4i-1}{d-2} \equiv 0$ for all i by Lemma C1 and so $e(d, 4k + 2) \equiv 0$. If $d = 4t$, then $e(d, 4k + 2) \equiv \sum_{i=0}^k \binom{4t+4i-1}{4t-2} \equiv \sum_{i=0}^k \binom{t+i-1}{t-1} = \binom{t+k}{t}$ by Lemmas C1 and C2. If $d = 4t + 2$, then $e(d, 4k + 2) \equiv \sum_{i=0}^k \binom{4t+4i+1}{4t} \equiv \sum_{i=0}^k \binom{t+i}{t} = \binom{t+k+1}{t+1}$ by Lemmas C1 and C2.

(iv) $e(d, 4k + 3) = \sum_{i=0}^{2k+1} (2k-i+2) \binom{d+2i-2}{d-2} \equiv \sum_{0 \leq i \leq 2k, i \text{ odd}} \binom{d+2i-2}{d-2} = \sum_{i=0}^k \binom{d+4i}{d-2}$. If $d = 2r$, then $e(d, 4k + 3) \equiv \sum_{i=0}^k \binom{2r+4i}{2r-2} \equiv \sum_{i=0}^k \binom{r+2i}{r-1}$ by Lemma C1. If $r = 2t$, i.e. $d = 4t$, then $e(d, 4k + 3) \equiv \sum_{i=0}^k \binom{2t+2i}{2t-1} \equiv 0$ by Lemma C1. If $r = 2t + 1$, i.e. $d = 4t + 2$, then $e(d, 4k + 3) \equiv \sum_{i=0}^k \binom{2t+2i+1}{2t} \equiv \sum_{i=0}^k \binom{t+i}{t} = \binom{t+k+1}{t+1}$ by Lemmas C1 and C2. If $d = 2r + 1$, then $e(d, 4k + 3) \equiv \sum_{i=0}^k \binom{2r+4i+1}{2r-1} \equiv \sum_{i=0}^k \binom{r+2i}{r-1}$ by Lemma C1. This coincides with the $d = 2r$ case, and so $e(d, 4k + 3) \equiv 0$ for $d = 4t + 1$ and $e(d, 4k + 3) \equiv \binom{t+k+1}{t+1}$ for $d = 4t + 3$. \square

The parity of $s(d, n)$ can be evaluated in terms of a single binomial coefficient without splitting into cases for n and $d \pmod{4}$. However, in order to make the results more easily comparable to Proposition C3, we will also state the result in these terms.

Proposition C4. *Let $d, n \geq 2$. Then $s(d, n) \equiv 1 + \binom{d+n}{d}$. In particular, we have that:*

(i) If $n = 4k$, then

$$s(d, n) \equiv 1 + \binom{t+k}{t}, \quad \text{if } d = 4t, 4t + 1, 4t + 2 \text{ or } 4t + 3.$$

(ii) If $n = 4k + 1$, then

$$s(d, n) \equiv \begin{cases} 1 + \binom{t+k}{t}, & \text{if } d = 4t \text{ or } 4t + 2 \\ 1, & \text{if } d = 4t + 1 \text{ or } 4t + 3. \end{cases}$$

(iii) If $n = 4k + 2$, then

$$s(d, n) \equiv \begin{cases} 1 + \binom{t+k}{t}, & \text{if } d = 4t \text{ or } 4t + 1 \\ 1, & \text{if } d = 4t + 2 \text{ or } 4t + 3. \end{cases}$$

(iv) If $n = 4k + 3$, then

$$s(d, n) \equiv \begin{cases} 1 + \binom{t+k}{t}, & \text{if } d = 4t \text{ or } 4t + 2 \\ 1, & \text{if } d = 4t + 1 \text{ or } 4t + 3. \end{cases}$$

It follows from Lemma C1 that, if n is odd and d is odd, then $s(d, n)$ is odd.

Proof. $s(d, n) = \sum_{i=0}^{n-1} (-1)^{n+i+1} \binom{d+i}{d-1} \equiv 1 + \sum_{i=-1}^{n-1} \binom{d+i}{d-1} \equiv 1 + \binom{n+d}{d}$, by Lemma C2. To evaluate $s(d, n)$ in the various values of $n, d \bmod 4$, we repeatedly apply Lemma C1. \square

REFERENCES

- [1] A. Adem and R. J. Milgram, *Cohomology of finite groups*, second ed., Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 309, Springer-Verlag, Berlin, 2004.
- [2] S. Akbulut and R. Kirby, *A potential smooth counterexample in dimension 4 to the Poincaré conjecture, the Schoenflies conjecture, and the Andrews-Curtis conjecture*, *Topology* **24** (1985), 375–390.
- [3] J. J. Andrews and M. L. Curtis, *Free groups and handlebodies*, *Proc. Amer. Math. Soc.* **16** (1965), 192–195.
- [4] A. Bak, V. Petrov, and G. Tang, *Stability for quadratic K_1* , *K-Theory* **30** (2003), 1–11, Special issue in honor of Hyman Bass on his seventieth birthday. Part I.
- [5] H. Bass, *Unitary algebraic K-theory*, Algebraic K-theory, III: Hermitian K-theory and geometric applications (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972), Lecture Notes in Math., Vol. 343, Springer, Berlin-New York, 1973, pp. 57–265.
- [6] S. Bauer, *The homotopy type of a 4-manifold with finite fundamental group*, Algebraic topology and transformation groups (Göttingen, 1987), Lecture Notes in Math., vol. 1361, Springer, Berlin, 1988, pp. 1–6.
- [7] G. E. Bredon, *Introduction to compact transformation groups*, Academic Press, New York, 1972, Pure and Applied Mathematics, Vol. 46.
- [8] K. S. Brown, *Cohomology of groups*, Graduate Texts in Mathematics, vol. 87, Springer-Verlag, New York, 1994, Corrected reprint of the 1982 original.
- [9] W. Browning, *Finite CW-complexes of cohomological dimension 2 with finite abelian π_1* , ETH preprint (23 pages), 1979.
- [10] ———, *Homotopy types of certain finite CW-complexes with finite fundamental group*, Ph.D. thesis, Cornell University, 1979.
- [11] A. Conway, D. Crowley, M. Powell, and J. Sixt, *Stably diffeomorphic manifolds and modified surgery obstructions*, 2022, *Mém. Soc. Math. Fr.*, to appear. arXiv:2109.05632.
- [12] A. Conway, D. Crowley, M. Powell, and J. Sixt, *Simply connected manifolds with large homotopy stable classes*, *J. Aust. Math. Soc.* **115** (2023), 172–203.
- [13] C. W. Curtis and I. Reiner, *Methods of representation theory. Vol. I*, Pure and Applied Mathematics, John Wiley & Sons, Inc., New York, 1981, With applications to finite groups and orders, A Wiley-Interscience Publication.
- [14] ———, *Methods of representation theory. Vol. II*, Pure and Applied Mathematics (New York), John Wiley & Sons, Inc., New York, 1987, With applications to finite groups and orders, A Wiley-Interscience Publication.
- [15] L. E. Dickson, *Linear groups: With an exposition of the Galois field theory*, Dover Publications, Inc., New York, 1901.
- [16] S. K. Donaldson, *An application of gauge theory to four-dimensional topology*, *J. Differential Geom.* **18** (1983), 279–315.
- [17] Y. A. Drozd and A. I. Plakosh, *Cohomologies of finite abelian groups*, *Algebra Discrete Math.* **24** (2017), 144–157.
- [18] M. N. Dyer, *Homotopy classification of (π, m) -complexes*, *J. Pure Appl. Algebra* **7** (1976), 249–282.
- [19] ———, *On the essential height of homotopy trees with finite fundamental group*, *Compositio Math.* **36** (1978), 209–224.
- [20] M. Freedman, V. Krushkal, and T. Lidman, *Spineless 5-manifolds and the deformation conjecture*, 2024, arXiv:2401.03498.
- [21] R. E. Gompf, *Killing the Akbulut-Kirby 4-sphere, with relevance to the Andrews-Curtis and Schoenflies problems*, *Topology* **30** (1991), 97–115.

- [22] I. Hambleton, *Two remarks on Wall's D_2 problem*, Math. Proc. Cambridge Philos. Soc. **167** (2019), 361–368.
- [23] ———, *A stability range for topological 4-manifolds*, Trans. Amer. Math. Soc. **376** (2023), 8769–8793.
- [24] I. Hambleton and A. Hildum, *Topological 4-manifolds with right-angled Artin fundamental groups*, J. Topol. Anal. **11** (2019), 777–821.
- [25] ———, *Topological 4-manifolds with right-angled Artin fundamental groups: corrigendum*, Journal of Topology and Analysis (2024), doi: 10.1142/S1793525324500249.
- [26] I. Hambleton and M. Kreck, *On the classification of topological 4-manifolds with finite fundamental group*, Math. Ann. **280** (1988), 85–104.
- [27] ———, *Cancellation results for 2-complexes and 4-manifolds and some applications*, Two-dimensional homotopy and combinatorial group theory, London Math. Soc. Lecture Note Ser., vol. 197, Cambridge Univ. Press, Cambridge, 1993, pp. 281–308.
- [28] ———, *On the classification of topological 4-manifolds with finite fundamental group: corrigendum*, Math. Ann. **372** (2018), 527–530.
- [29] I. Hambleton, M. Kreck, and P. Teichner, *Topological 4-manifolds with geometrically two-dimensional fundamental groups*, J. Topol. Anal. **1** (2009), 123–151.
- [30] I. Hambleton, A. Ranicki, and L. R. Taylor, *Round L -theory*, J. Pure Appl. Algebra **47** (1987), 131–154.
- [31] I. Hambleton and L. R. Taylor, *A guide to the calculation of the surgery obstruction groups for finite groups*, Surveys on surgery theory, Vol. 1, Ann. of Math. Stud., vol. 145, Princeton Univ. Press, Princeton, NJ, 2000, pp. 225–274.
- [32] J. Harlander, *Some aspects of efficiency*, Groups—Korea '98 (Pusan), de Gruyter, Berlin, 2000, pp. 165–180.
- [33] F. E. A. Johnson, *Stable modules and the $D(2)$ -problem*, London Mathematical Society Lecture Note Series, vol. 301, Cambridge University Press, Cambridge, 2003.
- [34] D. Kasprowski, M. Land, M. Powell, and P. Teichner, *Stable classification of 4-manifolds with 3-manifold fundamental groups*, J. Topol. **10** (2017), 827–881.
- [35] D. Kasprowski, J. Nicholson, and B. Ruppik, *Homotopy classification of 4-manifolds whose fundamental group is dihedral*, Algebr. Geom. Topol. **22** (2022), 2915–2949.
- [36] D. Kasprowski, M. Powell, and A. Ray, *Counterexamples in 4-manifold topology*, EMS Surv. Math. Sci. **9** (2022), 193–249.
- [37] D. Kasprowski, M. Powell, and B. Ruppik, *Homotopy classification of 4-manifolds with finite abelian 2-generator fundamental groups*, Math. Proc. Cambridge Philos. Soc. **177** (2024), 263–283.
- [38] D. Kasprowski, M. Powell, and P. Teichner, *Algebraic criteria for stable diffeomorphism of spin 4-manifolds*, 2020, Mem. Amer. Math. Soc., to appear. arXiv:2006.06127.
- [39] D. Kasprowski and P. Teichner, \mathbb{CP}^2 -stable classification of 4-manifolds with finite fundamental group, Pacific J. Math. **310** (2021), 355–373.
- [40] M.-A. Knus, *Quadratic and Hermitian forms over rings*, Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 294, Springer-Verlag, Berlin, 1991, With a foreword by I. Bertuccioni.
- [41] M. Kreck, *Surgery and duality*, Ann. of Math. (2) **149** (1999), 707–754.
- [42] M. Kreck and J. A. Schafer, *Classification and stable classification of manifolds: some examples*, Comment. Math. Helv. **59** (1984), 12–38.
- [43] M. P. Latiolais, *Homotopy and homology classification of 2-complexes*, Two-dimensional homotopy and combinatorial group theory, London Math. Soc. Lecture Note Ser., vol. 197, Cambridge Univ. Press, Cambridge, 1993, pp. 97–124.
- [44] R. Lee, *Computation of Wall groups*, Topology **10** (1971), 149–176.
- [45] J. A. Lees, *The surgery obstruction groups of $C. T. C. Wall$* , Advances in Math. **11** (1973), 113–156.
- [46] P. A. Linnell, *Minimal free resolutions and (G, n) -complexes for finite abelian groups*, Proc. London Math. Soc. (3) **66** (1993), 303–326.
- [47] W. Metzler, *Über den Homotopietyp zweidimensionaler CW-Komplexe und Elementartransformationen bei Darstellungen von Gruppen durch Erzeugende und definierende Relationen*, J. Reine Angew. Math. **285** (1976), 7–23.
- [48] J. Milnor, *Whitehead torsion*, Bull. Amer. Math. Soc. **72** (1966), 358–426.
- [49] C. Nagy, J. Nicholson, and M. Powell, *Generalised doubles and simple homotopy types of high dimensional manifolds*, 2024, arXiv:2409.03082.
- [50] J. Nicholson, *On CW-complexes over groups with periodic cohomology*, Trans. Amer. Math. Soc. **374** (2021), 6531–6557.
- [51] ———, *Stably free modules and the unstable classification of 2-complexes*, 2023, Math. Ann., to appear. arXiv:2108.02220.
- [52] ———, *Cancellation for (G, n) -complexes and the Swan finiteness obstruction*, Int. Math. Res. Not. IMRN (2024), 11832–11864.
- [53] ———, *Projective modules and the homotopy classification of (G, n) -complexes*, Algebr. Geom. Topol. **24** (2024), 2245–2284.
- [54] A. A. Ranicki, *Algebraic L -theory. I. Foundations*, Proc. London Math. Soc. (3) **27** (1973), 101–125.
- [55] J. A. Schafer, *Fiber squares and complexes*, J. Pure Appl. Algebra **106** (1996), 153–171.

- [56] W. Scharlau, *Quadratic and Hermitian forms*, Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 270, Springer-Verlag, Berlin, 1985.
- [57] A. J. Sieradski and M. N. Dyer, *Distinguishing arithmetic for certain stably isomorphic modules*, J. Pure Appl. Algebra **15** (1979), 199–217.
- [58] A. J. Sieradski, *A semigroup of simple homotopy types*, Math. Z. **153** (1977), 135–148.
- [59] ———, *Algebraic topology for two-dimensional complexes*, Two-dimensional homotopy and combinatorial group theory, London Math. Soc. Lecture Note Ser., vol. 197, Cambridge Univ. Press, Cambridge, 1993, pp. 51–96.
- [60] R. G. Swan, *Minimal resolutions for finite groups*, Topology **4** (1965), 193–208.
- [61] ———, *Groups with no odd dimensional cohomology*, J. Algebra **17** (1971), 401–403.
- [62] M. J. Taylor, *Locally free classgroups of groups of prime power order*, J. Algebra **50** (1978), 463–487.
- [63] P. Teichner, *Topological 4-manifolds with finite fundamental group*, Ph.D. thesis, University of Mainz, Germany, 1992, Shaker Verlag, ISBN 3-86111-182-9.
- [64] C. T. C. Wall, *On simply-connected 4-manifolds*, J. London Math. Soc. **39** (1964), 141–149.
- [65] ———, *Finiteness conditions for CW-complexes*, Ann. of Math. (2) **81** (1965), 56–69.
- [66] ———, *Classification problems in differential topology. IV. Thickenings*, Topology **5** (1966), 73–94.
- [67] ———, *Surgery on compact manifolds*, London Mathematical Society Monographs, No. 1, Academic Press, London-New York, 1970.
- [68] ———, *Classification of Hermitian Forms. VI. Group rings*, Ann. of Math. (2) **103** (1976), 1–80.
- [69] P. J. Webb, *The minimal relation modules of a finite abelian group*, J. Pure Appl. Algebra **21** (1981), 205–232.
- [70] J. H. C. Whitehead, *Simplicial Spaces, Nuclei and m -Groups*, Proc. London Math. Soc. (2) **45** (1939), 243–327.
- [71] J. S. Williams, *Free presentations and relation modules of finite groups*, J. Pure Appl. Algebra **3** (1973), 203–217.

DEPARTMENT OF MATHEMATICS & STATISTICS, MCMASTER UNIVERSITY L8S 4K1, ONTARIO, CANADA
 Email address: hambleton@mcmaster.ca

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF GLASGOW, UNITED KINGDOM
 Email address: john.nicholson@glasgow.ac.uk