

# Matrices for general linear transformations

Goal: To associate a matrices to linear transformations between finite-dimensional vector spaces.

The process: Suppose that  $T : V \rightarrow W$  is a linear transformation from an  $n$ -dimensional vector space  $V$  to an  $m$ -dimensional vector space  $W$ .

- 1 Fix a basis  $B$  for  $V$  and  $B'$  for  $W$ .
- 2 Construct an  $m \times n$  matrix  $A$  such that

$$A[x]_B = [T(x)]_{B'}$$

$A$  is called the matrix for  $T$  with respect to  $B$  and  $B'$  and we will denote it by  $[T]_{B',B}$ .

- 3 Suppose  $B = \{u_1, u_2, \dots, u_n\}$ . Form  $A$  with column vectors  $[T(u_1)]_{B'}, [T(u_2)]_{B'}, \dots, [T(u_n)]_{B'}$ .

# Notational issues

Fix  $T : V \rightarrow W$  is a linear transformation from an  $n$ -dimensional vector space  $V$  to an  $m$ -dimensional vector space  $W$ .

- 1 Notice that

$$[T]_{B',B}[x]_B = [T(x)]_{B'}$$

This is the real reason for writing the subscripts in this order.

- 2 If  $T$  is a linear operator i.e. if  $V = W$ , then we will write  $[T]_B$  for  $[T]_{B,B}$ .

# Change of basis

## The problem

Suppose we are given two bases

$$B = \{u_1, u_2, \dots, u_n\} \text{ and } B' = \{u'_1, u'_2, \dots, u'_n\}$$

for an  $n$ -dimensional vector space  $V$ ; how are  $B$  and  $B'$  related?

## The solution

Let  $P$  be the  $n \times n$  matrix given by

$$P = ([u'_1]_B, [u'_2]_B, \dots, [u'_n]_B)$$

Then  $[v]_B = P[v]_{B'}$  for all  $v \in V$ .  $P$  is called the transition matrix from  $B'$  to  $B$

# Change of basis, cont'd

## Theorem

*If  $P$  is the transition matrix from  $B'$  to  $B$  and  $Q$  is the transition matrix from  $B$  to  $B'$  then  $Q = P^{-1}$ .*

## Change of basis, cont'd

- So the matrix representing the identity transformation,  $I : V \rightarrow V$ , with respect to  $B$  and  $B'$  is just the change of basis matrix  $P$ .

### Theorem

*If  $T : V \rightarrow V$  is a linear operator on a finite-dimensional vector space  $V$  and  $B$  and  $B'$  are two bases for  $V$  then*

$$[T]_{B'} = P^{-1}[T]_B P$$

*where  $P$  is the change of basis matrix from  $B'$  to  $B$ .*

# Composition and inverse

## Theorem

If  $T_1 : U \rightarrow V$  and  $T_2 : V \rightarrow W$  are linear transformations between finite-dimensional vector spaces and  $B, B'$  and  $B''$  are bases for  $U, V$  and  $W$  respectively then

$$[T_2 \circ T_1]_{B'',B} = [T_2]_{B'',B'} [T_1]_{B',B}$$

## Theorem

If  $T : V \rightarrow V$  is a linear operator and  $B$  is a basis for  $V$  then  $T$  is one-to-one iff  $[T]_B$  is invertible. If  $T$  is one-to-one then

$$[T^{-1}]_B = [T]_B^{-1}$$