

Lecture 30

6.B Linear functionals and a worked out Example

Observation: F is an F -vector space. So, for any vector space V , $\mathcal{L}(V, F)$ makes sense

Defⁿ A linear functional on V is any element of $\mathcal{L}(V, F)$

Ex Let $V = F^3$ and define

$$\varphi: V \rightarrow F \text{ by } \varphi((x, y, z)) = 3x + 12y - 2z$$

Note: If $\langle -, - \rangle$ denotes the standard Euclidean inner product on F^3 , then

$$\varphi((x, y, z)) = 3x + 12y - 2z = \langle (x, y, z), \underline{(3, 12, -2)} \rangle$$

Ex Let $V = \mathcal{P}_2(\mathbb{R})$ and define

$$\psi: V \rightarrow \mathbb{R} \quad \text{by} \quad \psi(p(x)) = \int_0^1 p(x) dx$$

$$\begin{aligned} \text{So } \psi(a_0 + a_1x + a_2x^2) &= \int_0^1 a_0 + a_1x + a_2x^2 dx \\ &= a_0x + \frac{a_1x^2}{2} + \frac{a_2x^3}{3} \Big|_0^1 = a_0 + \frac{a_1}{2} + \frac{a_2}{3} \end{aligned}$$

Note Can define an inner product $\langle -, - \rangle$ on $\mathcal{P}_2(\mathbb{R})$
b,

$$\langle p(x), q(x) \rangle = \int_0^1 p(x)q(x) dx$$

$$\text{So } \psi(p(x)) = \langle p(x), 1 \rangle = \int_0^1 p(x) \cdot 1 dx$$

(Riesz Representation Theorem) Let V be a fin. dim. inner product space. If ϕ is a linear functional, i.e. $\phi \in \mathcal{L}(V, F)$, then there exists a unique $u \in V$ such that

$$\phi(v) = \langle v, u \rangle \text{ for all } v \in V$$

exists by
G.S.

Proof Let e_1, \dots, e_n be an orthonormal basis of V .

Then

$$v = \langle v, e_1 \rangle e_1 + \langle v, e_2 \rangle e_2 + \dots + \langle v, e_n \rangle e_n \text{ for all } v \in V$$

$$\begin{aligned} \text{So } \phi(v) &= \phi(\langle v, e_1 \rangle e_1 + \dots + \langle v, e_n \rangle e_n) && \downarrow \phi \text{ is linear} \\ &= \langle v, e_1 \rangle \phi(e_1) + \dots + \langle v, e_n \rangle \phi(e_n) \\ &= \langle v, \overline{\phi(e_1)} e_1 \rangle + \dots + \langle v, \overline{\phi(e_n)} e_n \rangle && \downarrow \text{prop of inner prod} \\ &= \langle v, \overline{\phi(e_1)} e_1 + \dots + \overline{\phi(e_n)} e_n \rangle && \downarrow \text{"} \end{aligned}$$

Set $u = \overline{\phi(e_1)} e_1 + \dots + \overline{\phi(e_n)} e_n$. So

$$\phi(v) = \langle v, u \rangle \text{ for all } v \in V$$

To show u is unique, suppose

$$\phi(v) = \langle v, u_1 \rangle = \langle v, u_2 \rangle \text{ for some } u_1, u_2$$

$$\begin{aligned} \text{So } 0 &= \langle v, u_1 \rangle - \langle v, u_2 \rangle \\ &= \langle v, u_1 - u_2 \rangle \quad \text{for all } v \in V \end{aligned}$$

In particular, for $v = u_1 - u_2$,

$$0 = \langle u_1 - u_2, u_1 - u_2 \rangle \Leftrightarrow u_1 - u_2 = 0 \Leftrightarrow u_1 = u_2$$



Very nice result! Proof gives procedure to find u

① Find an orthonormal basis e_1, \dots, e_n for V (via G-S)

② Compute $\overline{\phi(e_1)}, \dots, \overline{\phi(e_n)}$

③ Return $u = \overline{\phi(e_1)}e_1 + \overline{\phi(e_2)}e_2 + \dots + \overline{\phi(e_n)}e_n$

Worked out Example

Show that there is a polynomial $g(x) \in \mathcal{P}_1(\mathbb{R})$ such that

$$\int_0^1 p(x)g(x) dx = \int_0^1 p(x) \sin(\pi x) dx \text{ for all } p(x) \in \mathcal{P}_1(\mathbb{R})$$

(the power of Riesz Representation!)

Note: • $\mathcal{P}_1(\mathbb{R})$ is an inner product space with

$$\langle p(x), g(x) \rangle = \int_0^1 p(x)g(x) dx$$

• The map $\varphi: \mathcal{P}_1(\mathbb{R}) \rightarrow \mathbb{R}$ given by

$$\varphi(p(x)) = \int_0^1 p(x) \sin(\pi x) dx \text{ is a linear functional}$$

hypotheses
of
Riesz
Rep. Thm
are met

Riesz Rep Thm \Rightarrow exists some $g(x) \in \mathcal{P}_1(\mathbb{R})$ such that

$$\int_0^1 p(x) \sin(\pi x) dx = \varphi(p(x)) \stackrel{\uparrow}{=} \langle p(x), g(x) \rangle = \int_0^1 p(x)g(x) dx$$

Riesz Repres

We use the procedure to find $g(x)$

Step 1 Find an orthonormal basis of $\mathcal{P}_1(\mathbb{R})$

A basis for $\mathcal{P}_1(\mathbb{R})$ is $\{1, x\}$. Now apply Gram-Schmidt. Let $v_1 = 1$, $v_2 = x$

$$\bullet e_1 = \frac{v_1}{\|v_1\|} = 1$$

$$\bullet e_2 = \frac{v_2 - \langle v_2, e_1 \rangle e_1}{\|v_2 - \langle v_2, e_1 \rangle e_1\|}$$

$$\text{Calculations: } \langle v_2, e_1 \rangle = \langle x, 1 \rangle = \int_0^1 x \, dx = \frac{x^2}{2} \Big|_0^1 = \frac{1}{2}$$

$$\text{So } v_2 - \langle v_2, e_1 \rangle e_1 = x - \frac{1}{2} \cdot 1 = x - \frac{1}{2}$$

$$\|v_2 - \langle v_2, e_1 \rangle e_1\| = \|x - \frac{1}{2}\| = \sqrt{\langle x - \frac{1}{2}, x - \frac{1}{2} \rangle}$$

$$= \sqrt{\int_0^1 \left(x - \frac{1}{2}\right)^2 dx} = \sqrt{\int_0^1 x^2 - x + \frac{1}{4} dx}$$

$$= \sqrt{\frac{x^3}{3} - \frac{x^2}{2} + \frac{1}{4}x \Big|_0^1} = \frac{1}{\sqrt{12}}$$

$$\text{So } e_2 = \frac{x - \frac{1}{2}}{\frac{1}{\sqrt{12}}} = \sqrt{12} \left(x - \frac{1}{2}\right)$$

Orthonormal basis of $\mathcal{P}_1(\mathbb{R})$ is $e_1 = 1$, $e_2 = \sqrt{12} \left(x - \frac{1}{2}\right)$

$$e_1 = 1$$

$$e_2 = \sqrt{12} \left(x - \frac{1}{2}\right) \leftarrow \text{orthonormal basis for } \mathcal{P}_1(\mathbb{R})$$

Step 2 Compute $\overline{\varphi(e_1)}, \overline{\varphi(e_2)}$
Note $F = \mathbb{R}$, $\overline{\varphi(e_1)} = \varphi(e_1)$, $\overline{\varphi(e_2)} = \varphi(e_2)$

$$\varphi(1) = \int_0^1 \sin(\pi x) dx = -\frac{\cos(\pi x)}{\pi} \Big|_0^1$$

$$= -\frac{\cos(\pi)}{\pi} + \frac{\cos(0)}{\pi} = \boxed{\frac{2}{\pi}}$$

$$\begin{aligned}
\phi(\sqrt{12}(x - \frac{1}{2})) &= \int_0^1 \sqrt{12}(x - \frac{1}{2}) \sin(\pi x) dx \\
&= \sqrt{12} \int_0^1 x \sin(\pi x) dx - \frac{\sqrt{12}}{2} \int_0^1 \sin(\pi x) dx \\
&= \sqrt{12} \left[-x \frac{\cos(\pi x)}{\pi} + \frac{\sin \pi x}{\pi} \Big|_0^1 \right] - \frac{\sqrt{12}}{2} \left[-\frac{\cos \pi x}{\pi} \Big|_0^1 \right] \\
&= \sqrt{12} \left[\frac{-\cos(\pi) + 0}{\pi} \right] - \frac{\sqrt{12}}{2} \cdot \frac{2}{\pi} \\
&= \sqrt{12} \left(\frac{-1(-1)}{\pi} \right) - \frac{\sqrt{12}}{\pi} = 0
\end{aligned}$$

So $\overline{\phi(e_2)} = \phi(e_2) = 0$

Step 3 Return $\cdot u = \overline{\phi(e_1)}e_1 + \overline{\phi(e_2)}e_2$

$$u = \frac{2}{\pi} \cdot 1 + 0 \cdot e_2 = \frac{2}{\pi}$$

Conclusion: For all $p(x) \in \mathcal{P}_1(\mathbb{R})$

$$\int_0^1 p(x) \sin(\pi x) dx = \int_0^1 p(x) \cdot \frac{2}{\pi} dx$$

Key ideas * Linear functionals
* Riesz Representation

